

Proceedings of the Iowa Academy of Science

Volume 94 | Number

Article 7

1987

Life History of *Elsinoe panici*

A. W. Gabel
Iowa State University

L. H. Tiffany
Iowa State University

Let us know how access to this document benefits you

Copyright ©1987 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Gabel, A. W. and Tiffany, L. H. (1987) "Life History of *Elsinoe panici*," *Proceedings of the Iowa Academy of Science*, 94(4), 121-127.

Available at: <https://scholarworks.uni.edu/pias/vol94/iss4/7>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

Life History of *Elsinoë panici*¹

A.W. GABEL² and L.H. TIFFANY

Department of Botany, Iowa State University, Ames, Iowa 50011

Elsinoë panici Tiffany and Mathre is widespread on native *Panicum virgatum* L. in Iowa. In June *Sphaceloma* (imperfect stage) conidia develop from acervuli on overwintered stromata of wind-dispersed leaf litter and culms. Germ tubes penetrate between epidermal cells of young leaves, and 2 wk later tiny yellowish, subepidermal stromata are visible. Secondary conidia develop from acervuli on these stromata within 2 wk and are spread to adjacent leaves. Stromata become black, raised, shiny, ovoid-elongate patches, composed of thick-walled pseudoparenchymatous cells. Chlorosis or necrosis does not develop around the stromata. Stromata are visible on all leaves throughout the season. Acervuli, appearing as white spots in black stromata, open by irregular tears in the epidermis. Ascstromata develop on lower leaves infrequently from late July into September. They are tan-bronze, raised, ovoid-elongate, softer, and more loosely structured than stromata. Asci mature at different times. Ascospores are discharged from the bitunicate asci and germinate directly by budding or by germ tubes.

INDEX DESCRIPTORS: *Elsinoë panici*, *Panicum virgatum*, life cycle.

In 1961, a leaf-spotting pathogen on *P. virgatum* was described as *E. panici* by Tiffany and Mathre. The genus *Elsinoë* exhibits a simple, undifferentiated, innate, intra- or sub-epidermal ascostroma. Members of the genus produce hyaline to yellowish ascospores with three transverse septa and longitudinal septa in some cells in globose to elliptical bitunicate asci (Luttrell, 1973).

The only other reports of *Elsinoë*, or of the closely related genus *Myriangiium*, on a member of the Gramineae are of *M. bambusae* Rick on *Phyllostachys puberula* Munro in China (Tai, 1931) and *E. sacchari* Lo on *Saccharum officinarum* L. from Taiwan (Lo, 1964) and Florida (Todd, 1960). All other known hosts for *Elsinoë* spp. are herbaceous or woody dicots. Most *Elsinoë* spp. are tropical or subtropical. However, temperate-zone diseases caused by *Elsinoë* spp. do occur (Brook, 1973; Jenkins, 1932; Jenkins et al., 1946; Jones, 1924; and Massey and Jenkins, 1935). Most studies have concerned life histories, symptoms, inoculations, and growth in culture with economically important hosts. *E. panici* has never been observed on other hosts, including other species of *Panicum*.

There has been recent interest in *P. virgatum* because of its use as summer and fall forage. The purpose of this study was to elucidate a life cycle for *E. panici* and to provide background information for assessing disease losses and selecting plant material in breeding programs.

MATERIALS AND METHODS

Field Observations

Leaf collections were made regularly from August 1981 to December 1984 at various sites (Fig. 1). Diseased leaves were catalogued and either pressed or killed and fixed. Leaves with ascstromata (stromata that consistently developed asci and ascospores) and stromata (stromata that did not usually develop asci, but often formed acervuli) were placed in moist chambers at 4, 15, and 25 C for 3-5 days and observed at regular intervals for development.

Overwintering and spring development were observed by placing 100 leaves, with ascstromata or stromata from fall 1983 collections, in nylon mesh bags attached to stakes anchored on the soil surface in a reconstructed prairie near Ames, Iowa. Samples were removed the following March, May, and June, examined directly for conidia and/or ascospores, and then placed in moist chambers at 4, 15, and 25 C for 3-5 days and observed for conidia or ascospores.

Ascospore maturity was followed from August to October 1984 by removing 15-20 ascstromata from the lowest or next lowest leaf and observing ascospores of 100 asci from each sample. Ascospores were considered mature if transverse septa were observed. This maturity information was used to determine persistence of the sexual system.

Inoculation of *Panicum virgatum*

Isolates for inoculation were obtained from green infected leaves of *P. virgatum* collected at various sites. The leaves were dipped in 95% ethyl alcohol and immersed in 0.525% NaOCl for 5 min. Small pieces of leaves were placed on potato dextrose agar (PDA) amended with an 80 mg/l solution of streptomycin sulfate. Isolates were incubated in the dark or in laboratory light at 25 C and transferred to PDA every 4 wk. Cultures were initiated by spreading 0.3 ml of a conidial suspension over surfaces of PDA plates and then holding plates for 3 wk at 25 C. Conidia were harvested from plates with deionized water, concentrations determined with a hemacytometer and used as inoculum.

Detached Leaves — Leaves of plants grown from rhizomes of previously infected plants from Kalsow State Preserve (IKP) in Pocahontas County and leaves of plants of the cultivar Blackwell (BP) grown from 1979 seed were detached, and 2 drops of a conidial suspension containing 25×10^6 spores/ml of water were placed on




Fig. 1. Sites where infected *Panicum virgatum* plants that did not develop sexual structures (●) and plants that did develop sexual structures (○) were collected from August 1981 to December 1984. Disease development was monitored each year, and sexual structures developed at Doolittle Prairie (⊙).

¹Journal Paper No. 12625 of the Iowa Agriculture and Home Economics Experiment Station, Ames, Iowa, Project No. 0110.

²Present address: Division of Science and Mathematics, Black Hills State College, Spearfish, SD 57783.

each leaf. Leaves were placed in humidity chambers and incubated in laboratory light at 25 C. Half of the leaves from each source were inoculated on the adaxial surface and the remaining leaves were inoculated on the abaxial surface. Samples were removed periodically up to 15 days after inoculation, cleared in saturated chloral hydrate for 7 days, stained 20 min in 0.1% cotton blue in water, and mounted in polyvinyl alcohol-glycerin-lactic acid (PVLG) (Omar et al., 1979). Spore germination and early stomatal development were observed.

Greenhouse — Inoculated plants were sprayed with conidia suspended in deionized water until runoff. Control plants were sprayed with water. Plants were considered infected when a stroma was observed on any leaf.

In experiment one, inocula from isolates of *E. panici* obtained from six Iowa prairies were adjusted to 15×10^6 conidia/ml. Each inoculum was sprayed individually and in all possible pairs on 4-mo-old BP. There were 2 replicate pots per treatment. Plants were covered with plastic bags for 72 hr after inoculation, and moist treatment was continued each night for 6 wk by placing pots in a $1.6 \text{ m} \times 1 \text{ m} \times 1 \text{ m}$ humidity box equipped with a wet sphagnum floor and a mister. Stromata were observed for sexual structures for 4 mo.

In experiment two, infection and disease development were compared among BP, IKP, and plants grown from rhizomes of previously uninfected Kalsow State Preserve plants (UKP). Plants were sprayed with 25×10^6 conidia/ml obtained from a Kalsow State Preserve isolate. Half of the plants from each source were covered with plastic bags for 72 hr. The remainder were left uncovered. Moist treatment for the covered plants was continued at night in the humidity box for 4 wk.

In experiment three, 17×10^6 conidia/ml were used to inoculate five pots of IKP and 2 pots of BP that had approximately equal numbers of leaves. All plants were covered with plastic bags for 72 hr immediately after inoculation, and also at night for 4 mo. Disease progression on emerging plants and their individual leaves was monitored and marked approximately every 10 days. Different tillering rates between IKP and BP resulted in unequal numbers of leaves assessed.

Spore Studies

Conidia — Conidial germination was observed in spore suspensions on slides in sterile moist chambers, on water agar (WA) or on PDA plates incubated at 25 C.

The effect of temperature on conidial germination was observed by preparing 0.3 ml of a 1×10^6 conidia/ml deionized water suspension from a 3-wk-old culture of a greenhouse reisolat, spreading on 3- or 4-day-old WA plates and incubating in the dark at 5, 10, 15, 20, 25, and 30 ± 1 C. One hundred spores per plug were observed on each of three plugs, randomly cut with a #6 corkborer from three replicate plates from each temperature after 24, 72, and 120 hr. Spore germination and chlamyospore formation were noted.

Ascospores — To study ascospore discharge and germination, ascostromata were removed aseptically from leaves, added to 2 ml of sterile deionized water, and the suspension spread on WA slides placed in sterile moist chambers. Slides were held at 4 or 25 C and observations were made periodically for 12-48 hr.

RESULTS

Field Observations

Panicum virgatum infected with *E. panici* was collected at 43 Iowa sites in 26 counties (Fig. 1). Commercial varieties planted along roads were monitored for disease but were not observed to be infected with *E. panici*. *E. panici* was never found on any other host, including other species of *Panicum*.

Overwintering — Conidia were observed on stromata on overwintered *P. virgatum* residue in early spring of 1982-1984. Conidiophores



Fig. 2. Mature stromata of *Elsinoë panici* on adaxial leaf surface of *Panicum virgatum*, collected on August 17, 1983, from Freda Haffner Kettlehole State Preserve (Bar equals 4.0 mm).

in acervuli on 1983 culms and leaves of *P. virgatum* overwintered in mesh bags contained conidia from April to early June, 1984, or produced them when overwintered material was placed in moist chambers. Ascostromata produced in Fall 1983 were recovered occasionally from overwintered material next spring, but were less intact than stromata. No asci or ascospores were present, but some conidia were observed.

Infection and disease progression — The first signs of infection on emerging leaves of *P. virgatum* at Doolittle Prairie were raised, translucent spots approximately 1×1 mm observed during mid-June to early July. Acervuli developed in these stromata after 1 wk. By mid-July, when inflorescences of *P. virgatum* plants were expanded, nearly 30% of leaves randomly collected were infected. Over 50% of leaves were infected by mid-August. Most stromata occurred on abaxial leaf surfaces, but moderate to heavily infected leaves had stromata on both surfaces (Fig. 2). Conidia were recovered from acervuli developed on stromata throughout the growing season. Lower leaves were infected first, and the fungus progressed slowly up the plant. By late summer or early fall, leaves, leaf sheaths, culms, the rachis, glumes, and lemmas were infected on heavily diseased plants. Some plants near but not in contact with diseased plants remained uninfected throughout the season. Disease patterns were similar during each year of the study.

Ascostromata were first observed on leaves collected from three prairies in August 1983. In 1984, ascostromata were observed on leaves collected from the same and from two additional sites (Fig. 1). Usually ascostromata developed separately, but occasionally were with larger asexual stromata.

Percentages of leaves with ascostromata collected from the Kalsow State Preserve from August 11 to October 5, 1984 are presented in Fig. 3. Percentages of #1 (lowest) and #2 (next to lowest) leaves with ascostromata remained considerably higher than for other leaves on the same plants. Similar patterns were seen from ascostromata collections at two other sites during the same period.

By August, ascostromata contained flattened eroded areas where mature asci had released ascospores. Mature asci were observed on leaves from Kalsow State Preserve from July 25 to September 28, 1984 (Fig. 4). Percentages of #1 leaves with functional ascostromata varied as the ascostromata matured and eroded. Number 2 leaves had similar patterns for comparable collection dates. A similar pattern of ascostroma, ascus, and ascospore development was observed at the two other sites during this period. The sexual system evidently continues to function from July to September as continued production of asci




Fig. 3. Percentages of #1 leaves (lowest) to #6 leaves (top) of *Panicum virgatum* with ascostromata of *Elsinoë panici* collected from Kalsow State Preserve, August-October, 1984.

occurs, replacing those that had been present in upper eroded areas of ascostromata.

Establishment of *E. panici* on Inoculated Plants

Detached Leaves — Detached inoculated leaves were cleared to examine germination and early hyphal development. Typically, conidia germinated within 3 days after inoculation. Conidia germinated by short, thick germ tubes, which darkened and developed into irregularly torulose hyphae, the width occupying about half the width of an epidermal long cell. Hyphae grew parallel to the long axis of epidermal cells and between epidermal cell walls. Hyphae quickly modified into masses of blocky cells comprising the young stromata. Each cell measured approximately $2.5 \times 2.5 \mu\text{m}$. Most young stromata were located over vascular bundles and associated with silica cells or trichomes. Conidial germination and infection of abaxial and adaxial surfaces were similar and consistent on the two different sources of *P. virgatum*.

Greenhouse — **Experiment one.** Stromata were observed on the abaxial surfaces of leaves on 4-mo-old BP 14 days after inoculation with a concentration of 15×10^6 conidia. Stromata developed on leaves inoculated with all six isolates and pairs of isolates, and conidia were produced in acervuli 2-3 wk after inoculation. With continued high humidity, new secondary infections developed after 1 mo on leaves that had emerged since inoculation. Stromata were examined periodically for sexual stages during the 4 mo after inoculation, but none was observed. Conidia were consistently present.

Experiment two. Typical stromata developed 3 wk after inoculation with 25×10^6 conidia/ml on BP, IKP, and UKP receiving high humidity. Inoculated plants without humidity treatment were not diseased.




Fig. 4. Percentages of mature asci of *Elsinoë panici* from 15-20 ascostromata of #1 (lowest) and #2 (next to lowest) leaves of *Panicum virgatum* collected at Kalsow State Preserve, July 25-September 29, 1984.

Table 1. Numbers of healthy and infected leaves occurring from secondary infections following inoculations with *Elsinoë panici* on *Panicum virgatum* Kalsow and Blackwell plants, February 5, 1984.

Plants	Days after inoculation																			
	41		55		62		70		76		84		90		99		108		119	
	H	D	H	D	H	D	H	D	H	D	H	D	H	D	H	D	H	D	H	D
Kalsow	N		52	12	69	17	95	19	M	M	109	56	106	84	259	119	246	147	186	239
Blackwell			N		33	0	33	0	33	0	33	0	M	M	M	M	25	8	16	21

H = number healthy leaves
 D = number of diseased leaves
 N = new plants emerging; tagged
 M = missing data

Experiment three. All pots of IKP and BP developed stromata with conidia 17 days after inoculation with 17×10^6 conidia/ml. Fifty-five days after inoculation, 5% of the newly emerged leaves of IKP were infected (Table 1). Emerging plants expressed disease approximately 10 days after emergence. Seventy days after inoculation, 17% of the leaves on IKP plants were diseased. The percentage of diseased leaves decreased for IKP plants as leaves emerged at a rapid rate. Ninety days after inoculation, 44% of leaves on IKP plants were infected. Leaves of BP plants showed no secondary infections until 119 days after inoculation, when disease had increased on BP and IKP leaves to nearly equal percentages.

In all greenhouse inoculations, the disease progressed slowly in individual plants and spread from plant to plant in the same pot. Leaves became diseased at different rates, but lower leaves were infected before upper ones. Only after the disease was well established on abaxial surfaces were symptoms seen on the adaxial surface. Uninoculated IKP plants in adjacent pots remained uninfected throughout the experiment. Ascstromata were not observed.

Spore Studies

Conidia — Hyaline to lightly pigmented conidia from any source, such as field collections, cultures from diseased leaves, budding ascospores, or subcultures were heterogeneous in size and shape (Fig. 5). Most conidia were slightly elongate and uninucleate (Fig. 6). On diseased leaves, conidia were produced from flask-shaped conidiophores lining acervuli, but in culture, conidia were either produced infrequently from randomly occurring conidiophores (Fig. 7) or more often budded from germ tubes, chlamydo-spores, or conidia (Figs. 8, 9, 10). Conidia germinated by germ tubes or by budding. Germination by germ tubes was typically monopolar, but occasionally a germ tube would emerge from the side of a conidium, and bipolar germination occasionally was observed (Figs. 11, 12, 8).

Germination of conidia was extremely variable among experiments conducted to determine temperature effects (Fig. 13). Germination was highest at 5-15 C, declined at 20-25 C, and increased at 30 C. The same reisolate, subcultured for each experiment, was used for the first three experiments. A different isolate was used for experiment four, and this may account for the relatively low percentage germination of this isolate. Because of the extreme variation among experiments, they were not combined as replications to satisfy conditions required for an analysis of variance.

Between 25-35 C, many conidia enlarged and became round with dense cytoplasm. These structures were interpreted as chlamydo-spores formed by direct conversion of a conidium. These chlamydo-spores were similar to detached terminal and intercalary chlamydo-spores formed in mycelial cultures.

Ascospores — Ascii from ascostromata collected August 17, 1984, from Doolittle Prairie spontaneously released ascospores in water mounts. Ascospores germinated readily in water by budding elongate

conidia (Fig. 14). Most young buds were $3.3 \times 3.3-4.4 \mu\text{m}$, but grew to typical conidial size. One-septate ascospores germinated as readily as 2- or 3-septate ascospores. Each cell of the ascospore was capable of germinating, and sometimes some cells of a single ascospore produced germ tubes while other cells budded conidia. Conidia formed during germination continued to develop conidia by budding. Occasionally, ascospores still within an ascus would germinate by forming germ tubes which grew through the ascus walls.

CONCLUSIONS AND DISCUSSION

Life Cycle

Panicum virgatum plants in Iowa resume growth from early to late June. About the same time, when temperatures are favorable and during periods of high humidity and dew, conidia develop from acervuli in overwintered stromata in host debris. Because conidia are thin-walled spores, they probably do not survive the winter. No ascospores were observed from overwintered stromata or ascostromata; thus, conidia seem to be the primary inoculum.

During June, splashing rain and dew spread conidia to leaves of young *P. virgatum* plants where conidia germinate during cool (10-15 C) periods of high humidity or dew, usually during the night and early morning hours. Spores can germinate from 5 to 30 C. Germination at higher temperatures is important in establishing secondary infections later in the season.

Conidia germinate, producing hyphae that infect leaves and modify into stromata by 3 days after inoculation. Conidia may form resistant chlamydo-spores at 25-35 C. Conidia budded from chlamydo-spores at 30 C may increase inoculum when this temperature occurs in the field during the summer. Conidia germinate by either forming germ tubes or budding, but chlamydo-spores germinate only by budding elongate conidia.

One to two wk after leaf emergence, tiny, translucent microscopic stromata appear on leaves. Young stromata can develop acervuli and produce conidia within 1-2 wk. Thus, approximately every 4 wk, a new generation of conidia can develop and be available to infect new leaves. Conidia are disseminated by rain and dew.

Stromata increase in number as the season progresses. Colonization by *E. panici* is first observed on lower leaves and progresses up the plant, eventually involving all leaves. Stromata become black and firm as they mature, and by August, diseased leaves are obvious because of the numerous shiny black stromatal areas. Acervuli on stromata continue to produce conidia. Although no chlorosis or necrosis is associated with spots, some leaves occasionally display a red pigmentation around the periphery of the black stromata.

In addition to stromata, softer ascostromata containing asci and ascospores may develop, primarily on the lower two leaves of plants, from late July until the first frost. These two types of stromata differ in stromal morphology (Gabel, 1985), distribution among sites, predic-

tability of development any year in the field, and leaves colonized.

Shortly after release, ascospores germinate by germ tubes or by budding conidia. Thus, ascospores could be a source of inoculum in late summer or early fall. As early as August, many ascostromatal areas are eroding, accompanied by ascospore release from mature asci. Some ascostromata may overwinter and can produce conidia in the spring.

In fall, diseased leaves become part of the prairie litter or remain on the persistent culms. *E. panici* does not colonize such dead host material. In spring, conidia produced from stromata on partially decomposed and scattered leaf debris are the source of primary inoculum. When spring burning of prairies as a management practice removes overwintering plant debris, the fungus is usually eliminated for the following growing season.

The life cycle of *E. panici* is similar to that of other species of *Elsinoë*. Conidia of *E. corni* Jenkins and Bitanc. (Jenkins et al., 1953) and *E. ampelina* Shear (Brook, 1973) served as primary inoculum. However, ascospores as well as conidia were primary inoculum of *E. veneta* (Burk.) Jenkins (Jones, 1924).

Optimum temperature for conidial germination was different for some species, for example 22-26 C for *E. veneta* (Jones, 1924) compared with 10-15 C for *E. panici*.


Two species, *E. fawcettii* Bitanc. and Jenkins (Whiteside, 1975) and *E. brasiliensis* Bitanc. and Jenkins (Zeigler and Lozano, 1983) produce two types of conidia. Hyaline conidia germinate by germ tubes or budding and colored conidia only by budding. Colored conidia of these species could be similar in function to the chlamydospores of *E. panici*.

Incubation periods of 6 days for *E. veneta* (Jones, 1924), 10-14 days for *E. brasiliensis* (Zeigler and Lozano, 1983), and 7-14 days reported here for *E. panici* are similar. Brook (1973) reported a decrease of incubation period for *E. ampelina* with an increase in temperature.

In contrast with conidia of *E. ampelina* which initiated infections up to 10 m in 4 mo (Brook, 1973), conidia of *E. panici* were not widely disseminated.

The sequence of infections of *E. veneta* on new cane growth (Jones, 1924) is very similar to the progression observed with *E. panici*.

Elsinoë panici does not elicit a chlorotic or necrotic response on its



Figs. 5-12. *Elsinoë panici* conidia.

Fig. 5. Wet mount of heterogeneous conidia from culture (Bar equals 10 μ m). Fig. 6. Elongate conidium with a single nucleus from a squash of field material, stained with Whittman's aceto-iron hematoxylin observed with phase-contrast optics (Bar equals 10 μ m). Fig. 7. Wet mount of conidia formed from conidiophores in culture (Bar equals 10 μ m). Fig. 8. Wet mount of elongate conidium with bipolar germ tubes budding (Bar equals 10 μ m). Fig. 9. Wet mount of elongate conidia budding from chlamydospores cultured on potato dextrose agar at 35 C (Bar equals 10 μ m). Fig. 10. Wet mount of elongate conidia budding from chlamydospores cultured on potato dextrose agar and observed with phase-contrast optics (Bar equals 10 μ m). Fig. 11. Wet mount of monopolar germination of conidium in culture (Bar equals 10 μ m). Fig. 12. Wet mount of conidium producing a germ tube (see arrow) from the median area (Bar equals 10 μ m).




Fig. 13. Conidia germination means at different temperature averaged over 24, 72, and 120 hr for each of 4 experiments (1, 2, 3, and 4) conducted with isolates of *Elsinoë panici* from Doolittle Prairie. Each point represents % germination of 2,700 spores.

graminaceous host. Only a red discoloration was associated with stromata of *E. sacchari* on specimens (No. 541 and 522) on loan from the New York Botanical Garden Mycological Herbarium (NY). *Myriangium bambusae* specimens (NY) developed stromata with no associated chlorosis or necrosis. However, many species of *Elsinoë* attacking dicotyledonous hosts produce necrotic areas (Jenkins, 1931a, 1932; Jenkins et al., 1946; and Massey and Jenkins, 1935).

Ascospores of *E. panici* germinate to form germ tubes or bud, functioning similarly to ascospores of *M. bambusae* (Tai, 1931), *E. canavaliae* Rac. (Jenkins, 1931a, b), *E. veneta* (Jones, 1924), and *E. brasiliensis* (Zeigler and Lozano, 1983) if microconidia, conidia, and sprout cells of these fungi are interpreted as similar structures.

Distribution and Persistence

A collection of diseased *P. virgatum* in Story County, Iowa, made by C.E. Bessey in 1882 was determined to be infected with *E. panici* (Tiffany and Mathre, 1961), documenting the disease in central Iowa over 100 years. Collections of *E. panici* on *P. virgatum* made during the late 1950s to early 1960s document the disease from Story, Pocahontas, Boone, and Webster counties in central and west-central Iowa (Tiffany and Mathre, 1961). Collections made during the present study document the disease from these same counties and confirm that *E. panici* frequently infects native *P. virgatum* throughout Iowa. *P. virgatum* is widely distributed in Iowa and is common in many prairie remnants (Pohl, 1966). The few collections of *P. virgatum* from northeastern Iowa examined in the present study were not diseased. Collections from north-central Nebraska (Keya Paha and Holt counties) and western South Dakota (Lawrence and Harding counties) indicate that the disease may be widespread.




Fig. 14. Ascospores of *Elsinoë panici* collected from 1984 ascostromata germinating by budding and germ tubes after 24 h in deionized water (Bar equals 10 µm).

ACKNOWLEDGEMENTS

The senior author thanks the Sigma Xi Scientific Research Society for a grant, which partly supported this study. We are grateful to Black Hills State College support services, for their assistance in the preparation of this manuscript. This work is part of a dissertation by the senior author submitted in partial fulfillment of the Ph.D. degree requirement at Iowa State University.

REFERENCES

BROOK, P.A. 1973. Epidemiology of grapevine anthracnose caused by *Elsinoë ampelina*. N.Z.J. Agric. Res. 16:333-342.
 GABEL, A. 1985. Life history, development, and host-parasite relations of *Elsinoë panici* Tiffany and Mathre. Ph.D. Dissertation, Microfilm No. 85-24653, Iowa State Univ., Ames.
 JENKINS, A.E. 1931a. Scab of *Canavalia* caused by *Elsinoë canavaliae*. J. Agric. Res. 42(1):1-12.
 ———. 1931b. Lima bean scab caused by *Elsinoë*. J. Agric. Res. 42(1):13-23.
 ———. 1932. Rose anthracnose caused by *Sphaceloma*. J. Agric. Res. 45:321-337.
 ———, M.J. FORSELL, and L.G. BOYLE. 1946. Identity and known distribution of *Elsinoë piri* in Washington and Oregon. Phytopathology 36(6):458-461.
 ———, J.H. MILLER, and G.H. HEPTING. 1953. Spot anthracnose and other leaf and petal spots of flowering dogwood. Natl. Hortic. Mag. 32(2):57-69.
 JONES, L. 1924. Anthracnose of cane fruits and its control on black raspberries in Wisconsin. Wis. Agric. Exp. Stn. Bull. 59. 25 pp.
 LO, T.C. 1964. *Elsinoë* stage of *Sphaceloma sacchari*. Proc. Biol. Soc. Wash. 77:1-4.
 LUTTRELL, E.S. 1973. Loculoascomycetes. pp. 135-210. In G.C. Ainsworth, F.K. Sparrow, and A.S. Sussman (eds.) The fungi an advanced treatise. Vol. IVA. Academic Press, New York, New York.
 MASSEY, L.M., and A.E. JENKINS. 1935. Scab of violet caused by *Sphaceloma*. Cornell (Ithaca) Agric. Exp. Stn. Mem. 176. 9 pp.
 OMAR, M.B., L. BOLLAND, and W.A. HEATHER. 1979. A permanent mounting medium for fungi. Bull. Br. Mycol. Soc. 13:13-32.
 POHL, R.W. 1966. The grasses of Iowa. Iowa State J. Sci. 40(4):341-566.

- TAL, F.L. 1931. Observations on the development of *Myriangiium bambusae* Rick. Sinensia 1(10):147-164.
- TIFFANY, L.H., and J.H. MATHRE. 1961. A new species of *Elsinoë* on *Panicum virgatum*. Mycologia 53(6):600-604.
- TODD, E.H. 1960. *Elsinoë* diseases of sugarcane in Florida. Plant Dis. Rep. 44(3):153.
- WHITESIDE, J.O. 1975. Biological characteristics of *Elsinoë fawcettii* pertaining to the epidemiology of sour orange scab. Phytopathology 65:1170-1175.
- ZEIGLER, R.S., and J.C. LOZANO. 1983. The relationship of some *Elsinoë* and *Sphaceloma* species pathogenic on cassava and other Euphorbiaceae in Central and South America. Phytopathology 73(2):293-300.