

7-31-2020

Modeling the Griffiths Phase in Manganese Intercalated Tantalum Disulfide - Slides

Aaron Janaszak
University of Northern Iowa

Lukas Stuelke
University of Northern Iowa

See next page for additional authors

Follow this and additional works at: <https://scholarworks.uni.edu/ugswork>

Let us know how access to this document benefits you

Recommended Citation

Janaszak, Aaron; Stuelke, Lukas; and Shand, Paul, "Modeling the Griffiths Phase in Manganese Intercalated Tantalum Disulfide - Slides" (2020). *Undergraduate Student Work*. 20.
<https://scholarworks.uni.edu/ugswork/20>

This Open Access Undergraduate Student Work is brought to you for free and open access by UNI ScholarWorks. It has been accepted for inclusion in Undergraduate Student Work by an authorized administrator of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

Authors

Aaron Janaszak, Lukas Stuelke, and Paul Shand

Simulating Griffiths Phase Behavior

Aaron Janaszak and Lukas Stuelke

UNI Summer Undergraduate Research Program

Mentor: Dr. Paul Shand





What is the Griffiths Phase?


- In a paramagnet, at high temperatures, inverse magnetization follows the Curie-Weiss law and is linear.
- At lower temperatures, clusters align and magnetization deviates from the Curie-Weiss law.

Inverse Magnetization is linear at higher temperatures


In the Griffiths phase, this linearity is interrupted as shown by the exaggerated curve




Fully Occupied Crystalline Lattice



Clusters begin to dominate, displaying Griffiths phase behavior at low temperatures




Some Moments Removed



Purpose


Can we reproduce the Griffiths Phase behavior using a computer model?

And if so, what variables influence this behavior the most?




Methods & Data

- Langevin function used to simulate clusters in MATLAB
- Data collection at many values of the external magnetic field




Comparison Between Experimental Results and Computer Model

Experimental Data



Computational Model





Any Questions?

The Griffiths Phase is a result of a spontaneous increase in the overall magnetization of the system primarily influenced by the rare, large clusters transitioning to a ferromagnetic state at the Curie Temperature. We were able to create a MATLAB program that exhibits this trend by using the Langevin function to simulate clusters.

Future Work:

- Exploration of variables and distribution types

Acknowledgements:

- Dr. Paul Shand
- Paul White
- UNI Physics Department
- US DOE Grant #DE-SC0020334

