

3-2019

## Broadening the Use of Perennial Native Vegetation for Nutrient Reduction on Marginal Farmland

Justin Meissen  
*University of Northern Iowa*

Ashley Kittle

*See next page for additional authors*

*Let us know how access to this document benefits you*

Copyright ©2019 Justin Meissen, Ashley Kittle, Laura Jackson

Follow this and additional works at: [https://scholarworks.uni.edu/tpc\\_facpub](https://scholarworks.uni.edu/tpc_facpub)



Part of the [Plant Sciences Commons](#)

---

### Recommended Citation

Meissen, Justin; Kittle, Ashley; and Jackson, Laura, "Broadening the Use of Perennial Native Vegetation for Nutrient Reduction on Marginal Farmland" (2019). *UNI Tallgrass Prairie Publications and Reports*. 22.

[https://scholarworks.uni.edu/tpc\\_facpub/22](https://scholarworks.uni.edu/tpc_facpub/22)

This Report is brought to you for free and open access by the Publications and Reports at UNI ScholarWorks. It has been accepted for inclusion in UNI Tallgrass Prairie Publications and Reports by an authorized administrator of UNI ScholarWorks. For more information, please contact [scholarworks@uni.edu](mailto:scholarworks@uni.edu).

**Offensive Materials Statement:** Materials located in UNI ScholarWorks come from a broad range of sources and time periods. Some of these materials may contain offensive stereotypes, ideas, visuals, or language.

---

**Authors**

Justin Meissen, Ashley Kittle, and Laura Jackson

# Broadening the Use of Perennial Native Vegetation for Nutrient Reduction on Marginal Farmland

Justin Meissen, Ashley Kittle, Laura Jackson | Mar 2019

justin.meissen@uni.edu

## Project Highlights

- **Key prairie species establish well even in dry conditions**
  - Native warm and cool season grasses (e.g. big bluestem, wild rye) establish well in marginal dry soil, along with important summer and fall wildflowers (e.g. ox-eye, sunflowers)
- **Seed mixes customized for dry soils result in more ecological functions at similar price**
  - Dry adapted wildflowers established better than their medium to wet soil counterparts, resulting in more options for wildlife and pollinators- the entire spring forb functional group was missing from the non-customized mix
- **Cost-effectiveness of native perennial vegetation comparable in productive vs marginal soils**
  - Plants per dollar similar for many native grass and forb species

## Background

### Why Should Perennial Vegetation Nutrient Reduction Practices Target Marginal Lands?

- Lower barrier to adoption, often best choice economically when paired with conservation programs
- Reduce inputs altogether, puts unproductive land to use through multiple ecological benefits

### Partnerships to Broaden Implementation Research

- Pheasants Forever, NRCS, Fayette County Conservation, TPC
- Multiple stakeholders using same retired farmland as research site for ag-oriented perennial vegetation questions

### Objective: Compare Establishment and Cost Effectiveness for Different Seed Mixes That Differ In Soil Type Customization On Dry Marginal Soils

- Randomized design ( $n=4$ ) planted November 2017 next to Wapsipinicon River near Fairbank, IA
- Two seed mixes from TPC seed calculator: 1) medium soil (\$365/ac, 46 species), 2) dry soil(\$368/ac, 49 species)

Figure 1. Study layout.



Figure 2. Marginal lands context.




Figure 3. Site in Fall 2018.

