

1983

A Statewide Screening For Acid Rainfall In Iowa

Jack O. Kennedy
University of Iowa

Andrew E. Granston
University of Iowa

Roger C. Splinter
University of Iowa

Let us know how access to this document benefits you

Copyright ©1983 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Kennedy, Jack O.; Granston, Andrew E.; and Splinter, Roger C. (1983) "A Statewide Screening For Acid Rainfall In Iowa," *Proceedings of the Iowa Academy of Science*, 90(2), 41-43.

Available at: <https://scholarworks.uni.edu/pias/vol90/iss2/3>

This Research is brought to you for free and open access by the IAS Journals & Newsletters at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

Offensive Materials Statement: Materials located in UNI ScholarWorks come from a broad range of sources and time periods. Some of these materials may contain offensive stereotypes, ideas, visuals, or language.

A Statewide Screening For Acid Rainfall In Iowa

JACK O. KENNEDY, ANDREW E. GRANSTON, JR., AND ROGER C. SPLINTER

University Hygienic Laboratory, The University of Iowa, Iowa City, Iowa 52242

From April through October 1980, a statewide screening for acid rainfall was conducted in Iowa. Ninety-seven National Weather Service observers were provided with pH color-indicating strips and measured and recorded the pH of every precipitation sample. Results of the screening indicate pH values of rainfall ranged from a low of 4.0 to a high of 7.0. The pH values 5.7 and 5.9 were observed most often and represented 66% of the 4,197 values measured. Approximately 80% of the rainfall pH values fell in the 5.7 to 7.0 range and 20% in the acid rainfall range (5.6 or less). Median pH values calculated for the 97 sampling locations ranged from 5.1 to 6.2, with medians of 5.7 and 5.9 occurring most frequently. With the exception of one 3-county area, most of the 11 sampling locations demonstrating acid rainfall (median values 5.6 or less) were widely separated and probably represented localized problems. Three adjacent counties located in southeast Iowa had low median pH values and this area is recommended for future study.

INDEX DESCRIPTORS: Iowa, acid rainfall, pH values, National Weather Service

For Iowa, an intensely agricultural state, the need for precipitation to initiate and sustain plant growth and to replenish streams, lakes, and ground water supplies is ever present. Much concern has been expressed in the last few years regarding precipitation and changes in its acidity (pH). Pure rainwater in equilibrium with carbon dioxide has a pH value of 5.65. Recent reviews (1, 2) of available data indicate that precipitation in a large region of North America is well below 5.65 (rain with a pH below 5.6 is considered acidic rain). The increasing acidity of rainfall is believed to be the result of the accumulation of certain acids in the atmosphere. These acids are produced by dissolution of sulfur and nitrogen oxides in precipitation and are created primarily from the combustion of fossil fuels. Originally, acid rainfall was found in the eastern and north-eastern portions of the United States, which are highly industrialized areas utilizing fossil fuels to a relatively high degree. Recent information, however, indicates that "acid precipitation has spread measurably southward and westward in the United States." (1) Researchers in Minnesota have observed acid rainfall (3) and are concerned about potential damage to both terrestrial and aquatic ecosystems. The Minnesota Legislature, recognizing the extent and severity of the acid rain problem, passed a bill designated to identify, control, and abate acid rain. In Iowa, very little is known about acid rainfall. Tabatabai, at Iowa State University, has conducted research on the nutrient content and pH of samples of rainfall from seven locations around the state. He found that the average pH value of rainfall and snowmelt samples was about 6 (3, 4). In an effort to expand and improve on the limited data base, we conducted a statewide screening for acid rainfall.

MATERIALS AND METHODS

The National Weather Service maintains a voluntary group of observers throughout Iowa to record daily temperature and precipitation. Upon obtaining approval from the National Weather Service, an informational mailing was sent to 150 observers, requesting their assistance in determining rainfall pH. Over 125 positive responses were received, from which 120 were selected and provided with the necessary instructions and materials. The approximate locations of all participating observers are shown in Figure 1. The study began in April and continued through October 1980.

In order to obtain as much statewide information as possible, utilize non-scientific personnel, and measure pH as soon as possible following precipitation, pH color-indicator sticks were used to determine the rain pH. The pH-indicator sticks (colorpHast indicator sticks, MCIB Manufacturing Chemicals, Inc., Cincinnati, Ohio) covered a range from 4.0 to 7.0 with a 0.2 sensitivity from 5.3 to 5.9 and a 0.3 to 0.5 sensitivity on all other readings. The manufacturer has established color codes for the pH values 4.0, 4.4, 4.76, 5.0, 5.3,




Fig. 1. Map of Iowa counties showing approximate rainfall sampling locations

5.5, 5.7, 5.9, 6.2, 6.5 and 7.0. A pH reading was obtained by immersing the stick in rain water, which then developed a color that was compared to a standard color chart with associated pH values. The pH values and rainfall amounts were recorded on data sheets and returned when the study was completed. Accuracy of the observers' readings was determined by sending laboratory prepared buffers for pHs 4.0, 5.0 and 6.0 to selected observers. Of the 3 buffers sent, 100% correctly measured the 4.0, and 95% correctly measured the 5.0 and 6.0.

RESULTS AND DISCUSSION

Data return frequency was somewhat lower than anticipated; data sheets were received from 101 out of 120 stations (84%). Four of the 101 were unusable (no name, forgot to do tests, etc.) leaving 97 that were used in developing data for this report. The 97 locations (Figure 1) represent most areas of the state and 69 of the Iowa's 99 counties. Statewide pH values ranged from a low of 4.0 to a high of 7.0. Figure 2 is a bar graph representing the number of occurrences of each pH value between 4.0 and 7.0. The pH values 5.7 and 5.9 were observed most often (39% and 28% of the time, respectively) and represented 66% of the 4,197 values measured. Approximately 79% of the values




Fig. 2. Number of rainfall samples for each pH value

observed were in the range from 5.7 to 7.0. Of the 21% occurring in the acid rainfall range (below 5.7), slightly over 1% (43) were below pH 5.0. In assessing these data, it was noted that several values were reported for which there was no manufacturer's color code. Upon review, it was determined that over 95% of the values were based on the color chart provided, whereas approximately 4.7% were best estimates between two given values. In addition, the mean pH value (computed by averaging concentrations and converting to pH) for all pH values was 5.62, which compares favorably to the value of 5.65 for rainfall in equilibrium with carbon dioxide.

Because of the difficulties inherent in the calculation of mean pH values, median values were used in this report. Median values were calculated for each of the 97 locations and ranged from 5.1 to 6.2 (Figure 3). Eighty-two (84%) of the median values were pH 5.7 or 5.9, while 11 (11%) were less than 5.7 and 4 (4%) were greater than 5.9. Of the 11 locations with medians less than 5.7, 2 locations had median values of 5.6, 5 had 5.5, 3 had 5.3 and 1 had 5.1. The medians were recorded geographically (Figures 4 and 5) for trend analysis. Figure 4 identifies locations having median pH values of 5.7




Fig. 3. Number of rainfall sampling locations of each median pH value




Fig. 4. Map of Iowa counties showing approximate sampling locations with median rainfall pHs of 5.7 or higher

or greater. Eighty-six locations appear in Figure 4 and geographically cover the entire state. The 11 sampling locations with median values below 5.7 are shown in Figure 5. With the exception of the 3-county area in southeast Iowa, locations with median values below 5.7 are widely separated and scattered throughout the state.

In evaluating for trends, another type of analysis applied to these data was frequency of occurrence of acid rainfall. For each station the number of pH values 5.6 or less was tabulated and divided by the total number of values reported for that station. The resulting ratio from the 97 sampling stations are expressed as percentages in Table 1.

From these data it is apparent that sampling locations with frequencies higher than 80% are significantly different from the other locations. Moreover, locations with frequencies of 80% or higher




Fig. 5. Map of Iowa counties showing approximate sampling locations with median rainfall pHs of 5.6 or lower

Table 1. Frequency of Occurrence of Acid Rainfall

Frequency of Acid Rainfall (Percentage of values 5.6 or less)	Number of Sampling Locations Having Stated Percentage
0 to 10%	43
11 to 20%	19
21 to 30%	14
31 to 40%	6
41 to 50%	5
51 to 60%	3
61 to 70%	1
71 to 80%	0
81 to 90%	1
91 to 100%	5

Table 2. Sampling locations with median pH values 5.6 or less and acid rainfall frequency.

Sampling Locations	Median pH	Frequency of Acid Rainfall (%)
Page	5.6	69
Butler	5.6	50
Sioux	5.5	100
Henry	5.5	98
Calhoun	5.5	56
Monona	5.5	54
Dubuque	5.5	53
Clinton	5.3	92
Washington	5.3	91
Allamakee	5.3	85
Keokuk	5.1	100

represent areas where acid rainfall was clearly occurring the majority of the time during the sampling period. Table 2 lists the 11 sampling locations with median values of 5.6 or less and their corresponding frequencies of acid rainfall. This method of analysis better defines the problem areas. In the preliminary evaluation the 7 median values of 5.5 and 5.6 were considered borderline for acid rainfall. From this analysis, Sioux and Henry County locations show a much higher frequency of acid rainfall than the other 5.5 and 5.6 locations and should be grouped with the lower median pH locations (Clinton, Washington, Allamakee and Keokuk) for evaluation. The data suggest that if acid rainfall is exerting a significant effect in Iowa, it would be at these 6 locations. Three of these locations (Sioux, Clinton, and Allamakee) are geographically isolated from one another, indicating that the source of the low values is probably independent for each. In addition, there are several other sampling locations near those counties with higher (less acidic) median values, and this also indicates a localized problem. One may speculate that the low value seen in Clinton County may be a result of the highly industrialized area located along that reach of the Mississippi River. The 3 other sampling locations with low median values and high frequencies of acid rainfall are in close geographical proximity, occurring in adjacent counties (Henry, Washington, and Keokuk).

The Keokuk County location had the lowest median value statewide with 26 of its 56 values less than pH 5.0 and the highest frequency of acid rainfall (100%). It is in the area defined by these 3 counties that Iowa's most significant acid rainfall effect may be occurring and this area should be considered for future study.

ACKNOWLEDGEMENTS

We would like to especially acknowledge the help and cooperation of all the participating National Weather Service observers without whom this study could not have been conducted.

REFERENCES

1. GLASS, N. R., G. E. GLASS and P. J. RENNIE. 1979. Effect of acid precipitation. *Environmental Science and Technology* 13:1350-1355.
2. U.S. ENVIRONMENTAL PROTECTION AGENCY. 1980. Acid rain. EPA-600/9-79-036. Office of Research and Development. Washington, D. C.
3. MINNESOTA POLLUTION CONTROL AGENCY. 1980. Minnesota Water Quality-305(b) report. Roseville, Minnesota.
4. TABATABAI, M. A. and J. M. LAFLEN. 1976. Nitrogen and sulfur content and pH of precipitation in Iowa. *Journal of Environmental Quality* 5:108-112.
5. TABATABAI, M. A. and J. M. LAFLEN. 1976. Nutrient content of precipitation over Iowa. Proceedings of the First International Symposium on Acid Precipitation and the Forest Ecosystem. USDA Forest Service. General Technical Report. NE-23:293-308.