

1970

## Distribution of Soils by Natural Drainage Class and by Slope Class for Iowa Counties

E. C. A. Runge

*University of Illinois at Urbana-Champaign*

R. I. Dideriksen

F. F. Riecken

*Iowa State University*

*Let us know how access to this document benefits you*

Copyright ©1970 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

---

### Recommended Citation

Runge, E. C. A.; Dideriksen, R. I.; and Riecken, F. F. (1970) "Distribution of Soils by Natural Drainage Class and by Slope Class for Iowa Counties," *Proceedings of the Iowa Academy of Science*, 77(1), 61-85.

Available at: <https://scholarworks.uni.edu/pias/vol77/iss1/12>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact [scholarworks@uni.edu](mailto:scholarworks@uni.edu).

## Distribution of Soils by Natural Drainage Class and by Slope Class for Iowa Counties<sup>1</sup>

E. C. A. RUNGE, R. I. DIDERIKSEN, and F. F. RIECKEN<sup>2</sup>

*Abstract.* Natural drainage class (namely, poor, imperfect, and well drained) and slope class are estimated for Iowa counties through a 2% ¼-section sample soil survey. Estimate of soil drainage class is given also by slope class. Counties vary widely in the amount of poorly drained soils. Kossuth County has 56.4% of poorly drained soils, while Allamakee has less than 1%. Land use management and conservation considerations are discussed briefly.

The degree to which Iowa soils become wet or near water-saturated affects their capability for crop use. It also affects their use and management for other purposes; for example, as sites for homes, roads, airports, and solid waste disposal.

Iowa counties differ widely in the amount of soil that is seasonally wet. It is the objective of this study to present information on county and state distribution of soils of differing slope and drainage class.

Three degrees of natural wetness are used in this study. They are corn crop-production biased and are: (a) poorly drained, (b) imperfectly or somewhat poorly drained, and (c) well drained. Poorly drained soils are commonly seasonally water-saturated. If cultivated, they are usually tile drained, if permeable enough. Imperfectly drained soils may be in need of some drainage improvement in some seasons. Well drained soils are seldom in need of drainage improvement. Criteria for placing soils in a drainage class are discussed later.

Seven slope classes of soils are used, and slope class of drainage class is given.

### DATA

This study used the data collected from a 2%, stratified, random sample of the land area in Iowa. Soil mapping started in 1947. Each soil mapped was identified by soil profile, slope class, and erosion class. The Soil Conservation Service was responsible for mapping three individual ¼-section tracts per township. This sampling technique was later adopted by the USDA to study the

<sup>1</sup> Journal Paper No. J-6555 of the Iowa Agriculture and Home Economics Experiment Station, Ames. Project No. 1191. Contribution from the Departments of Agronomy, Iowa State University, and University of Illinois at Urbana-Champaign.

<sup>2</sup> Associate Professor of Soils, University of Illinois at Urbana-Champaign (formerly Assistant Professor, Iowa State University); State Soil Scientist, SCS, Indiana (formerly Assistant State Soil Scientist, SCS, Iowa); and Professor of Soils, Iowa State University.

soils and the conservation needs throughout the entire country. The nationwide study has been named the National Conservation Needs Inventory. The Conservation Needs Inventory data for Iowa were presented elsewhere (Iowa Conservation Needs Committee, 1963).

Two papers have presented some soil drainage and soil slope data (Arnold et al. 1960; Slusher et al. 1961). The "depressional" soils data of north-central Iowa of Slusher et al. are included in our "poorly drained" soils. Geometrical properties of these depressions have been investigated by Haan and Johnson (1967). The soil slope data of Arnold et al. are essentially similar to the slope data of this paper.

The report is presented in three sections. In the first section, soils are identified as belonging to one of three natural drainage classes: poor—needing drainage; imperfectly, or somewhat poorly, drained—sometimes needing drainage, particularly in the more northern areas of Iowa; and well drained—not needing drainage. Included in this section is a discussion of the criteria used in establishing natural drainage class.

The second section gives the percentage of the drainage classes of each county in each of the various slope classes (Table 1). The estimated acreage of each of the various mutual drainage and slope classes may be determined by multiplying the percentage of each class by the total acres in the county.

The third section gives some implications of the slope-drainage data to use, management, and conservation of Iowa soil resources.

#### IDENTIFICATION OF SOILS BY NATURAL DRAINAGE CLASS

Each soil mapped in the sample soil survey was assigned to one of the three natural drainage classes. The color of the subsoil, usually the 12-to 36-inch layer, is considered diagnostic and is given major emphasis in placing the soil in a drainage class. Important colors considered include those affected by the content, form, and distribution of iron in the subsoil (Dideriksen, 1966; Runge and de Leon, 1967). Surface horizons are also used, but overall they are not as important as the subsoil. Most soils of the study are easily assigned to a drainage class, but some are difficult to place. This is particularly true of the soils formed from alluvium. Where color gave indefinite criteria for determining drainage class, other soil morphological features, landscape position, and experience of soil scientists with the soil in question were used in placing it in a drainage class.

In Iowa, most poorly drained soils have gray to olive gray subsoils. These subsoil horizons have Munsell color hues of 2.5Y to 5Y with values between 3 and 7 and chroma of 2 or less. If values

Runge et al.: Distribution of Soils by Natural Drainage Class and by Slope Clas

1970]

DISTRIBUTION OF SOILS

63

Table 1. The Distribution of Soils by Slope and Natural Drainage Class for all Iowa Counties.

County and Total Acres in County	Natural Drainage	Percent Totals	% of County in Respective Slope Classes							Excluded Area
			0-1%	2-4%	5-8%	9-13%	14-17%	18-24%	25+%	
Adair 364,166	All classes	100.0	8.9	24.3	22.8	30.9	7.8	2.6	—	2.7
	Poor	7.49	3.95	2.96	0.10	0.48	—	—	—	—
	Imperfect	15.75	2.89	8.27	1.16	3.33	0.10	—	—	—
	Well	74.04	2.08	13.06	21.50	27.11	7.65	2.64	—	—
Adams 272,641	All classes	100.0	17.7	26.6	20.4	23.1	7.0	2.2	0.8	2.2
	Poor	15.29	9.21	2.83	1.83	1.30	0.12	—	—	—
	Imperfect	19.91	5.18	11.08	2.13	1.52	—	—	—	—
	Well	62.61	3.30	12.72	16.42	20.29	6.86	2.15	0.87	—
Allamakee 423,681	All classes	100.0	9.0	7.9	17.1	31.9	1.0	1.0	28.8	2.3
	Poor	0.07	—	.07	—	—	—	—	—	—
	Imperfect	2.48	1.47	1.01	—	—	—	—	—	—
	Well	94.17	7.55	6.85	17.05	31.92	1.00	1.01	28.79	—
Appanoose 334,720	All classes	100.0	12.8	21.0	21.6	15.7	15.7	6.9	6.0	0.3
	Poor	11.94	11.35	.57	0.02	—	—	—	—	—
	Imperfect	36.70	—	18.42	17.33	0.95	—	—	—	—
	Well	51.06	1.47	2.08	4.21	14.70	15.70	6.90	6.00	—
Audubon 286,721	All classes	100.0	6.4	26.3	16.2	34.3	10.7	4.8	0.2	1.1
	Poor	6.11	6.04	0.03	—	0.04	—	—	—	—
	Imperfect	17.81	.34	15.73	1.21	0.36	0.09	0.08	—	—
	Well	74.98	.02	10.57	15.00	33.85	10.65	4.73	0.16	—
Benton 459,522	All classes	100.0	9.4	53.2	19.8	6.5	3.7	2.2	0.6	4.6
	Poor	16.56	3.70	12.86	—	—	—	—	—	—
	Imperfect	14.02	4.18	8.60	1.24	—	—	—	—	—
	Well	64.88	1.60	31.71	18.60	6.48	3.72	2.18	0.59	—

Table 1. Continued

County and Total Acres in County	Natural Drainage	Percent Totals	% of County in Respective Slope Classes							Excluded Area
			0-1%	2-4%	5-8%	9-13%	14-17%	18-24%	25+%	
Black Hawk 363,519	All classes	100.0	36.0	42.6	13.0	0.5	0.1	0.1	0.1	7.6
	Poor	18.16	13.78	4.38	—	—	—	—	—	—
	Imperfect	23.73	13.45	10.18	0.11	—	—	—	—	—
	Well	50.45	8.75	28.06	12.94	0.46	0.07	0.06	0.11	—
Boone 366,725	All classes	100.0	33.5	48.3	10.0	1.6	0.6	—	3.4	3.1
	Poor	33.97	31.56	2.41	—	—	—	—	—	—
	Imperfect	19.56	1.74	17.82	—	—	—	—	—	—
	Well	43.76	0.16	27.99	10.00	1.62	0.61	—	3.38	—
Bremer 280,962	All classes	100.0	48.0	43.6	3.6	0.8	0.2	0.5	0.2	3.1
	Poor	24.70	21.65	3.05	—	—	—	—	—	—
	Imperfect	45.43	22.37	22.93	0.13	—	—	—	—	—
	Well	26.78	4.02	17.65	3.47	0.80	0.15	0.53	0.16	—
Buchanan 364,160	All classes	100.0	27.4	59.9	7.7	0.4	0.5	—	0.3	3.9
	Poor	17.05	11.75	5.30	—	—	—	—	—	—
	Imperfect	30.75	8.02	22.43	0.30	—	—	—	—	—
	Well	48.34	7.67	32.14	7.37	0.39	0.52	—	0.25	—
Buena Vista 369,917	All classes	100.0	45.0	41.6	7.5	1.3	0.8	0.4	2.8	0.7
	Poor	29.75	29.40	0.35	—	—	—	—	—	—
	Imperfect	19.29	13.33	5.95	0.01	—	—	—	—	—
	Well	50.35	2.24	35.28	7.47	1.36	0.83	0.40	2.77	—
Butler 372,480	All classes	100.0	44.1	37.2	10.1	3.7	0.5	0.5	0.2	3.7
	Poor	32.34	28.82	3.52	—	—	—	—	—	—
	Imperfect	24.20	10.98	12.73	0.49	—	0.01	—	—	—
	Well	39.75	4.32	20.92	9.63	3.75	0.50	0.47	0.16	—

Runge et al.: Distribution of Soils by Natural Drainage Class and by Slope Clas

Table 1. Continued

County and Total Acres in County	Natural Drainage	Percent Totals	% of County in Respective Slope Classes							Excluded Area
			0-1%	2-4%	5-8%	9-13%	14-17%	18-24%	25+ %	
Calhoun 366,720	All classes	100.0	61.3	31.8	2.9	0.3	0.3	0.3	0.1	3.0
	Poor	52.10	52.10	—	—	—	—	—	—	—
	Imperfect	26.04	6.60	19.44	—	—	—	—	—	—
	Well	18.88	2.63	12.39	2.91	0.33	0.33	0.25	0.04	—
Carroll 367,360	All classes	100.0	15.3	40.2	21.1	14.8	3.3	0.8	0.7	3.8
	Poor	12.05	11.88	0.17	—	—	—	—	—	—
	Imperfect	19.92	3.02	16.26	0.64	—	—	—	—	—
	Well	64.20	0.43	23.68	20.51	14.82	3.31	0.79	0.66	—
Cass 357,666	All classes	100.0	9.0	33.5	18.0	31.0	8.2	0.3	0.1	—
	Poor	8.35	6.21	1.63	0.01	0.20	0.30	—	—	—
	Imperfect	16.04	0.08	15.76	0.20	—	—	—	—	—
	Well	75.61	2.71	16.08	17.72	30.82	7.98	0.27	0.03	—
Cedar 374,401	All classes	100.0	16.6	40.6	22.0	9.9	4.9	2.1	0.6	3.3
	Poor	6.87	6.65	0.22	—	—	—	—	—	—
	Imperfect	16.45	6.99	9.46	—	—	—	—	—	—
	Well	73.30	2.98	30.95	21.95	9.91	4.90	2.10	0.61	—
Cerro Gordo 368,640	All classes	100.0	45.7	34.9	6.9	2.7	1.6	1.6	0.6	6.0
	Poor	32.83	31.19	1.64	—	—	—	—	—	—
	Imperfect	22.80	8.79	13.91	0.10	—	—	—	—	—
	Well	38.36	5.79	19.38	6.79	2.72	1.57	1.55	0.56	—
Cherokee 366,720	All classes	100.0	16.3	46.2	22.1	2.9	1.6	1.5	4.4	5.0
	Poor	11.30	8.83	2.47	—	—	—	—	—	—
	Imperfect	13.78	2.86	10.48	0.44	—	—	—	—	—
	Well	65.61	4.65	33.27	21.65	2.87	1.63	1.54	4.38	—

1970]

DISTRIBUTION OF SOILS

65

Table 1. Continued

County and Total Acres in County	Natural Drainage	Percent Totals	% of County in Respective Slope Classes							Excluded Area
			0-1%	2-4%	5-8%	9-13%	14-17%	18-24%	25+%	
Chickasaw 323,200	All classes	100.0	37.9	52.7	5.4	0.1	0.1	—	—	3.9
	Poor	27.33	22.47	4.86	—	—	—	—	—	—
	Imperfect	24.15	9.74	13.83	0.58	—	—	—	—	—
	Well	44.64	5.72	34.01	4.79	0.10	0.02	—	—	—
Clarke 274,563	All classes	100.0	13.2	18.0	25.0	14.3	16.4	6.7	2.2	4.2
	Poor	10.62	7.26	2.59	0.59	0.18	—	—	—	—
	Imperfect	40.67	0.43	14.04	21.79	4.30	0.11	—	—	—
	Well	44.61	5.55	1.39	2.65	9.86	16.26	6.72	2.18	—
Clay 367,356	All classes	100.0	54.3	35.5	7.2	1.7	0.4	0.2	0.6	0.1
	Poor	19.80	18.40	1.40	—	—	—	—	—	—
	Imperfect	33.66	24.54	9.12	—	—	—	—	—	—
	Well	46.43	11.37	25.00	7.18	1.74	0.40	0.18	0.56	—
Clayton 509,440	All classes	100.0	5.1	15.5	21.7	23.4	13.7	6.3	8.9	5.4
	Poor	1.30	1.00	0.30	—	—	—	—	—	—
	Imperfect	3.71	1.37	2.32	0.02	—	—	—	—	—
	Well	89.67	2.80	12.88	21.67	23.37	13.70	6.29	8.96	—
Clinton 444,800	All classes	100.0	30.5	27.6	14.4	10.9	5.2	5.4	1.0	5.0
	Poor	16.17	15.74	0.11	—	0.05	0.12	0.15	—	—
	Imperfect	15.10	10.97	4.13	—	.01	—	—	—	—
	Well	63.77	3.79	23.39	14.38	10.89	5.04	5.25	1.03	—
Crawford 458,245	All classes	100.0	8.7	19.1	21.0	29.9	11.1	5.5	1.7	2.9
	Poor	3.91	3.91	—	—	—	—	—	—	—
	Imperfect	5.39	0.07	5.22	1.10	—	—	—	—	—
	Well	86.75	4.75	13.89	19.89	29.94	11.05	5.45	1.78	—

Runge et al.: Distribution of Soils by Natural Drainage Class and by Slope Class

Table 1. Continued

County and Total Acres in County	Natural Drainage	Percent Totals	% of County in Respective Slope Classes							Excluded Area
			0-1%	2-4%	5-8%	9-13%	14-17%	18-24%	25+%	
Dallas 382,083	All classes	100.0	27.3	43.1	10.2	6.3	5.0	2.6	3.1	2.4
	Poor	25.40	23.28	2.10	0.02	—	—	—	—	—
	Imperfect	12.25	1.91	10.32	—	0.02	—	—	—	—
	Well	59.97	2.14	30.66	10.17	6.22	5.03	2.64	3.11	—
Davis 325,760	All classes	100.0	22.9	22.3	20.0	16.1	9.2	4.6	1.6	3.3
	Poor	21.87	18.75	3.12	—	—	—	—	—	—
	Imperfect	43.15	0.31	18.48	16.57	7.79	—	—	—	—
	Well	31.69	3.92	0.66	3.43	8.26	9.23	4.61	1.58	—
Decatur 339,196	All classes	100.0	11.9	17.6	20.7	18.9	20.4	6.6	1.4	2.5
	Poor	12.60	8.06	4.47	0.07	—	—	—	—	—
	Imperfect	24.61	—	11.81	12.40	0.40	—	—	—	—
	Well	60.24	3.82	1.32	8.19	18.46	20.45	6.65	1.35	—
Delaware 366,720	All classes	100.0	18.2	46.6	15.9	6.7	4.0	1.7	3.4	3.5
	Poor	7.27	4.21	3.06	—	—	—	—	—	—
	Imperfect	20.41	7.28	13.11	0.02	—	—	—	—	—
	Well	68.86	6.75	30.47	15.88	6.72	3.95	1.69	3.40	—
Des Moines 270,723	All classes	100.0	33.0	26.7	16.9	6.8	4.4	2.5	3.9	5.8
	Poor	17.63	16.49	1.14	—	—	—	—	—	—
	Imperfect	15.69	12.04	3.35	0.30	—	—	—	—	—
	Well	60.77	4.45	22.21	16.64	6.81	4.40	2.50	3.86	—
Dickinson 255,806	All classes	100.0	32.8	38.4	15.9	3.6	2.3	0.9	0.4	5.7
	Poor	22.91	20.77	2.14	—	—	—	—	—	—
	Imperfect	22.25	8.99	13.23	0.03	—	—	—	—	—
	Well	49.20	3.07	23.07	15.87	3.61	2.31	0.85	0.43	—

19701

DISTRIBUTION OF SOILS

67



Table 1. Continued

County and Total Acres in County	Natural Drainage	Percent Totals	% of County in Respective Slope Classes							Excluded Area
			0-1%	2-4%	5-8%	9-13%	14-17%	18-24%	25+%	
Dubuque 398,083	All classes	100.0	3.5	12.2	18.1	31.5	18.0	4.7	6.3	5.7
	Poor	1.22	0.73	0.41	—	—	—	—	—	—
	Imperfect	2.24	0.41	1.54	—	—	—	—	—	—
	Well	92.76	2.41	8.33	18.10	31.54	17.99	4.65	6.34	—
Emmet 257,280	All classes	100.0	36.2	31.5	19.6	6.0	1.8	3.8	0.1	1.0
	Poor	28.03	25.87	4.52	—	—	—	—	—	—
	Imperfect	17.72	9.60	9.67	—	—	—	—	—	—
	Well	53.31	3.73	20.19	19.60	6.53	1.80	3.80	0.17	—
Fayette 465,924	All classes	100.0	12.9	53.5	13.5	6.6	6.0	3.0	3.3	1.2
	Poor	12.58	2.25	10.33	—	—	—	—	—	—
	Imperfect	11.11	4.52	6.24	0.35	—	—	—	—	—
	Well	75.05	6.14	36.87	13.14	6.60	6.03	2.96	3.29	—
Floyd 321,920	All classes	100.0	48.0	42.8	4.3	0.4	0.5	0.1	—	3.9
	Poor	29.02	24.50	4.46	0.06	—	—	—	—	—
	Imperfect	35.16	15.38	19.04	0.74	—	—	—	—	—
	Well	32.00	8.24	19.38	3.46	0.40	0.47	0.05	—	—
Franklin 375,040	All classes	100.0	41.2	42.2	10.7	2.4	0.5	0.3	0.1	2.6
	Poor	30.78	29.33	1.45	—	—	—	—	—	—
	Imperfect	22.00	7.49	14.22	0.29	—	—	—	—	—
	Well	44.66	4.37	26.54	10.42	2.40	0.53	0.33	0.07	—
Fremont 333,440	All classes	100.0	38.7	19.3	9.5	20.6	2.9	2.5	2.2	4.3
	Poor	12.52	12.52	—	—	—	—	—	—	—
	Imperfect	11.21	6.37	4.17	0.09	0.58	—	—	—	—
	Well	71.93	19.83	15.11	9.38	20.05	2.94	2.45	2.17	—

Runge et al.: Distribution of Soils by Natural Drainage Class and by Slope Clas

Table 1. Continued

County and Total Acres in County	Natural Drainage	Percent Totals	% of County in Respective Slope Classes							Excluded Area
			0-1%	2-4%	5-8%	9-13%	14-17%	18-24%	25+%	
Greene 364,160	All classes	100.0	42.8	41.6	7.1	1.3	0.7	0.1	2.4	4.0
	Poor	39.75	39.75	—	—	—	—	—	—	—
	Imperfect	19.05	2.80	16.25	—	—	—	—	—	—
	Well	37.22	0.27	25.38	7.15	1.26	0.74	0.04	2.38	—
Grundy 320,631	All classes	100.0	35.0	48.8	13.1	0.8	0.1	—	—	2.2
	Poor	24.99	24.96	0.03	—	—	—	—	—	—
	Imperfect	27.51	8.53	18.91	0.07	—	—	—	—	—
	Well	45.27	1.49	29.85	13.03	0.82	0.08	—	—	—
Guthrie 381,441	All classes	100.0	19.1	26.7	18.5	17.3	9.7	2.6	3.9	2.2
	Poor	12.77	10.62	2.07	0.04	0.04	—	—	—	—
	Imperfect	18.11	7.72	8.30	0.07	2.02	—	—	—	—
	Well	66.88	0.74	16.32	18.37	15.24	9.67	2.54	4.00	—
Hamilton 369,882	All classes	100.0	58.2	33.1	4.4	1.1	0.4	0.4	2.4	0.1
	Poor	50.97	48.99	1.98	—	—	—	—	—	—
	Imperfect	24.33	9.13	15.20	—	—	—	—	—	—
	Well	24.67	0.05	15.95	4.41	1.14	0.37	0.37	2.36	—
Hancock 366,105	All classes	100.0	59.1	27.3	8.3	1.8	1.4	0.4	0.4	1.3
	Poor	48.26	43.86	4.40	—	—	—	—	—	—
	Imperfect	20.22	10.92	9.29	0.01	—	—	—	—	—
	Well	30.24	4.35	13.60	8.31	1.83	1.35	0.40	0.43	—
Hardin 367,360	All classes	100.0	37.4	38.3	14.9	3.2	0.7	0.6	0.1	4.8
	Poor	30.80	30.58	0.22	—	—	—	—	—	—
	Imperfect	17.24	5.05	12.19	—	—	—	—	—	—
	Well	47.26	1.74	25.90	14.94	3.25	0.65	0.60	0.09	—

1970]

DISTRIBUTION OF SOILS

69

Table 1. Continued

County and Total Acres in County	Natural Drainage	Percent Totals	% of County in Respective Slope Classes							Excluded Area
			0-1%	2-4%	5-8%	9-13%	14-17%	18-24%	25+%	
Harrison 447,376	All classes	100.0	33.1	13.4	13.6	11.4	10.7	7.6	8.1	2.1
	Poor	9.69	9.36	0.33	—	—	—	—	—	—
	Imperfect	7.52	5.77	1.17	0.58	—	—	—	—	—
	Well	80.72	17.99	11.90	13.07	11.39	10.70	7.57	8.10	—
Henry 281,604	All classes	100.0	25.8	26.8	23.5	10.0	6.0	2.7	0.6	4.6
	Poor	22.59	21.57	1.02	—	—	—	—	—	—
	Imperfect	23.92	3.95	9.57	7.58	2.82	—	—	—	—
	Well	48.89	0.31	16.16	15.90	7.19	6.03	2.65	0.65	—
Howard 301,438	All classes	100.0	36.1	53.7	6.6	0.8	0.1	—	0.1	2.6
	Poor	30.77	21.16	9.61	—	—	—	—	—	—
	Imperfect	36.69	9.84	24.82	1.92	0.07	—	—	0.04	—
	Well	29.87	5.09	19.26	4.65	0.73	0.14	—	—	—
Humboldt 378,404	All classes	100.0	52.4	39.3	2.5	0.6	0.6	—	0.8	3.8
	Poor	24.76	24.61	0.15	—	—	—	—	—	—
	Imperfect	20.80	1.28	19.52	—	—	—	—	—	—
	Well	50.66	26.47	19.63	2.50	0.64	0.60	—	0.82	—
Ida 276,542	All classes	100.0	5.4	33.3	29.2	24.4	5.6	1.8	0.3	—
	Poor	5.78	3.86	1.92	—	—	—	—	—	—
	Imperfect	8.10	0.12	7.60	0.38	—	—	—	—	—
	Well	86.11	1.42	23.78	28.85	24.40	5.56	1.76	0.33	—
Iowa 373,760	All classes	100.0	23.7	20.6	15.1	21.0	10.4	3.9	2.5	2.8
	Poor	7.41	7.32	—	—	—	0.03	0.06	—	—
	Imperfect	21.96	9.64	9.60	—	1.34	1.38	—	—	—
	Well	67.86	6.78	10.96	15.09	19.70	9.02	3.82	2.54	—

Runge et al.: Distribution of Soils by Natural Drainage Class and by Slope Clas

1970]

DISTRIBUTION OF SOILS

71

Table 1. Continued

County and Total Acres in County	Natural Drainage	Percent Totals	% of County in Respective Slope Classes							Excluded Area
			0-1%	2-4%	5-8%	9-13%	14-17%	18-24%	25+%	
Jackson 414,120	All classes	100.0	8.2	8.7	14.8	29.8	17.9	8.3	11.9	0.4
	Poor	1.41	1.06	0.32	0.03	—	—	—	—	—
	Imperfect	5.59	0.87	4.60	0.09	0.03	—	—	—	—
	Well	92.65	6.33	3.82	14.63	29.80	17.93	8.29	11.85	—
Jasper 471,040	All classes	100.0	20.3	25.7	20.7	20.6	9.1	2.2	0.4	1.0
	Poor	12.42	10.40	1.54	—	0.19	0.29	—	—	—
	Imperfect	30.12	6.37	14.72	7.35	1.68	—	—	—	—
	Well	56.49	3.48	9.42	13.39	18.71	8.86	2.24	0.39	—
Jefferson 279,044	All classes	100.0	25.0	22.2	25.0	9.9	9.5	5.4	0.2	2.8
	Poor	21.09	21.09	—	—	—	—	—	—	—
	Imperfect	33.50	—	13.69	17.15	2.57	0.09	—	—	—
	Well	42.59	3.90	8.47	7.84	7.30	9.44	5.41	0.23	—
Johnson 396,800	All classes	100.0	23.0	25.4	16.1	15.0	6.6	2.2	2.2	9.5
	Poor	7.51	6.98	0.49	—	—	0.04	—	—	—
	Imperfect	17.71	10.56	5.98	0.01	1.02	0.06	0.08	—	—
	Well	65.34	5.45	18.92	16.07	14.03	6.50	2.12	2.25	—
Jones 375,308	All classes	100.0	12.4	33.3	21.4	12.9	6.7	7.5	2.2	3.6
	Poor	5.52	3.49	2.03	—	—	—	—	—	—
	Imperfect	14.09	6.38	7.71	—	—	—	—	—	—
	Well	76.86	2.57	23.55	21.37	12.94	6.76	7.46	2.21	—
Keokuk 370,582	All classes	100.0	25.5	29.7	22.3	17.9	3.8	0.8	—	—
	Poor	1.67	1.57	0.10	—	—	—	—	—	—
	Imperfect	20.49	15.93	4.47	—	0.09	—	—	—	—
	Well	77.84	8.04	25.13	22.28	17.83	3.75	0.81	—	—

Table 1. Continued

County and Total Acres in County	Natural Drainage	Percent Totals	% of County in Respective Slope Classes							Excluded Area
			0-1%	2-4%	5-8%	9-13%	14-17%	18-24%	25+%	
Kossuth 626,557	All classes	100.0	35.4	50.5	8.4	1.0	0.3	0.5	0.2	3.7
	Poor	56.44	34.68	21.76	—	—	—	—	—	—
	Imperfect	13.00	0.38	12.62	—	—	—	—	—	—
	Well	26.81	0.29	16.11	8.37	0.98	0.26	0.55	0.25	—
Lee 348,800	All classes	100.0	19.3	33.3	15.9	8.5	2.8	2.7	8.4	9.1
	Poor	17.35	15.61	1.18	0.26	0.12	0.14	0.02	—	—
	Imperfect	43.08	0.45	20.28	14.41	7.94	—	—	—	—
	Well	30.47	3.25	11.84	1.21	0.45	2.72	2.64	8.38	—
Linn 458,883	All classes	100.0	15.0	42.8	21.3	4.3	2.4	1.4	1.6	11.2
	Poor	7.79	4.00	3.79	—	—	—	—	—	—
	Imperfect	19.61	8.02	11.55	0.04	—	—	—	—	—
	Well	63.38	2.99	27.43	21.31	4.26	2.43	1.40	1.56	—
Louisa 267,521	All classes	100.0	38.2	26.1	11.4	5.4	5.5	5.4	3.6	4.4
	Poor	20.89	20.12	0.64	0.13	—	—	—	—	—
	Imperfect	10.86	9.31	1.40	0.15	—	—	—	—	—
	Well	63.81	8.77	24.09	11.06	5.44	5.53	5.36	3.56	—
Lucas 277,760	All classes	100.0	15.0	28.1	18.7	10.4	9.4	14.9	3.5	—
	Poor	12.7	11.9	0.8	—	—	—	—	—	—
	Imperfect	32.5	2.1	17.7	12.7	—	—	—	—	—
	Well	54.8	1.0	9.6	6.0	10.4	9.4	14.9	3.5	—
Lyon 376,326	All classes	100.0	24.4	51.6	17.1	2.8	0.6	0.5	2.0	1.0
	Poor	10.27	9.76	0.56	—	—	—	—	—	—
	Imperfect	13.18	6.26	6.92	—	—	—	—	—	—
	Well	75.43	8.39	44.06	17.09	2.80	0.60	0.53	1.96	—

Runge et al.: Distribution of Soils by Natural Drainage Class and by Slope Clas

Table 1. Continued

County and Total Acres in County	Natural Drainage	Percent Totals	% of County in Respective Slope Classes							Excluded Area
			0-1%	2-4%	5-8%	9-13%	14-17%	18-24%	25+%	
Madison 361,596	All classes	100.0	24.4	19.8	22.2	10.5	7.0	7.5	2.2	6.4
	Poor	12.09	9.41	2.55	0.13	—	—	—	—	—
	Imperfect	16.00	9.01	2.97	2.64	1.38	—	—	—	—
	Well	65.53	5.96	14.28	19.41	9.13	7.04	7.46	2.25	—
Mahaska 366,073	All classes	100.0	16.9	26.3	27.4	19.9	3.8	1.1	—	4.6
	Poor	16.18	14.50	1.65	—	—	—	—	—	—
	Imperfect	13.54	1.69	9.59	1.86	0.45	—	—	—	—
	Well	65.50	0.75	15.00	25.43	19.44	3.85	1.13	—	—
Marion 364,805	All classes	100.0	15.9	20.7	27.1	21.4	7.3	2.1	0.2	5.3
	Poor	10.56	7.23	2.91	0.39	0.03	—	—	—	—
	Imperfect	26.39	2.31	13.45	9.48	1.15	—	—	—	—
	Well	57.77	6.38	4.34	17.21	20.26	7.35	2.06	0.17	—
Marshall 367,341	All classes	100.0	22.6	29.4	29.6	14.1	3.8	0.4	0.1	0.1
	Poor	8.14	6.90	1.24	—	—	—	—	—	—
	Imperfect	12.47	10.94	7.53	—	—	—	—	—	—
	Well	73.36	4.74	20.65	29.55	14.14	3.82	0.37	0.09	—
Mills 285,440	All classes	100.0	18.0	25.1	18.4	20.2	8.6	3.8	1.6	4.3
	Poor	8.23	7.86	0.37	—	—	—	—	—	—
	Imperfect	9.41	1.95	7.15	—	0.13	0.18	—	—	—
	Well	78.04	8.20	17.53	18.43	20.07	8.38	3.84	1.59	—
Mitchell 298,880	All classes	100.0	61.1	30.5	2.3	1.2	0.4	0.3	0.3	3.9
	Poor	20.30	20.07	0.17	0.06	—	—	—	—	—
	Imperfect	28.45	22.09	5.99	0.37	—	—	—	—	—
	Well	47.35	18.92	24.31	1.84	1.24	0.35	0.34	0.35	—

Table 1. Continued

County and Total Acres in County	Natural Drainage	Percent Totals	% of County in Respective Slope Classes							Excluded Area
			0-1%	2-4%	5-8%	9-13%	14-17%	18-24%	25+%	
Monona 454,398	All classes	100.0	46.9	8.5	7.8	7.9	11.7	5.3	9.7	2.2
	Poor	22.25	22.25	—	—	—	—	—	—	—
	Imperfect	10.39	10.39	—	—	—	—	—	—	—
	Well	65.07	14.29	8.49	7.80	7.85	11.65	5.34	9.65	—
Monroe 278,402	All classes	100.0	10.9	18.4	24.2	18.5	11.0	10.3	4.4	2.3
	Poor	12.09	7.15	4.55	0.30	0.09	—	—	—	—
	Imperfect	34.68	1.58	11.03	19.44	2.63	—	—	—	—
	Well	51.03	2.18	2.82	4.47	15.78	10.98	10.35	4.45	—
Montgomery 270,080	All classes	100.0	19.2	31.8	12.9	28.5	2.1	0.9	0.5	4.1
	Poor	9.11	8.60	—	0.08	0.43	—	—	—	—
	Imperfect	17.61	2.04	12.28	0.26	2.99	0.04	—	—	—
	Well	69.21	8.53	19.56	12.59	25.13	2.06	0.86	0.48	—
Muscatine 287,964	All classes	100.0	42.4	32.3	12.4	5.7	3.0	1.5	2.5	0.2
	Poor	16.44	14.99	2.45	—	—	—	—	—	—
	Imperfect	10.95	9.27	1.54	0.14	—	—	—	—	—
	Well	71.43	18.16	28.28	12.26	5.72	2.97	1.50	2.54	—
O'Brien 368,260	All classes	100.0	30.8	61.0	4.0	0.8	0.1	0.6	1.0	1.7
	Poor	14.15	13.17	0.98	—	—	—	—	—	—
	Imperfect	24.64	9.27	15.01	0.19	0.17	—	—	—	—
	Well	59.51	8.35	44.98	3.79	0.63	0.14	0.64	0.98	—
Osceola 254,705	All classes	100.0	38.7	47.8	8.9	1.6	0.7	0.5	0.1	1.7
	Poor	16.70	16.60	0.10	—	—	—	—	—	—
	Imperfect	26.55	16.28	10.15	0.12	—	—	—	—	—
	Well	55.05	5.84	37.54	8.79	1.57	0.70	0.55	0.5	—

Runge et al.: Distribution of Soils by Natural Drainage Class and by Slope Clas

Table 1. Continued

County and Total Acres in County	Natural Drainage	Percent Totals	% of County in Respective Slope Classes							Excluded Area
			0-1%	2-4%	5-8%	9-13%	14-17%	18-24%	25+%	
Page 342,400	All classes	100.0	21.8	25.5	15.7	29.9	2.1	0.3	0.4	4.3
	Poor	11.13	10.95	0.01	0.07	0.10	—	—	—	—
	Imperfect	25.96	4.22	15.68	0.76	5.25	0.05	—	—	—
	Well	58.60	6.59	9.76	14.91	24.55	2.03	0.33	0.43	—
Palo Alto 362,894	All classes	100.0	51.5	34.6	10.9	2.2	0.5	0.2	0.1	—
	Poor	40.14	36.56	3.58	—	—	—	—	—	—
	Imperfect	16.97	7.48	9.47	0.02	—	—	—	—	—
	Well	42.88	7.46	21.53	10.87	2.21	0.53	0.20	0.08	—
Plymouth 552,344	All classes	100.0	11.8	31.7	35.0	11.5	3.9	2.7	2.3	1.1
	Poor	8.50	7.67	0.83	—	—	—	—	—	—
	Imperfect	7.50	0.15	6.85	0.34	0.16	—	—	—	—
	Well	82.86	3.96	24.05	34.62	11.32	3.88	2.68	2.35	—
Pocahontas 371,843	All classes	100.0	67.0	25.2	4.9	0.8	0.3	0.3	0.3	1.2
	Poor	52.66	52.45	0.21	—	—	—	—	—	—
	Imperfect	20.54	12.74	7.80	—	—	—	—	—	—
	Well	25.57	1.83	17.18	4.89	0.75	0.28	0.33	0.31	—
Polk 380,154	All classes	100.0	30.0	34.3	12.0	5.8	3.2	1.5	0.3	12.9
	Poor	18.37	17.37	1.00	—	—	—	—	—	—
	Imperfect	22.13	8.98	11.59	—	0.06	—	1.50	—	—
	Well	46.55	3.64	21.67	12.03	5.72	3.23	—	0.26	—
E. Pottawattamie 299,520	All classes	100.0	13.7	33.8	12.6	29.2	5.6	1.5	0.9	2.7
	Poor	7.11	6.51	0.01	0.02	0.43	0.14	—	—	—
	Imperfect	11.61	0.64	10.03	0.68	0.16	0.10	—	—	—
	Well	78.61	6.60	23.77	11.87	28.57	5.41	1.47	.92	—

1970]

DISTRIBUTION OF SOILS

75



Table 1. Continued

County and Total Acres in County	Natural Drainage	Percent Totals	% of County in Respective Slope Classes							Excluded Area
			0-1%	2-4%	5-8%	9-13%	14-17%	18-24%	25+%	
W. Pottawattamie 330,240	All classes	100.0	23.2	23.6	11.2	17.9	10.7	4.1	2.0	7.3
	Poor	4.45	4.45	—	—	—	—	—	—	—
	Imperfect	1.70	1.52	0.18	—	—	—	—	—	—
	Well	86.53	17.25	23.43	11.18	17.90	10.65	4.12	2.00	—
Poweshiek 376,960	All classes	100.0	16.9	27.7	22.0	16.8	9.4	2.9	0.2	4.1
	Poor	8.11	7.35	0.18	0.05	0.40	0.13	—	—	—
	Imperfect	27.66	6.75	13.75	0.12	6.09	0.95	—	—	—
	Well	60.11	2.78	13.73	21.86	10.32	8.34	2.85	0.23	—
Ringgold 344,320	All classes	100.0	5.3	23.3	30.3	24.2	11.5	1.1	0.1	4.3
	Poor	10.22	3.98	3.36	2.68	0.20	—	—	—	—
	Imperfect	52.40	0.58	17.78	24.23	9.81	—	—	—	—
	Well	33.05	0.77	2.15	3.37	14.14	11.54	1.06	0.02	—
Sac 371,032	All classes	100.0	28.6	50.3	13.7	4.1	1.3	0.5	0.6	0.9
	Poor	20.73	19.83	0.90	—	—	—	—	—	—
	Imperfect	16.75	6.55	10.10	0.04	—	—	0.06	—	—
	Well	61.61	2.26	39.27	13.62	4.13	1.27	0.42	0.64	—
Scott 289,920	All classes	100.0	23.3	34.5	17.5	6.2	3.5	4.0	1.5	9.5
	Poor	7.31	7.31	—	—	—	—	—	—	—
	Imperfect	20.01	10.43	9.58	—	—	—	—	—	—
	Well	63.16	5.52	24.89	17.53	6.17	3.51	4.04	1.50	—
Shelby 375,683	All classes	100.0	10.0	30.7	7.8	35.0	11.3	2.3	0.3	2.6
	Poor	2.81	2.76	—	—	0.03	0.02	—	—	—
	Imperfect	22.72	1.38	20.75	0.32	0.17	0.03	0.07	—	—
	Well	71.86	5.87	9.93	7.50	34.75	11.22	2.26	0.33	—

Runge et al.: Distribution of Soils by Natural Drainage Class and by Slope Clas

Table 1. Continued

County and Total Acres in County	Natural Drainage	Percent Totals	% of County in Respective Slope Classes							Excluded Area
			0-1%	2-4%	5-8%	9-13%	14-17%	18-24%	25+%	
Sioux 484,790	All classes	100.0	19.6	49.2	22.2	4.3	0.3	0.1	1.0	3.3
	Poor	7.16	6.41	0.74	0.01	—	—	—	—	—
	Imperfect	11.88	5.43	6.45	—	—	—	—	—	—
	Well	77.73	7.78	41.98	22.21	4.34	0.31	0.07	1.04	—
Story 363,520	All classes	100.0	39.0	38.3	9.4	3.2	2.0	1.6	0.4	6.1
	Poor	38.13	35.06	3.07	—	—	—	—	—	—
	Imperfect	14.46	1.79	12.58	—	—	—	0.09	—	—
	Well	41.36	2.11	22.67	9.40	3.19	1.99	1.56	0.44	—
Tama 460,800	All classes	100.0	43.9	31.3	—	10.2	6.3	5.6	—	2.7
	Poor	5.29	5.29	—	—	—	—	—	—	—
	Imperfect	6.53	5.93	0.35	—	0.25	—	—	—	—
	Well	85.56	32.64	30.96	—	10.00	6.31	5.65	—	—
Taylor 337,919	All classes	100.0	21.6	22.5	22.7	27.2	2.1	0.7	0.3	2.9
	Poor	19.58	9.22	0.65	9.35	0.36	—	—	—	—
	Imperfect	12.82	10.04	0.51	2.27	—	—	—	—	—
	Well	64.70	2.35	21.33	11.06	26.85	2.08	0.72	0.31	—
Union 272,640	All classes	100.0	12.1	26.0	25.9	19.8	7.4	2.9	1.1	4.8
	Poor	10.41	5.80	4.31	0.30	—	—	—	—	—
	Imperfect	19.55	2.88	11.51	5.10	—	0.06	—	—	—
	Well	65.23	3.36	10.19	20.53	19.79	7.32	2.91	1.13	—
Van Buren 313,597	All classes	100.0	17.5	16.2	18.4	12.3	10.5	16.0	6.2	2.9
	Poor	14.01	13.04	0.96	0.01	—	—	—	—	—
	Imperfect	38.74	0.28	14.37	17.08	6.99	0.02	—	—	—
	Well	44.34	4.19	0.91	1.26	5.34	10.45	15.98	6.21	—

1970]

DISTRIBUTION OF SOILS

77

Table 1. Continued

County and Total Acres in County	Natural Drainage	Percent Totals	% of County in Respective Slope Classes							Excluded Area
			0-1%	2-4%	5-8%	9-13%	14-17%	18-24%	25+%	
Wapello 280,327	All classes	100.0	25.4	27.4	20.8	10.6	6.7	3.8	3.3	2.0
	Poor	17.09	14.26	2.83	—	—	—	—	—	—
	Imperfect	36.09	1.17	16.03	16.33	2.56	—	—	—	—
	Well	44.81	9.96	8.54	4.46	8.02	6.71	3.76	3.34	—
Warren 364,820	All classes	100.0	18.6	28.0	23.9	15.2	8.2	5.3	0.5	0.3
	Poor	18.33	13.13	5.20	—	—	—	—	—	—
	Imperfect	21.84	4.03	10.56	7.07	0.13	0.05	—	—	—
	Well	59.46	1.44	12.20	16.86	15.05	8.16	5.28	0.47	—
Washington 363,493	All classes	100.0	30.7	27.7	21.9	14.8	3.2	0.9	0.8	—
	Poor	24.49	24.23	0.24	0.02	—	—	—	—	—
	Imperfect	6.64	1.01	5.59	0.04	—	—	—	—	—
	Well	68.88	5.47	21.85	21.84	14.76	3.25	0.93	0.78	—
Wayne 340,477	All classes	100.0	11.3	27.7	29.2	18.3	10.5	0.5	0.1	2.5
	Poor	15.17	10.69	4.27	0.21	—	—	—	—	—
	Imperfect	30.06	—	3.81	17.83	8.42	—	—	—	—
	Well	52.36	0.58	19.66	11.20	9.86	10.49	0.54	0.03	—
Webster 459,519	All classes	100.0	47.4	41.5	2.9	0.7	0.7	0.7	2.7	3.4
	Poor	43.39	40.24	3.10	—	0.05	—	—	—	—
	Imperfect	24.60	3.08	21.46	0.06	—	—	—	—	—
	Well	28.61	4.13	16.91	2.80	0.69	0.67	0.74	2.67	—
Winnebago 257,280	All classes	100.0	43.9	33.1	16.1	5.1	1.7	0.1	0.1	—
	Poor	45.64	39.93	5.71	—	—	—	—	—	—
	Imperfect	15.67	3.65	11.98	0.04	—	—	—	—	—
	Well	38.69	0.28	15.44	16.04	5.12	1.68	0.09	0.04	—

Runge et al.: Distribution of Soils by Natural Drainage Class and by Slope Clas

Table 1. Continued

County and Total Acres in County	Natural Drainage	Percent Totals	% of County in Respective Slope Classes							Excluded Area
			0-1%	2-4%	5-8%	9-13%	14-17%	18-24%	25+%	
Wineshiek 440,284	All classes	100.0	5.6	27.9	26.4	20.6	7.8	5.1	6.5	0.1
	Poor	6.46	2.33	4.07	0.06	—	—	—	—	—
	Imperfect	8.37	1.97	6.02	0.38	—	—	—	—	—
	Well	85.12	1.35	17.84	25.96	20.61	7.81	5.19	6.45	—
Woodbury 561,291	All classes	100.0	28.1	10.2	18.1	20.8	13.3	3.2	1.4	4.9
	Poor	10.27	10.27	—	—	—	—	—	—	—
	Imperfect	5.39	3.44	1.35	0.60	—	—	—	—	—
	Well	79.48	14.40	8.83	17.49	20.82	13.29	3.22	1.43	—
Worth 256,640	All classes	100.0	64.0	23.5	5.6	1.7	1.3	0.1	0.1	3.8
	Poor	41.67	41.34	0.33	—	—	—	—	—	—
	Imperfect	19.73	15.09	4.57	0.07	—	—	—	—	—
	Well	34.83	7.59	18.65	5.50	1.65	1.26	0.14	0.04	—
Wright 369,280	All classes	100.0	55.6	31.4	5.8	1.6	0.9	0.2	0.1	4.4
	Poor	51.65	50.60	1.05	—	—	—	—	—	—
	Imperfect	20.28	2.30	17.89	0.09	—	—	—	—	—
	Well	23.76	2.72	12.46	5.73	1.64	0.92	0.21	0.08	—

1970]

DISTRIBUTION OF SOILS

79

are 3 or 4, then chroma are typically 1 or less. Exceptions occur, however, and then observational experience and other morphological features of the soil are used. For example, in soils as the alluvial Wabash series (Oschwald et al. 1965), the subsoil color is black to very dark gray and obscures the olive gray colors diagnostic of poor drainage. Then the combination of 0 to 1 chroma and clay textures in the surface and subsoil layers, as well as the commonly observed water saturation of the subsoil, were used to place the Wabash series in the poorly drained drainage class.

The soils identified as being imperfectly drained generally have mottled brown and gray or partly gleyed subsoil horizons. Mottled subsoil horizons have Munsell hues of 10YR to 5Y with values between 4 and 7 and chroma of 2 or higher. If chroma are between 3 and 6, then mottles of 2 chroma or less are present. High and low chroma colors (yellow and brown to gray) are mixed randomly in the soil matrix. The percentage of yellow and brown to gray color is interpreted to be an indication of the natural soil drainage condition. Soils in Iowa having a high percentage of gray colors and a low percentage of brown colors are less well drained than soils having a low percentage of gray colors and a high percentage of brown colors.

Well drained soils in Iowa usually have yellow-brown or brown subsoil horizons.

Exceptions to the use of gray or mottled subsoil color for poor and imperfect drainage, respectively, have been noted by Ruhe et al. (1955) and Dideriksen (1966). They noted that some well drained soils have gray or mottled subsoils. These soils are on sideslopes where geological erosion has exposed the gray or sometimes mottled materials. The gray materials are stratigraphically related to geochemical weathering zones in the parent materials. They are an indicator of drainage before geological erosion. The colors are commonly relict and are not related to the present natural soil drainage.

In general, poorly drained soils must be drained before they will produce high yields of crops such as corn. Drainage improves tillage and harvesting operations as well as rooting depth of crops. When the soil has an internal permeability of moderate or higher, drainage can be improved by placing tile below the frost line. Soils with very slow internal permeability do not drain rapidly enough for tile to function adequately. Examples of the latter soils are the Edina and Haig soils of southern Iowa and the Luton soils of the Missouri River bottom land (Figure 1; Oschwald et al. 1965). Drainage can be improved on these soils by surface ditches or land grading to a ditch or tile inlet.

Imperfectly drained soils on slightly convex slopes are usually




Figure 1. Principal Soil Association Areas of Iowa

not tilled in Iowa. However, in the Kenyon-Floyd-Clyde and Cresco-Lourdes-Clyde soil associations in northeastern Iowa (Fig. 1) the imperfectly drained soils are often tilled. In part, it is thought that farmers tile these soils so they may work their fields earlier in the spring and thereby increase the length of the growing season. Farmers with large equipment on large tracts of land want to till and plant the field without interference of a few slightly wet areas.

Well drained soils do not require any improvement in drainage. A more serious problem with the well drained soils is water erosion resulting from rapid runoff.

DISTRIBUTION OF SOILS BY DRAINAGE AND BY SLOPE CLASSES

In Figure 2 a summary is given of the distribution of soils by natural drainage class given in Table 1 for each of the counties in Iowa. The percentage of the soils in each of the three natural drainage classes for each county is proportional to the area of the county covered by the respective drainage class symbol. For example, the percentages of soils in Story County in the respective drainage classes are 38.1% as poorly, 14.5% as imperfectly, and 41.4% as well drained, and are shown as black, horizontal hachuring and unhachured, respectively. By contrast, the soils in Allamakee County are virtually all well drained. Kossuth County has the largest percentage of poorly drained soils of any county, about 56%.




Figure 2. Distribution of Soils by Drainage Class for Iowa counties

The percentage of the soils in the county occurring in each of the slope classes (0-1%, 2-4%, 5-8%, 9-13%, 14-17%, 18-24%, 25%+, and excluded area) is given in Table 1. These seven slope classes were further grouped into the three slope classes: 0-4.5%, 4.6-8.5%, and greater than 8.6% slope. The percentage of each of the soils in a county within each of these three slope classes is shown in Figure 3. Figures 2 and 3 are correlated; counties having




Figure 3. Distribution of Soils by Slope Class for Iowa counties

large percentages of well drained soils are the same counties with large percentages of soils having slopes greater than 8.6 percent. Some counties do have some poorly drained soils on steeper slopes. These are mainly in southern and southeastern Iowa where the underlying Kansan till clayey paleosol outcrops on the steep side-slopes.

Figure 3 graphically shows the average slope of individual counties and also of the state. Those counties with relatively low average slope occur in northwestern and north-central Iowa. Soils with the greatest average slope are in northeastern Iowa adjacent to the Mississippi River. The soils in Allamakee County have the following slope: 16.9 percent have slopes less than 4.5%, 17.1 percent have slopes between 4.6 and 8.5%, and 63.7 percent have slopes greater than 8.6%.

#### USE, MANAGEMENT, AND CONSERVATION CONSIDERATIONS

The purpose of this paper has been to present data on the natural drainage and slope distribution of soils of Iowa counties. Differences among Iowa counties in natural drainage and slopes of soil resources are evident. As a result, different use, management, and conservation considerations are to be expected in different counties. For example, Allamakee County has mostly well-drained, quite sloping soils, and water erosion control practices are an important conservation need.

In constrast, most north-central Iowa counties have a preponderance of poorly drained, level soils (Figs. 2, 3). On such soils, farming practices include tile and surface drainage. Fall plowing is also a common practice to aid in winter freezing and thawing to form a more granular structure for the seedbed and to aid drying of the surface soil for earlier planting of crops. Recent technology has resulted in almost continuous row crops with extensive use of nitrogen fertilizer. Wind erosion has been most common in soil areas that use fall plowing practices. This problem and some suggestions for its amelioration have been discussed by Moldenhauer and Duncan (1969).

Natural soil drainage also affects non-farming uses of soils. For example, poorly and imperfectly drained soils have seasonally water-saturated subsoils. This wetness severely limits their use for solid waste disposal (land fill) as there would be a potential hazard to contamination of the ground water. They are also severely limited for septic field use as the effluent could not discharge into a wet soil, and there would be a hazard of contamination of the water supplies recharged through water-saturated subsoils. Though most well drained soils of Iowa have fewer limitations for these uses, there are exceptions. For example, well drained soils with



porous limestone present in the subsoil may also have severe limitation because of potential contamination of water. For this kind of information, soil type must be considered. Soil type was mapped in the sample soil survey but is not presented in this paper.

#### DETAILED COUNTY SOIL SURVEYS

For more precise and county-wide information on soil, slope, and drainage than provided by the sample soil survey, detailed county soil survey reports should be consulted. The county soil reports also include soil type information. The more recent county soil survey reports, as that of Winneshiek County (Kittleson and Dideriksen, 1968), also include a wide range of interpretations on the suitability for and limitations of the different soils for various uses. The detailed county soil surveys are made cooperatively by the Soil Conservation Service of the U. S. Department of Agriculture and the Iowa Agriculture and Home Economics Experiment Station of Iowa State University. Detailed county soil surveys are made on aerial photograph base maps usually of 4-inch-to-the-mile scale. Publication is usually also on this or a slightly smaller scale. Sixteen detailed county soil surveys have been published, twelve are in one stage or another of publication, and soil mapping is in progress in an additional twenty-two counties. County Extension and County Soil Conservation District Offices, or the Publications Distribution Office of Iowa State University, Ames, may be contacted for information on availability of detailed county soil surveys.

#### SUMMARY

The distribution of soils by natural drainage class and by slope groups is given. Problems associated with drainage class and slope class are quantified for individual counties in Iowa. Various individuals and groups may find this information helpful in their programs. This information can be especially useful to individuals responsible for broad resource planning. County Extension Councils, Soil Conservation Districts, county and regional planning groups, and county supervisors are examples of individuals who may be able to use the individual county data in their program planning and decision making sessions.

#### Literature Cited

- ARNOLD, R. W., L. E. TYLER, & F. F. RIECKEN. 1960. *Iowa Acad. Sci. Proc.* 67: 260-267.
- DIDERIKSEN, RAYMOND I. 1966. *Soil landscape relationships of the Mahaska topo-sequence*. Unpublished M.S. thesis. Iowa State University Library, Ames.
- HAAN, C. T. & H. P. JOHNSON. 1967. *Iowa State J. Sci.* 42: 149-160.

- IOWA CONSERVATION NEEDS COMMITTEE. 1963. *Iowa soil and water conservation needs inventory*. Coop. Ext. Serv., Ames.
- KITTLESON, KENNETH K. & RAYMOND I. DIDERIKSEN. 1968. *Soil survey of Winneshiek County, Iowa*. U. S. Dept. Agr.
- MOLDENHAUER, W. C. & E. R. DUNCAN. 1969. *Principles and methods of wind-erosion control in Iowa*. Iowa Agr. and Home Econ. Exp. Sta. Spec. Rept. 62. Ames.
- OSCHWALD, W. R., F. F. RIECKEN, R. I. DIDERIKSEN, W. H. SCHOLTES, and F. W. SCHALLER. 1965. *Principal soils of Iowa*. Spec. Rept. 42.
- RUHE, R. V., R. C. PRILL and F. F. RIECKEN. 1955. *Soil Sci. Soc. Amer. Iowa Coop. Ext. Serv., Ames. Proc. 19*: 345-347
- RUNGE, E. C. A. & L. DE LEON. 1967. *Iowa Acad. Sci. Proc.* **67**: 232-236.
- SLUSHER, D. F., R. W. ARNOLD, & R. PROTZ. 1961. *Iowa Aca. Sci. Proc.* **68**: 380-396.