

1963

Trematodes of Passerine Birds from Chickasaw County, Iowa

Charles J. Ellis
State College of Iowa

Copyright ©1963 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Ellis, Charles J. (1963) "Trematodes of Passerine Birds from Chickasaw County, Iowa," *Proceedings of the Iowa Academy of Science*, 70(1), 486-492.

Available at: <https://scholarworks.uni.edu/pias/vol70/iss1/78>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

exist, certain of these might possibly be more resistant to in-activation than others.

Literature Cited

- Levin, A. J. 1940. *J. Parasit.* 26 (Suppl.): 31.
Mauss, E. A. 1940. *Am. J. Hyg., Sec. D*, 32: 80-83.
Offutt, E. P., Jr. 1941. *J. Parasit.* 27 (Suppl.): 24.
Oliver-Gonzalez, J. 1940. *Jour. Infect. Dis.* 67: 292-300.
Roth, H. 1941. *Acta. path. et microbiol. Scandinav.* 18: 160-167.

Trematodes of Passerine Birds from Chickasaw County, Iowa¹

CHARLES J. ELLIS¹

Abstract. One-hundred and twenty-five passerine birds were collected in Chickasaw County, Iowa, October, 1959, through August, 1960. Thirteen species of birds (19%) were infected with 14 genera of trematodes. The hosts, helminths and infection sites are tabulated.

Little is known concerning the trematode fauna of Iowa birds belonging to the order Passeriformes. Ward (1901) examined 33 Iowa passerines and reported only one bird harboring a trematode infection.

The present study concerns the trematodes of passerine birds collected October, 1959, through August, 1960, from the Goodale Conservation Area in northwest Chickasaw County, Iowa. This area encompasses approximately 21 acres and contains six major types of habitats: river, marsh, pond, thicket, deciduous woods and meadow.

Previous studies on trematodes of passerine birds indicate varying degrees of infection. In Czechoslovakia Ryšavy (1955a) studied 168 passerines (32 species) and found none of them infected with trematodes. Rankin (1946) as part of a larger helminth survey of birds and mammals in western Massachusetts examined five passerines and found none infected with trematodes. Sulgostowska (1958) examined 33 passerines in Poland and found two (6%) infected with trematodes. According to unpublished data from Iowa Lakeside Laboratory at Lake Okoboji, 161 passerine birds (27 species) have been examined and 59 (37%) were infected with trematodes. In the current study approximately 19% of the birds examined harbored adult trematodes.

Several studies on helminths of a single species of passerines

¹ State College of Iowa (Now at Iowa State University).

have been published. Thus, Boyd (1951) examined 300 starlings (*Sturnus vulgaris*) collected from various eastern and midwestern states and found only four infected with trematodes. Boyd, et al, (1956) found a 37% trematode infection rate in blue jays (*Cyanocitta cristata*) from New England. Ryšavy (1955b) examined 79 blackbirds (*Turdus merula*) and found 30% infected with trematodes. Daly (1959) in a study on the intestinal helminths of 103 crows (*Corvus brachyrhynchus paulus*) in Virginia found trematodes in only 6%. Joszt (1958) found 9.5% of the 190 house sparrows (*Passer domesticus*) from Warsaw, Poland, infected with trematodes.

Trematodes recovered during the present study were fixed in A.F.A., stained in Semichon's aceto-carmine and mounted in synthetic resin. Many were counterstained with fast green.

Table 1 summarizes the 125 birds examined, 24 of which (19%) harbored adult trematodes. Of these birds (representing 14 families and 38 species) those belonging to the family Fringillidae were the most abundant. Seventeen percent of them harbored adult trematodes.

Table 1. Avian Families Examined

	No. examined	No. Infected
Alaudidae (larks)	6	0
Corvidae (crows, jays)	4	2
Fringillidae (sparrows)	46	8
Hirundinidae (swallows)	1	1
Icteridae (meadowlarks, red-wing)	35	11
Mimidae (thrashers)	2	1
Paridae (chickadees)	6	0
Parulidae (warblers)	7	0
Sittidae (nuthatches)	3	0
Sturnidae (starlings)	1	0
Sylvidae (kinglets)	3	0
Troglodytidae (wrens)	2	0
Turdidae (thrushes)	1	0
Tyrannidae (flycatchers)	8	1
Totals	125	24

The lowest degree of infection was seen among members of the families Paridae, Parulidae, Sylvidae and Sittidae, none of which harbored trematodes. The small number examined, however, does not permit valid conclusions to be drawn regarding their ability to serve as definitive hosts.

The genera of trematodes recovered are listed in Table 2. Of these, those genera belonging to the family Dicrocoeliidae are the most abundant. A summary of hosts, helminths and sites is given in Table 3.

FAMILY DICROCOELIIDAE

Of all the trematode genera represented in the collection, those belonging to the family Dicrocoeliidae were most abund-

Table 2. Genera of Trematodes Recovered and Their Hosts

DICROCOELIIDAE	
<i>Brachylecithum</i>	(fox sparrow, brown thrasher, swamp sparrow)
<i>Conspicuum</i>	(common grackle, fox sparrow, eastern meadowlark)
<i>Lutztrema</i>	(eastern meadowlark, blue jay)
<i>Lyperosomum</i>	(brown thrasher)
<i>Zonorchis</i>	(fox sparrow)
ECHINOSTOMATIDAE	
<i>Echinostoma</i>	(common grackle, blue jay, eastern meadowlark, western meadowlark)
EUCOTYLIDAE	
<i>Tanaisia</i>	(song sparrow)
LECITHODENDRIIDAE	
<i>Ornithodendrium</i>	(eastern kingbird)
OPISTHORCHIIDAE	
<i>Opisthorchis</i>	(white-throated sparrow)
PLAGIORCHIIDAE	
<i>Plagiorchis</i>	(red-wing)
<i>Prosthogonimus</i>	(white-throated sparrow)
STRIGEIDAE	
<i>Strigea</i>	(rusty blackbird)
STOMYLOTREMATIDAE	
<i>Laterotrema</i>	(blue jay)
<i>Stomylotrema</i>	(barn swallow)

Table 3. Summary of Helminths Recovered

Host	Helminth	Site
Eastern kingbird (<i>Tyrannus tyrannus</i>)	<i>Ornithodendrium</i> sp.	Intestine
Barn swallow (<i>Hirundo rustica</i>)	<i>Stomylotrema</i> sp.	Intestine
Brown thrasher (<i>Toxostoma rufum</i>)	<i>Brachylecithum exochocotyle</i>	Liver
Blue jay (<i>Cyanocitta cristata</i>)	<i>Lyperosomum oswaldoi</i>	Liver
	<i>Lutztrema microstomum</i>	Liver
	<i>Echinostoma revolutum</i>	Intestine
	<i>Laterotrema</i> sp.	Intestine
Eastern meadowlark (<i>Sturnella magna</i>)	<i>Conspicuum icteridorum</i>	G. bladder
	<i>Brachylecithum</i> sp.	Liver
	<i>Lutztrema monenteron</i>	G. bladder
	<i>Echinostoma revolutum</i>	Intestine
	<i>Echinostoma</i> sp.	Intestine
Western meadowlark (<i>Sturnella neglecta</i>)	<i>Echinostoma revolutum</i>	Intestine
Red-wing (<i>Agelaius phoeniceus</i>)	<i>Plagiorchis noblei</i>	Liver
Rusty blackbird (<i>Euphagus carolinus</i>)	<i>Strigea</i> sp.	Intestine
Common grackle (<i>Quiscalus quiscula</i>)	<i>Conspicuum icteridorum</i>	G. bladder
	<i>Echinostoma</i> sp.	Intestine
White-throated sparrow (<i>Zonotrichia albicollis</i>)	<i>Opisthorchis</i> sp.	Intestine
	<i>Prosthogonimus</i> sp.	Intestine
Fox sparrow (<i>Passerella iliaca</i>)	<i>Brachylecithum nanum</i>	Liver
	<i>Conspicuum icteridorum</i>	G. bladder
	<i>Zonorchis</i> sp.	G. bladder
Swamp sparrow (<i>Melospiza georgiana</i>)	<i>Brachylecithum</i> sp.	Liver
Song sparrow (<i>Melospiza melodia</i>)	<i>Tanaisia zarudnyi</i>	Ureter
	<i>Zonorchis</i> sp.	G. bladder

ant. Five genera are represented: *Conspicuum*, *Lutztrema*, *Lyperosomum*, *Brachylecithum* and *Zonorchis*. Dicrocoeliids parasitize the gall bladder, liver, bile ducts and sometimes the pancreatic ducts of their hosts.

The only species of the genus *Conspicuum* recovered was *C. icteridorum* located within the gall bladder of the common grackle (*Quiscalus quiscula*), fox sparrow (*Passerella iliaca*) and the eastern meadowlark (*Sturnella magna*).

Lutztrema monenteron was recovered from the gall bladder of an eastern meadowlark. Varying dimensions are reported for this species by Price and McIntosh (1935), Denton and Byrd (1951) and by Mettrick (1958). My specimens agree more closely with those reported by Mettrick.

Lutztrema microstomum was recovered from the liver of a blue jay during this survey. Only one of four blue jays examined harbored this parasite which, according to Denton and Byrd (1951), is differentiated from *L. monenteron* by the former's narrower and smaller body, relatively larger and more anteriorly situated acetabulum, relatively shorter cecum, more tandem testes, greater distance between the acetabulum and the testes, and the pattern of its uterus in the pre-testicular region.

Three specimens of *Lyperosomum oswaldoi* were recovered from the bile duct of the brown thrasher (*Toxostoma rufum*). This host also was parasitized by *Brachylecithum exochocotyle*. Denton and Byrd (1951) also report a double infection of the brown thrasher with *L. oswaldoi* and *B. exochocotyle*.

More than 20 specimens of *Brachylecithum nanum* were recovered from the liver of a fox sparrow, but due to their fragility many were broken in removal.

The sixth dicrocoeliid species recovered was *Brachylecithum exochocotyle* mentioned previously from the bile duct of a brown thrasher. Denton and Byrd (1951) examined 41 of these hosts and found only one infected with this species of trematode.

One swamp sparrow (*Melospiza georgiana*) was infected with an unidentified species of *Brachylecithum*. The gall bladder and the bile ducts of two fox sparrows and one song sparrow (*Melospiza melodia*) were infected with members of another dicrocoeliid genus, *Zonorchis*.

FAMILY ECHINOSTOMATIDAE

Members of this family are noted for their apparent lack of host specificity both as adults and as larvae. These trematodes parasitize many kinds of vertebrates but particularly water birds. However, in this study, echinostomes were recovered from such birds as the eastern meadowlark, western meadow-

lark (*Sturnella neglecta*), common grackle and the blue jay, none of which is considered a water bird. In fact, all of these according to Bailey (1927) feed upon insects, grain and seeds. However, Pearson (1936) states that grackles also may eat salamanders, fish, crayfish and snails. The Goodale Conservation Area is marshy and includes numerous areas where birds could ingest those snails serving as intermediate hosts for the echinostome larvae.

All echinostomes found in this survey were taken from the intestine or cloaca. A double infection of *Echinostoma* sp. and *Lutztrema monenteron* was found in one eastern meadowlark.

Despite the fact that the genus *Echinostoma* seems to be one with a wide host specificity and a broad geographical range, the literature reveals few passerine hosts infected with these trematodes. Beaver (1937) refers to the species *E. revolutum* as being "as cosmopolitan in its choice of hosts as it is in geographical distribution. This lack of specificity in all of its parasitic stages is of course the most important reason for its wide distribution." Six of the 103 crows examined by Daly (1959) were infected with *E. revolutum*. Lumsden (1962) reports as a new host record *E. revolutum* from *Cassidix mesamexicanus* from Louisiana.

FAMILY EUCOTYLIDAE

As indicated in Table 2, this family is represented by only one species, *Tanaisia zarudnyi*. Two specimens were recovered from the ureters of the song sparrow which also harbored a *Zonorchis* infection. The identification of *T. zarudnyi* is based upon studies of Byrd and Denton (1950).

FAMILY LECITHODENDRIIDAE

Two specimens of the genus *Ornithodendrium* were recovered from the intestine of a single host, the eastern kingbird (*Tyrannus tyrannus*).

FAMILY OPISTHORCHIIDAE

This family is represented in this survey by one specimen of the genus *Opisthorchis* recovered from the cloaca of a white-throated sparrow (*Zonotrichia albicollis*). This infection was accompanied by a prosthoganimid in the terminal portion of the host's intestine.

The presence of an opisthorchid in a sparrow is difficult to explain since fish normally serve as second intermediate hosts for many members of this family.

FAMILY PLAGIORCHIIDAE

The same white-throated sparrow infected with *Opisthorchis* sp. was also infected with *Prosthogonimus* sp. The site of this

infection was the intestine despite the usual site for *Prosthogonimus* being in the oviduct or the bursa fabricius. Eight specimens of another member of this family, *Plagiorchis noblei*, were recovered from the cloaca of a red-winged blackbird (*Agelaius phoeniceus*).

FAMILY STOMYLOTREMATIDAE

One specimen of the genus *Stomylotrema* was recovered from the cloaca of a barn swallow (*Hirundo rustica*). Another stomylotrematid recovered belonged to the genus *Laterotrema* taken from the cloaca of a blue jay. However, its dimensions do not agree with those of previously described members of the genus.

FAMILY STRIGEIDAE

A single specimen belonging to the genus *Strigea* was recovered from the cloaca of a rusty blackbird (*Euphagus carolinus*).

The lack of uniformity shown by studies on trematode infections of passerines has already been noted. Such variation is undoubtedly due in a large measure to the food habits of the birds involved. Passerine birds inhabit a wide variety of terrestrial and semi-aquatic habitats. A more extensive trematode fauna is generally associated with semi-aquatic environments.

The present survey undertaken as a preliminary study on trematodes of Iowa passerines includes several species of avian hosts represented by only a single bird. Hence, any conclusions relative to dietary habits as indicative of degrees of trematode infection would be premature.

I am indebted to Dr. Martin L. Grant of the State College of Iowa for verifying many host identifications, to Dr. Martin J. Ulmer of Iowa State University for the trematode identifications and much other help and to the Chickasaw County Conservation Board for permission to do this study in the Goodale Conservation Area.

Literature Cited

- Bailey, Florence Merriam. 1927. Handbook of birds of the western United States. Houghton Mifflin Company, Boston and New York.
- Beaver, Paul Chester. 1937. Experimental studies on *Echinostoma revolutum* (Froelich) a fluke from birds and mammals. Ill. Biol. Mon. No. 15.
- Boyd, Elizabeth M. 1951. A survey of parasitism of the starling, *Sturnus vulgaris* L., in North America. J. Parasitol. 37:56-84.
- , Rosemarie L. Diminno and Carlita Nesslinger. 1956. Metazoan parasites of the blue jay *Cyanocitta cristata* L. J. Parasitol. 42:332-346.
- Byrd, E. E. and J. F. Denton. 1950. The helminth parasites of birds. I. A review of the trematode genus *Tanaisia* Skrjabin, 1924. Am. Midland Naturalist 43:32-57.
- Daly, E. F. 1959. A study of the intestinal helminths of the southern crow *Corvus brachyrhynchus paulus* in Virginia. Proc. Helminthol. Soc. Wash. 26:66.
- Denton, J. Fred and Elon E. Byrd. 1951. The helminth parasites of birds, III. Dicrocoeliid trematodes from North American birds. Proc.

- U. S. Nat. Mus. 101(3274):157-202.
- Joszt, I. 1958. Helminthofauna of a domestical (sic) sparrow (*Passer domesticus*) from Warsaw region. *Wiadomosci Parazytologiczne* 4 (5/6)690.
- Lumsden, Richard D. 1962. Four echinostome trematodes from Louisiana birds including the description of a new species. *Tulane Studies in Zoology* 9(5):301-308.
- Mettrick, David F. 1958. Helminth parasites of Hertfordshire birds I. *J. Helminthol.* 32 (1/2):49-64.
- Pearson, T. Gilbert. 1936. *Birds of America*. Garden City Books Garden City.
- Price, E. W. and Allen McIntosh. 1935. A new trematode, *Lyperosomum monenteron*, n. sp. (Dicrocoeliidae), from a robin. *Proc. Helminthol. Soc. Wash.* 2:63-64.
- Randkin, John Stewart. 1946. Helminth parasites of birds and mammals in western Massachusetts. *Am. Midland Naturalist* 35:756-768.
- Rysavy, Bohumil. 1955a. The parasitic worms of Passeriformes of the Lednice State Reservation. *Vestník Československé Zoologické Společnosti* 19(2):99-118 (Original not seen; abstracted in *Helminthological Abstracts* vol. 24, No. 960, 1955.)
- . 1955b. Motolice (Trematoda) u kosů černých (*Turdus merula* L.) v prazkém okolí (Die Saugwürmer (Trematoda) bei der Schwarzdrossel (*Turdus merula* L.) in der Prager Umgebung). *Zool. a Entomol. Listy*. Year 18, 4(3):271-274. (Original not seen; abstracted in *Helminthological Abstracts* vol. 24, No. 984, 1955.)
- Sulgotowska Teresa. 1958. Flukes of birds on Mamry Lake. *Wiadomosci Parazytologiczne* 4(5/6): 691-692.
- Ward, Henry B. 1901. Internal parasites of Nebraska birds. *Proc. Neb. Ornithol. Union* 63-70.

The Effect of Temperature Change on the Reaction Time of *Helisoma trivolvis* (Say)

PAUL K. GAUER¹

Abstract. A study of reaction speed, at different temperatures, of fresh-water snails (*Helisoma trivolvis*) is reported. Organic response to mechanical stimuli shows a definite relationship of reaction speed to the environmental temperature. There is an increase in rate of reaction from 40°F to 98°F; and a subsequent leveling off trend from 98°F to 110°F. Snail size has no appreciable effect upon reaction speed. Learning by experience is suggested.

The purpose of this research is to study the speed of reaction of the fresh-water snail in relation to environment. The hypothesis of this study is that environmental temperature has a direct effect on the speed of reaction of the fresh-water snail. The availability and convenient size of *Helisoma trivolvis* makes observation and experimentation with them practical.

Preliminary observation showed that the gastropods withdrew into their shells after being stimulated mechanically. In a short time they would again emerge from their shells; first the foot,

¹ Iowa Wesleyan College, Mt. Pleasant, Iowa.