2013

The Vascular Flora of Clay Prairie State Preserve (Butler County, Iowa): Recommendations to the Iowa State Preserve System

Edwin L. Freese

William R. Norris

Western New Mexico University

Let us know how access to this document benefits you

Copyright © Copyright 2013 by the Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/jias>

Part of the [Anthropology Commons](#), [Life Sciences Commons](#), [Physical Sciences and Mathematics Commons](#), and the [Science and Mathematics Education Commons](#)

Recommended Citation

Freese, Edwin L. and Norris, William R. (2013) "The Vascular Flora of Clay Prairie State Preserve (Butler County, Iowa): Recommendations to the Iowa State Preserve System," *Journal of the Iowa Academy of Science: JIAS*, 120(1-4), 21-34.

Available at: <https://scholarworks.uni.edu/jias/vol120/iss1/5>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Journal of the Iowa Academy of Science: JIAS by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

The Vascular Flora of Clay Prairie State Preserve (Butler County, Iowa): Recommendations to the Iowa State Preserve System

EDWIN L. FREESE¹ and WILLIAM R. NORRIS^{2*}

¹33493 S Ave., Adel, Iowa 50003

²Department of Natural Sciences, Western New Mexico University, PO Box 680, 1000 W. College Ave., Silver City, New Mexico 80062

Clay Prairie State Preserve (1.2 ha) is an upland prairie located in Butler County, IA, and represents one of the last remnants of black-soil tallgrass prairie in northeast Iowa. An annotated checklist of the vascular flora of Clay Prairie is presented here, based on a previously published flora of the preserve and recent field work conducted by the authors between 1997 to 2001, and 2008 to 2011. The preserve contains a diverse vascular flora representing 50 families, 145 genera and 214 taxa (174 native to Iowa). In comparison with 26 larger prairies (4 to 121 ha) protected in the Iowa state preserves system, the Clay Prairie flora ranks 10th in total number of native Iowa prairie plant taxa (135 taxa) and 10th in total number of native Iowa prairie plant taxa assigned high (7 to 10) values of the Iowa Conservation Coefficient (55 taxa). As a remnant of formerly much more extensive tallgrass prairie in Iowa, Clay Prairie protects individual plant taxa, plant assemblages, and ecological and evolutionary processes, while providing opportunities for people to experience native prairie. The floristic comparisons presented in this paper suggest that the floras of many other prairies protected in the Iowa State Preserves system are not thoroughly documented. Hence, we recommend that floristic studies of these other prairie preserves be undertaken to include documentation of the entire flora by voucher specimens. We also encourage site managers of prairies in the Iowa State Preserves system to prepare annual reports summarizing management and educational activities that occur in these preserves.

INDEX DESCRIPTORS: Clay Prairie State Preserve, cemetery, floristics, Iowa flora, prairie, Butler County, Iowa.

The loss and fragmentation of historic prairie vegetation, which once covered 85% of the state prior to settlement by Europeans but now occupies less than 0.1% of this state's land area, is well documented (Roosa 1978, Samson and Knopf 1994, Smith 1998). Remnant prairies occur all across the vast agricultural matrix that is Iowa. That small acreages of prairie were spared the plow at all is remarkable, and the circumstances responsible for the preservation of each prairie are unique. However, many prairie remnants in Iowa and elsewhere in the Midwest share a cultural history in that they owe their existence to their proximity to cemeteries (e.g., Betz 1989, Betz and Lamp 1992, Eilers 1966, Foster et al. 2010, Kerr and White 1981, Lantz 1969, Menges-Schaben 1998, Phillippe et al. 2010, Ruch et al. 2010, Schennum 1986, White 1978, 1988). These parcels of native grassland were thus afforded unintended but effective and long-term protection from conversion to crop fields or for other human use.

Clay Prairie State Preserve (Butler Co., IA), adjacent to the east border of Butler Center Cemetery, is one such prairie. Although a mere 1.2 ha in size (Herzberg and Pearson 2001), the vascular flora documented from Clay Prairie is remarkable in both its high diversity of prairie plant taxa and its floristic quality. In this paper, we present a checklist of the vascular flora of the prairie based on a previous study of Clay Prairie (Lantz 1969) as well as our own field work in the preserve (1997 to 2001; 2008 to 2013). Then, we compare the prairie flora of Clay Prairie to that of 26 other prairies which are protected as Iowa state preserves. We proceed to discuss the role of Clay Prairie as a natural vegetation remnant in conservation. Finally, we outline avenues of future

research that might be conducted on this and other small prairie remnants in Iowa.

STUDY SITE

Clay Prairie State Preserve (Butler Co., IA), is located at lat 42.69989° N and long 92.77171° W. It is rectangular in shape, and occupies 1.2 ha (3 ac) (Herzberg and Pearson 2001). The preserve is bounded by the cemetery proper on the west, blacktop on the north, gravel road to the east, and agricultural lands to the south (Fig. 1). Clay Prairie occurs about 0.80 km (0.5 mi) east of the historical town of Butler Center, which was abandoned in the late 19th century (Clay Prairie Preserve...[date unknown]) and today is a cornfield. The prairie was an unused part of Butler Center Cemetery (pers. comm. from W. V. Anthony to Kenneth Madden, Iowa State Preserves Board Ecologist, 1972) until 1961, when it was purchased by the University of Northern Iowa (UNI) with funds provided by the late agricultural equipment maker and UNI alumnus, Joseph B. Clay (Lantz 1969). Clay Prairie was dedicated as a biological state preserve on Dec. 28, 1976 (Roosa 1981), and is currently managed by the Biological Preserves Committee, Department of Biology, UNI (Herzberg and Pearson 2001).

Clay prairie occurs within a surface landform known as the Iowan Surface, formerly referred to as the Iowa Erosional Surface (Anderson 1998, Prior 1991). This landform is situated between the Des Moines Lobe to the west, noted for its bold landscape features (e.g., kettles, moraines) that mark its recent glacial history, and the Paleozoic Plateau to the east, recognized immediately by its rugged topography, frequently exposed Paleozoic sedimentary bedrock, and the near absence of glacial

* Corresponding author: William.Norris@wnmu.edu

Fig. 1. Aerial photograph (2011) of Clay Prairie State Preserve and surrounding landscape. Aerial imagery is from the National Agriculture Imagery Program (NAIP).

deposits (Prior 1991). In contrast, Prior (1991) describes the land surface of the Iowan Surface as “slightly inclined to gently rolling with long slopes, low relief, and open views to the horizon.” The terrain in the vicinity of Clay Prairie conforms to this description, being gently sloping (1 to 9%), and consisting of rolling hills and plains with elevation ranging from about 287 m to 317 m (940 to 1,040 ft) above sea level. The subdued landscape of the Iowan Surface, last visited by glaciers in pre-Illinoian times, reflects subsequent “episodes of weathering and soil development, erosion, and soil deposition” (Prior 1991).

The preserve is found on a north-facing hillside with elevations from 303 m (995 ft) in the northeast corner to 311 m (1,020 ft) in the southwest corner. Soils of this prairie remnant slope gradually down toward the northeast corner. Generally, the eastern edge is lowest and wettest from near the center to the northeast corner. The southwest and western half contain the driest soils. The soils are of the Kenyon-Clyde-Floyd Association (Buckner 1982, Web Soil Survey). About 90% of these soils in the preserve consist of Floyd Loam. The other approximately 10% of the preserve soils is classified as Kenyon Loam (part of northwest corner and a semicircular region in the south-central region of the preserve).

Climate for this region of Iowa is mesothermal moist subhumid (Thornwaite 1948). Climatic data available for Waterloo, IA (National Oceanic and Atmospheric Administration 1998), the nearest largest city to the preserve, approximate those for Clay Prairie State Preserve. Average yearly precipitation is about 841 mm (33.1 in). The average driest (21 mm [0.83 in]) and coldest (-10°C [14°F]) month is January, and the average wettest (119 mm [4.7 in]) and warmest (22.6°C [72.7°F]) month is July. The average frost-free growing season is from April 28 to October 4 (159 days).

Preserve Management

We compiled a partial record of management activities that have occurred at Clay Prairie State Preserve as documented in unpublished reports (Smith 1991, Camarata 1990, 1991, 1996) and personal communications. Although apparently never grazed (Smith 1991), Clay Prairie was hayed annually from an unspecified year through 1958 (Smith 1991) or until its purchase in 1961 (UNI Bio. Dept. website 2013). Smith (1991) reports that Clay Prairie was burned annually between 1976 and 1991, with half of the prairie burned every year, usually in late April. Additional prescribed burns of Clay Prairie occurred in 1996 (Camarata 1996), 2009, and at least twice since 2009 (pers. comm. to author from Greg Houseal, July 27, 2015; unreferenced).

Numerous measures have been implemented to control invasive plants at Clay Prairie. A diversion berm was constructed (1984 to 1985) just outside of the southern boundary fence of the preserve to control incoming silt and weed seeds from adjacent agricultural lands (pers. comm. to author from Paul Whitson, [date unknown] 1999; unreferenced). Cutting, weed-whipping and herbicide treatment have been implemented to control invasive woody plants including gray dogwood (*Cornus foemina* P. Miller ssp. *racemosa* [Lam.] J.S. Wilson), silver poplar (*Populus alba* L.), smooth sumac (*Rhus glabra* L.), Canada elderberry (*Sambucus canadensis* L.), and willow (*Salix* sp.). Herbicide treatment, hand pulling, cutting of seed heads, and mowing have been variously applied to control invasive grasses such as smooth brome (*Bromus inermis* Leysser), reed canary grass (*Phalaris arundinacea* L.) and forbs including giant ragweed (*Ambrosia trifida* L.), mustard (*Brassica* sp.), and wild parsnip (*Pastinaca sativa* L.) (Camarata 1990, 1991).

Several instances of native plant introductions are documented for Clay Prairie and potentially contribute to the current flora.

Fig. 2. Location of 27 Iowa state preserves with significant (> 30% of total area) prairie vegetation. See Table 1 for information about each preserve. Clay Prairie State Preserve indicated with a star.

Unspecified prairie plants were transplanted into the preserve in 1989, with little evidence of subsequent survival. In 1990, porcupine grass (*Hesperostipa spartea* [Trin.] Barkworth) seeds collected just north of the prairie and stratified in the UNI greenhouse were sown into the south side of Clay Prairie. During this same year, additional plant species raised from seed in the UNI greenhouse (seed source unspecified) were plugged into bare spots in the south end of the preserve, including side-oats grama (*Bouteloua curtipendula* Torrey), purple prairie clover (*Dalea purpurea* Vent.), porcupine grass, round-headed bush clover (*Lespedeza capitata* Michx.), and Indian grass (*Sorghastrum nutans* [L.] Nash). In June 1991, Canada wild rye (*Elymus canadensis* L.) and oat (*Avena* sp.) were seeded south of the south border of Clay Prairie to provide an “erosion filter”; the seed source of these plants is likewise unspecified (Camarata 1991).

PREVIOUS BOTANICAL STUDY OF CLAY PRAIRIE STATE PRESERVE

Cyrus W. Lantz provided the first established published list of vascular plants for Clay Prairie (Lantz 1969). The checklist includes 98 vascular plant taxa representing 33 families. The most diverse families were Poaceae (11 taxa), Fabaceae (11 taxa) and Asteraceae (20 taxa). Fourteen (14.3 %) of these taxa were non-native plant taxa. Lantz (1969) did not indicate that he collected voucher specimens during his study, and we have not encountered specimens collected at Clay Prairie by Lantz in any Iowa herbarium.

The occurrence of several other vascular plant taxa at Clay Prairie prior to this study and not reported by Lantz (1969) are documented by voucher specimens deposited in the Martin L. Grant Herbarium (ISTC) at UNI and the Ada Hayden Herbarium (ISC) at Iowa State University (ISU). Management reports for Clay Prairie (Camarata 1990, 1991, 1994) mention several additional plant taxa not reported by Lantz (1969) as present in the prairie in the late 1980s and early 1990s.

METHODS

We inventoried the vascular plants of Clay Prairie State Preserve during the growing seasons between 1997 and 2001 and again between 2008 and 2013, for a total of 29 total visits. We collected voucher specimens of all but eight vascular plant taxa

encountered during our field work, and deposited these in the Ada Hayden Herbarium (ISC). Freese also searched the Martin L. Grant Herbarium (ISTC) at UNI for specimens collected in the preserve. From this work, we compiled a checklist of vascular plants known from the preserve. We assigned abundance codes to plants we personally observed at Clay Prairie as follows:

Common: widely distributed and often found growing in large quantities in several different habitats.

Frequent: widespread but not abundant and usually found in only one type of habitat.

Infrequent: not widespread and often not found in places where it might be expected to occur.

Sparse: found in only one or a few places

Family concepts follow Judd et al. 2008. Nomenclature for plant taxa follows Flora of North America Committee (1993+; vols. 2-5, 7-8, and 19-26) and Eilers and Roosa (1994), with the former taking priority. The origin (native, non-native) status of each taxon is likewise based on the above references. Common names follow Eilers and Roosa (1994) and Swink and Wilhelm (1994), with the former taking priority.

We compared the richness and floristic quality of the prairie flora documented from Clay Prairie State Preserve with that documented from 26 other prairies protected in the Iowa State Preserve system (Fig. 2). These represent a broad variety of tallgrass and mid-grass prairie types in Iowa, and range in area from 3.8 ha to 319.7 ha (Table 1) as summarized by Rosburg (2001). All of these prairies comprise a substantial ($\geq 38\%$) portion of the total land area within the state preserve in which they occur (Table 1). We compiled the floras documented from these other 26 prairies from all (48) available sources (e.g., published floras, plant lists, and plot data presented in M.S. theses, Ph.D. dissertations, technical reports to sponsoring conservation agencies, and unpublished checklists) into a spreadsheet. Then, we compared the number of native “Iowa prairie taxa” documented from each of these 27 prairies (including Clay Prairie) protected in the Iowa state preserve system. We determined that a given native vascular plant taxon was an “Iowa prairie plant” if 1) it was stated to occur in prairie habitat in the Iowa checklist of vascular plants (Eilers and Roosa 1994), or 2) if it was included in a list of Iowa prairie plants compiled by Pearson (1991).

We also compared the number of native Iowa prairie plant taxa of high conservation concern in Clay Prairie with the other 26 prairies. To do this, we utilized Iowa Coefficient of Conservation (ICC) values assigned by a panel of experts to each native vascular plant taxon in the Iowa flora (Drobney et al. 2001). As broadly applied, coefficients of conservation range from 0 to 10, and reflect the likelihood that a given plant taxon will occur in natural habitat (Swink and Wilhelm 1994, Bourdaghs et al. 2006). Following Taft et al. (1997), who state that Illinois plant taxa assigned coefficient of conservation values of 7 to 10 associate mostly with (or are restricted to) natural areas, we tallied the number of Iowa prairie taxa with ICC values equal to 7, 8, 9 or 10 reported for each prairie to allow comparison. Although it was tempting to calculate and compare floristic quality indices and mean ICC values among all 27 prairies (e.g., Swink and Wilhelm 1994, Higgins et al. 2001), we decided against this because of potential effects of area on those metrics (Matthews 2003, Matthews et al. 2005) and especially because of the great range in area of prairie vegetation (1.3 ha to 121.4 ha) represented among the 27 prairies included in this comparison.

Finally, we compared the number of Iowa prairie plant taxa listed as endangered, threatened, or of special concern (Iowa

Table 1. Descriptions of 27 prairies protected in the Iowa State Preserve system. The number of each prairie corresponds to its location presented in Fig. 1. Prairie areas, primary prairie communities and secondary prairie communities are from Rosburg (2001), except for Cedar Hills Sand Prairie and Clay Prairie (pers. comm., Thomas Rosburg, Department of Biology, Drake University). LH = Loess Hills, NE = Northeast, NW = Northwest, TG = Tall Grass. '*' indicates a study for which voucher specimens were collected to document the majority of the flora reported.

Number	Name	Iowa County	Prairie Area (ha)	Primary Prairie Community	Secondary Prairie Community	Flora(s)
3	Anderson Prairie	Emmet	80.9	NW Wet-mesic TG	NW Dry-mesic TG	Moats 1988, 1989, *Bice and James 2013
19	Richard W. Pohl Memorial Preserve at Ames High Prairie	Story	4.0	Mesic TG	Dry-mesic TG	*Freckmann 1966, Norris 1995, *Thompson 2013
1	Cayler Prairie	Dickinson	60.7	NW Wet-mesic TG	NW Dry-mesic TG	*Aikman and Thorne 1956, Smith et al. 1991
14	Cedar Hills Sand Prairie	Black Hawk	10.2	Dry-mesic sand TG		*Crum 1972, Glenn-Lewin 1981b, *Freese 2012
13	Clay Prairie	Butler	1.2	Wet-mesic TG	Mesic TG	Lantz 1969
4	Crossman Prairie	Howard	4.0	NE mesic TG		Glenn-Lewin 1981a, Bockenstedt 2004
23	Dinesen Prairie	Shelby	8.1	Mesic TG		Rosburg 2005a
18	Doolittle Prairie	Story	10.1	Wet-mesic TG		*Woodley 1983, *Wetzel et al. 1999
8	Five Ridge Prairie	Plymouth	121.4	LH Mid-grass	LH Dry-mesic TG	Duxbury 1982
2	Freda Haffner Kettlehole	Dickinson	30.4	NW Mid-grass	NW Dry-mesic TG	Gerhardt 1940, *Freese and Platt 1991
5	Hayden Prairie	Howard	97.1	NE Mesic TG	NE Wet-mesic TG	Moyer 1953, Christiansen 1993
7	Hoffman Prairie	Cerro Gordo	14.2	Wet-mesic TG		Christiansen 1999a
12	Kalsow Prairie	Pocahontas	48.6	Mesic TG	Wet-mesic TG	Moyer 1953, Richards 1970 based on Brotherson 1969, Dornbush 2004
25	Kish-Ke-Kosh Prairie	Jasper	6.1	Dry-mesic TG		Kurtz 1976, Wagner 1980, *Kessler 2002
15	Liska-Stanek Prairie	Webster	8.1	Mesic TG	Wet-mesic TG	Wilson 1979
22	Manikowski Prairie	Clinton	16.2	Dry-mesic lime TG		Cady 2002
20	Marietta Sand Prairie	Marshall	6.1	Dry-mesic sand TG	Mesic TG	Leoschke 1982, Christiansen 1999b
9	Mount Talbot	Plymouth, Woodbury	36.4	LH Mid-grass	LH Dry-mesic TG	Pearson and Fleckenstein 1989, *Hazlett 2003
11	Nestor Stiles	Cherokee	4.0	Mesic TG		Schennum and Dudley 1983, Pearson and Leoschke 1987
21	Rock Island	Linn	4.0	Dry-mesic sand TG		Cady 2003
27	Rolling Thunder	Warren	24.3	Mesic TG		Mabry 2002
24	Sheeder Prairie	Guthrie	10.1	Mesic TG	Dry-mesic TG	Kennedy 1969, Roosa 1989, Rosburg 1996
10	Steele Prairie	Cherokee	80.9	Mesic TG	Wet-mesic TG	Pearson and Fleckenstein 1987
6	Stinson Prairie	Kossuth	12.1	NW Wet-mesic TG	NW Dry-mesic TG	Glenn-Lewin 1976, Crist 1978
16	Sylvan Runkel	Monona	121.4	LH Mid-grass	LH Dry-mesic TG	Pearson 1996, Rosburg 2005b
17	Turin Loess Hills	Monona	80.9	LH Mid-grass	LH Bluff Mid-grass	Novacek-Bates 1989
26	Williams Prairie	Johnson	8.1	Wet TG	Wet-mesic TG	*Sorenson 1962

Fig. 3. Relationship between total number of native Iowa prairie plant taxa versus total prairie area for 27 prairies protected in the Iowa state preserve system. Clay Prairie State Preserve is identified by an arrow. Each of the 27 prairies is characterized in Table 1.

Administrative Code 2002) in Clay Prairie with those documented from the other 26 preserves.

RESULTS

A total of 214 vascular plant taxa have been identified at Clay Prairie State Preserve, of which 174 (81%) are native to Iowa (Appendix A). Of the 214 taxa on this list, 162 are eudicots, 50 are monocots, and 2 are fern allies. This list includes representatives of 50 families and 145 genera. A total of 174 taxa (81 %) in the Clay Prairie flora are native plant taxa. The most diverse plant families represented in Clay Prairie are Asteraceae (45 taxa), Poaceae (30), Cyperaceae (14), Fabaceae (13), and Rosaceae (11). The most diverse plant genera are *Carex* (11 taxa), *Symphyotrichum* (6), *Solidago* (5), *Asclepias* (4), *Euphorbia* (4), and *Viola* (4). One of these taxa, smooth clustered sedge (*Carex aggregata* Mack.), is considered to be of special concern in Iowa (Iowa Administrative Code 2002).

We were unable to locate fourteen plant taxa from Lantz's previously compiled checklist (1969) on the preserve. These include seven native taxa (pasque flower [*Anemone patens* L. var. *multifida* Fritzell], purple coneflower [*Echinacea angustifolia* DC.], rockrose [*Helianthemum bicknellii* Fern.], June grass [*Coeleria macrantha* {Ledeb.} Schultes], bunch-flower [*Melanthium virginicum* L.], sand dropseed [*Sporobolus cryptandrus* {Torrey} A. Gray], heart-leaved meadow parsnip [*Zizia aptera* {Gray} Fern.]) and seven non-native taxa (*Carduus* sp., cypress spurge [*Euphorbia cyparissias* L.], alfalfa [*Medicago sativa* L.], red-seeded dandelion [*Taraxacum laevigatum* {Willdenow} de Candolle var. *erythrospermum* {Andrzejowski ex Besser} J. Weiss], cultivated red raspberry [*Rubus idaeus* L.], white clover [*Trifolium repens* L.], rabbit-foot clover [*T. arvense* L.]).

In contrast, we encountered 117 plant taxa (including 91 native taxa) during the current study that were not reported by Lantz (1969). Among these are three taxa planted in the preserve in the early 1990s (Camarata 1990, 1991): side-oats grama, Canada wild rye, and Indian grass. The seed sources of these taxa are not specified in these reports. Although not reported as occurring at Clay Prairie by Lantz (1969), Indian grass was apparently present in the preserve prior to its introduction reported by Camarata because it is listed as an associate of tall nut-rush (*Scleria triglomerata* Michx.) on the specimen label (ISC 365127) of the latter taxon collected in the preserve in the mid-1980s. Finally, our observation of hairy

Fig. 4. Relationship between total number of native Iowa prairie plant taxa assigned high (7, 8, 9 or 10) values of the Iowa Coefficient of Conservation (ICC) versus total prairie area for 27 prairies protected in the Iowa state preserve system. Clay Prairie State Preserve is identified by an arrow. Each of the 27 prairies is characterized in Table 1.

blue violet (*Viola sororia* Willd.) at Clay Prairie during the current study is probably due to its mid-1980s introduction in soil brought in to create the diversion berm just outside the southern edge of the preserve (pers. comm., Paul Whitson, fall 1999; unreferenced). Whitson noted subsequent hybridization of hairy blue violet with other violet taxa in the preserve.

Additional vascular plant taxa reported anecdotally by preserve workday managers (Smith 1991, Camarata 1990, 1991, 1994) were not reported by Lantz (1969) nor relocated by us at Clay Prairie. These include six native taxa (northern reed grass [*Calamagrostis stricta* {Timm} Koeler], downy painted cup [*Castilleja sessiliflora* Pursh], purple coneflower [*Echinacea purpurea* {L.} Moench], Maximilian's sunflower [*Helianthus maximilliani* Schrader], blazing star [*Liatris aspera* Michx.], Solomon's seal [*Polygonatum biflorum* {Walter} Ell.]) and five non-native taxa (velvet leaf [*Abutilon theophrasti* Medicus], tickseed coreopsis [*Coreopsis lanceolata* L.], canary grass [*Phalaris canariensis* L.], silver poplar, hedgerow rose [*Rosa rugosa* Thunb.]). Given the anecdotal nature of these reports, we did not include these taxa in the checklist (Appendix A).

The total number of Iowa prairie plant taxa ranged from 69 to 182 among the 27 Iowa state preserves considered in this study. Of these, Clay Prairie ranked 10th with 135 prairie plant taxa (Fig. 3). The total number of Iowa prairie plant taxa with ICC values = 7, 8, 9, or 10 ranged from 15 to 83 among these same 27 Iowa state preserve prairies. In this comparison, Clay Prairie ranked 10th among all of these prairies with 55 Iowa prairie plant taxa assigned high (7 to 10) ICC values (Fig. 4). No Iowa prairie plant in the Clay Prairie flora is currently considered to be endangered, threatened, or of special concern in the State of Iowa (note: Iowa special concern taxon smooth clustered sedge, mentioned above as occurring at Clay Prairie, is not an Iowa prairie plant). In this regard, the Clay Prairie flora is tied for last place along with four other prairies included in this comparison.

DISCUSSION

To truly comprehend how small Clay Prairie State Preserve is, consider that the next largest prairies featured in the Iowa state preserves system (Ames High Prairie, Crossman Prairie, Nestor Stiles Prairie, Rock Island) are more than three times larger (Table 1). The Iowa state preserve containing the largest amount of prairie, Five Ridge Prairie, contains more than 250 times as

much prairie as Clay Prairie. More than 200 prairies the size of Clay Prairie would fit inside one square mile (the common unit into which much of the Iowa landscape is subdivided by intersecting county roads).

Thus, it is remarkable that the Clay Prairie flora ranks so highly in floristic comparisons with other prairies in the Iowa state preserve system (Figs. 3, 4). One likely explanation for this result is that the Clay Prairie flora has been studied more comprehensively and for a longer period of time than many of the other prairies in the Iowa state preserve system. We strongly suspect that the floras of many of the other 26 prairies, if thoroughly inventoried, would be discovered to contain more plant taxa with prairie affinity and/or more prairie plant taxa with high ICC values (i.e., ICC = 7, 8, 9, or 10) than are currently documented from Clay Prairie.

Role of Clay Prairie State Preserve as Vegetation Remnant in Conservation

What is the conservation value of Clay Prairie State Preserve, a mere speck on the Iowa landscape? Saunders et al. (1991) described four reasons that natural vegetation remnants should be preserved. These include: a) *species oriented reasons*, in which individual species and their gene pools are preserved; b) *example oriented reasons*, in which remnants preserve particular ecosystems, communities, habitats and vegetation assemblages; c) *process oriented reasons*, by which remnants preserve important ecological and evolutionary processes; and d) *social oriented reasons*, in which remnants provide opportunities for people to experience nature in a variety of ways. We summarize how Clay Prairie fulfills the above four criteria below.

Clay Prairie possesses a diverse flora of prairie plants (Fig. 3) including many taxa of high conservation concern (Fig. 4). Populations of some of these plants have provided seed for prairie restoration projects elsewhere (Camarata 1991). What of other biota? Studies of insect diversity and biology conducted in prairie remnants in Iowa and adjacent states have focused on ants (Foster et al. 2010), bees (Slagle and Hendrix 2009), beetles (St. Pierre and Hendrix 2003, Larson and Work 2003, Larsen et al. 2003), butterflies (Debinski and Kelly 1998, Selser and Schramm 1992), grasshoppers (Bomar 2001, Bomar and Secrist 2002), leafhoppers (Hamilton 1995), moths (Summerville 2008, Summerville et al. 2006), and thrips (Gerdes 1982). Bryophytes (van der Linden and Farrar 1981) and fungi (Liberta and Anderson 1986, Tiffany and Knaphus 1995) have also been the focus of biotic inventories conducted on prairie remnants in the Midwest. To our knowledge, Clay Prairie State Preserve has not been the focus of any inventory of insects, fungi, or bryophytes. Scientists and conservation managers of Clay Prairie would benefit from such information.

In preserving an example of tallgrass prairie in Iowa, Clay Prairie satisfies criterion b. Shafer (1995) argues that although conservation strategies should focus on large vegetation conservation reserves, these large reserves should be augmented with smaller ones because the latter may be the only available representative of a given vegetation type. This justification for preservation of small preserves certainly applies in Iowa, where the majority (> 99.9 %) of historic tallgrass prairie vegetation no longer exists (Smith 1998).

Certainly ecological and evolutionary processes (criterion c) continue at Clay Prairie, though highly modified through the lens of human impacts on this prairie and surrounding landscape. Dozens of studies have appeared in the scientific literature that focus on various biological processes occurring in prairie remnants. Clay Prairie was included in at least one of these

studies (Wilsey et al. 2005) that focused on prediction of plant extinction rates in prairie remnants based on species-area curves. *The results of this study indicated that many more native prairie plant species were present in small prairie remnants than were predicted based on island biogeography theory* (MacArthur and Wilson 1967).

Finally, Clay Prairie State Preserve fulfills criterion d, justifying preservation of vegetation remnants in providing opportunities for people to experience prairie. Smith (1991), Camarata (1990, 1991, 1994, 1996, 1997, 1998) and Houseal (pers. comm. to author, July 27, 2015; unreferenced) describe numerous field trips and prairie management work days that took place at Clay Prairie in the early 1970s and through the late 2000s.

RECOMMENDATIONS

The high floristic quality of Clay Prairie State Preserve is clear, and its future preservation is ensured due to its status as an Iowa state preserve. To date, comprehensive floristic studies are published in the scientific literature for only eight prairies in the Iowa state preserves system: Ames High School Prairie (Freckmann 1966), Cayler Prairie (Aikman and Thorne 1956), Cedar Hills Sand Prairie (Crum 1972), Clay Prairie (Lantz 1969), Doolittle Prairie (Wetzel et al. 1998), Freda Haffner Kettlehole (Freese and Platt 1991), Stinson Prairie (Glenn-Lewin 1976), and Williams Prairie (Sorenson 1962). Floristic data for the other 19 prairie preserves (Table 1) are available as technical reports to several conservation agencies, university masters theses and PhD dissertations, and in a few cases as private plant lists compiled by citizen scientists with expertise in plant identification.

It took us more than two years to locate and obtain all of the floras and reports cited in Table 1. These vary widely in duration (multiple years versus several days), intensity (plants of high conservation priority versus entire flora), and extent (entire preserve versus plot data collected along a transect line). Obviously, the conservation management agencies who oversee prairies protected in the Iowa state preserve system should sponsor comprehensive floristic inventories of prairie remnants that they manage which span several years and focus on the entire flora. These agencies should then encourage publication of these floristic studies in peer-reviewed journals. Such careful characterization of the floras of prairie remnants in the Iowa state preserve system would provide data for sophisticated ecological studies of plant community and population dynamics (e.g., Wilsey et al. 2005). Likewise, we recommend that establishment and survey of permanent plots surveyed at multiple points in time (e.g., Rosburg 1996, Dornbush 2004) be undertaken to inform preserve managers of vegetation change in these prairie remnants, especially increasing or decreasing trends of exotic plant populations.

Only 12 of the 49 studies cited in Table 1 indicate that vouchers were collected to thoroughly document the featured floras. Thus, it is impossible to verify the occurrence of plant taxa reported for the majority of prairie remnants in the Iowa State Preserve system. To illustrate this problem, Lantz (1969) reported one species of purple coneflower (*Echinacea angustifolia*) from Clay Prairie in the 1960s, Camarata (1990) anecdotally noted the occurrence of another species of purple coneflower (*Echinacea purpurea*) in the preserve during the late 1980s and early 1990s, and Lammers vouchered pale coneflower (*Echinacea pallida* Nutt.) from the preserve in 1980 (subsequently encountered by us during the current study). Future floristic studies should adhere to rigorous standards, *especially comprehensive documentation of the flora via voucher specimens and/or digital photographs with subsequent deposit in university herbaria*, as outlined by Palmer and Richardson (2012).

Recent ecological studies have explored how prairie vegetation influences the insects that reside within it. For example, Panzer and Schwartz (1998) showed that native plant species richness explained a significant proportion of variation in conservative insect species richness in 50 prairie remnants near Chicago, IL. Other studies (e.g., Panzer 2002) have investigated how prairie management (i.e., burning) influences insect communities. Published studies of insect diversity in Iowa state prairie preserves (e.g., Larson and Work 2003, Larson et al. 2003, Slagle and Hendrix 2009) are limited. We recommend that, concurrent with increased effort to characterize the floras of prairie remnants in the Iowa state preserve system, an increased focus on insect inventory in these prairie remnants be undertaken by academicians, conservation agency biologists, and citizen scientists with expertise in insect identification.

Finally, the numerous management reports by Smith (1991) and Camarata (1990, 1991, 1994, 1996, 1997, 1998) provide extremely useful historical information about Clay Prairie, including a record of seed collection and reintroduction of native plants in the preserve, accounts of visits by school groups to the prairie, and a vegetation map locating regions of problematic woody plant occurrence. We encourage a resumption of annual reporting of activities at Clay Prairie to inform future generations of prairie enthusiasts about this prairie jewel.

ACKNOWLEDGEMENTS

Lawrence J. Eilers, Martin L. Grant, Thomas G. Lammers, and Mark Leoschke each collected important voucher specimens of plant taxa at Clay Prairie independent of this and the prior floristic study of this preserve. Ron Camarata, Greg Houseal, Daryl Smith, and Paul Whitson documented management history of Clay Prairie through written reports and personal communications to us. Cyrus W. Lantz introduced Freese to Clay Prairie on a field trip he led there during the summer of 1966. John Pearson provided us with numerous Iowa prairie floras and prepared several figures for use in this paper. Tom Rosburg compiled information about Iowa prairies that we utilized in this paper. Michael Palmer and three anonymous reviewers provided many helpful suggestions that have helped us to improve the quality of this paper. Mary E. Brown, Billie Hemmer, Mark Leoschke, Brian Hazlett, Steve Hendrix, Greg Houseal, Daryl Howell, Kelly Kindscher, Deb Lewis, Jean Prior, Tom Rosburg, Daryl Smith, Arnold Webster, Paul Whitson, Mark Widrechner, Gerould Wilhelm, Tre Wilson, and Scott Zager all assisted us with various other aspects of this project. We thank all of the above for their contributions to this study.

The floristic comparisons presented in this paper were possible because of the 49 botanical studies conducted in Iowa prairies by the authors and the following 43 individuals: John Aikman, Philip Bice, Paul Bockenstedt, Jack Brotherson, Tom Cady, Allen Crist, Paul Christiansen, Glenn Crum, Kathryn Davis, Matt Dornbush, R. Dudley, Alexis Duxbury, John Fleckenstein, Robert Freckmann, Charles Gerhardt, Brian Hazlett, Aubrie James, Carol Kern, Carl Kurtz, Cyrus Lantz, Mark Leoschke, David Glenn-Lewin, Brian Hazlett, Robert Kennedy, Jim Kessler, Cathy Mabry, Bob Moats, John Moyer, Jean Novacek-Bates, John Pearson, William Platt, Mary Richards, Dean Roosa, Tom Rosburg, Wayne Schennum, Daryl Smith, Paul Sorenson, Jimmie Thompson, Robert Thorne, Mark Wagner, Paul Wetzel, Tre Wilson, and Greg Woodley. We thank all of the above for their efforts to document historic prairie vegetation in Iowa.

This research was funded in part by a research grant from the Iowa State Preserves Board, Iowa Department of Natural Resources for the 1999-2000 field season.

LITERATURE CITED

- AIKMAN, J. M. and R. F. THORNE. 1956. The Cayler Prairie: an ecological and taxonomic study of a northwest Iowa prairie. *Proceedings of the Iowa Academy of Science* 63:177-200.
- ANDERSON, W. I. 1998. Iowa's Geological Past – three billion years of change. University of Iowa Press, Iowa City.
- BETZ, R. F. 1989. Species composition of old settler and sand prairie cemeteries in northern Illinois and northwestern Indiana. p. 79-87. *In*: D. D. Smith, and C. A. Jacobs (eds.). *Proceedings of the 12th North American Prairie Conference: recapturing a vanishing heritage*. University of Northern Iowa, Cedar Falls, IA.
- BETZ, R. F. and H. F. LAMP. 1989. Species composition of old settler silt-loam prairies. p. 33-39. *In*: T. B. Bragg, and J. Stubbendieck (eds.). *Prairie pioneers: history and culture: Proceedings of the Eleventh North American Prairie Conference*. Lincoln, NE.
- BICE, P. M. and A. R. M. JAMES. 2013. The vascular flora of Anderson Prairie State Preserve, Emmet County. Unpublished report, State Preserve Advisory Board, Iowa Department of Natural Resources, Des Moines, IA.
- BOCKENSTEDT, P. J. 2004. Botanical survey of Crossman Prairie State Preserve, Howard County, Iowa. Unpublished report, State Preserve Advisory Board, Iowa Department of Natural Resources, Des Moines, IA.
- BOMAR, C. R. 2001. Comparison of grasshopper (Orthoptera: Acrididae) communities on remnant and reconstructed prairies in western Wisconsin. *Journal of Orthoptera Research* 10:105-112.
- BOMAR, C. R. and E. SECRIST. 2002. Grasshoppers (Orthoptera: Acrididae) associated with prairie remnants in the lower Chippewa River State Natural Area (LCRSNA) of western Wisconsin. *Journal of Orthoptera Research* 11:37-41.
- BOURDAGHS, M., C. A. JOHNSTON, and R. R. REGAL. 2006. Properties and performance of the floristic quality index in Great Lakes coastal wetlands. *Wetlands* 26:718-735.
- BROTHERSON, J. D. 1969. Species composition, distribution, and phytosociology of Kalsow Prairie, a mesic tall-grass prairie in Iowa. Ph.D. Dissertation, Iowa State University, Ames, IA.
- BUCKNER, R. L. 1982. Soil survey of Butler County, Iowa. United States Department of Agriculture.
- CADY, T. F. 2002. Floristic inventory of Manikowski Prairie State Preserve. Unpublished report, State Preserves Advisory Board, Iowa Department of Natural Resources, Des Moines, IA.
- CADY, T. F. 2003. Floristic inventory of Rock Island State Preserve. Unpublished report, State Preserves Advisory Board, Iowa Department of Natural Resources, Des Moines, IA.
- CAMARATA, R. D. 1990. University of Northern Iowa biological preserve and greenhouse report 1989-1990. 113 p.
- CAMARATA, R. D. 1991. University of Northern Iowa biological preserve and greenhouse report 1990-1991. 98 p.
- CAMARATA, R. D. 1994. University of Northern Iowa biological preserve and greenhouse report 1993-1994. 180 p.
- CAMARATA, R. D. 1996. University of Northern Iowa biological preserve and greenhouse annual report 1995-1996. 180 p.
- CAMARATA, R. D. 1997. University of Northern Iowa biological preserve and greenhouse annual report 1996-1997. 107 p.
- CAMARATA, R. D. 1998. University of Northern Iowa biological preserve and greenhouse report 1997-1998. 93 p.
- CHRIST, A. M. 1978. A gradient analysis of a northern Iowa tall-grass prairie. M.S. Thesis, Iowa State University, Ames, IA.
- CHRISTIENSEN, P. 1993. Report on the flora and vegetation of Hayden Prairie, Howard County, Iowa. Unpublished report, State Preserves Advisory Board, Iowa Department of Natural Resources, Des Moines, IA.
- CHRISTIENSEN, P. 1999a. Report on the flora and vegetation of Hoffman Prairie, Cerro Gordo County, Iowa. Unpublished report, State Preserve Advisory Board, Iowa Department of Natural Resources, IA.

- CHRISTIANSEN, P. 1999b. Report on the flora and vegetation of Marietta Sand Prairie, Marshall County, Iowa. Unpublished report, State Preserves Advisory Board, Iowa Department of Natural Resources, Des Moines, IA.
- CLAY PRAIRIE PRESERVE, BUTLER CENTER. Pamphlet published by Biological Preserves Committee, Department of Biology, University of Northern Iowa, Cedar Falls [date unknown; accessed 2013 Aug 18; www.uni.edu/biology/clay].
- CRUM, G. H. 1972. Flora of a sand prairie in Black Hawk County, Iowa. *Proceedings of the Iowa Academy of Science* 78:81–87.
- DEBINSKI, D. D. and L. KELLY. 1998. Decline of Iowa populations of the regal fritillary (*Speyeria idalia*) Drury. *Journal of the Iowa Academy of Science* 105:16–22.
- DORNBUSH, M. E. 2004. Plant community change following fifty-years of management at Kalsow Prairie Preserve, Iowa, U.S.A. *American Midland Naturalist* 151:241–250.
- DROBNEY, P. D., G. S. WILHELM, D. G. HORTON, M. J. LEOSCHKE, D. Q. LEWIS, J. A. PEARSON, D. M. ROOSA, and D. D. SMITH. 2001. Floristic quality assessment for the state of Iowa. Unpublished report. Accessed 2013 Aug 18; <http://www.public.iastate.edu/~herbarium/coeffici.html>
- DUXBURY, A. 1982. The Loess Hills Nature Preserve. A report to the Nature Conservancy, Minneapolis, MN and Des Moines, IA.
- EILERS, L. J. 1966. The remnant prairie flora in northeast Iowa. *Proceedings of the Iowa Academy of Science* 73:33–43.
- EILERS, L. J. and D. M. ROOSA. 1994. The Vascular Plants of Iowa: an annotated checklist and natural history. University of Iowa Press, Iowa City.
- FLORA OF NORTH AMERICA EDITORIAL COMMITTEE (eds). 1993+. *Flora of North America north of Mexico*. Oxford University Press, New York (NY).
- FOSTER, J., J. TRAGER, and E. L. FREESE. 2010. A small prairie gem in Iowa. p. 22–34. *In*: B. Bursari (ed.). *Proceedings of the 21st North American Prairie Conference: the prairie meets the river*. Winona State University, Winona, MN.
- FRECKMANN, R. W. 1966. The prairie remnants of the Ames area. *Proceedings of the Iowa Academy of Science* 73:126–136.
- FREESE, E. L. 2012. Vascular plant species list, Cedar Hills Sand Prairie Preserve, Black Hawk County, Iowa. Unpublished report, State Preserve Advisory Board, Iowa Department of Natural Resources, Des Moines, IA.
- FREESE, E. L. and W. J. PLATT. 1991. Vascular flora of Arend's Kettle, Freda Haffner Kettlehole State Preserve. *Journal of the Iowa Academy of Science* 98:102–107.
- GERHARDT, C. E. 1940. An ecological survey of a large kettle hole. M.S. Thesis. University of Iowa, Iowa City.
- GERDES, C. F. 1982. Distribution of adult overwintering Thysanoptera in an Illinois prairie. *Journal of the Kansas Entomological Society* 55:418–422.
- GLENN-LEWIN, D. C. 1976. The vegetation of Stinson Prairie, Kossuth County, Iowa. *Proceedings of the Iowa Academy of Science* 83:88–93.
- GLENN-LEWIN, D. C. 1981a. Inventory and management plan for Crossman Prairie. Report Number 12 to The Nature Conservancy, Des Moines, IA.
- GLENN-LEWIN, D. C. 1981b. Inventory and management plan for Mark Sand Prairie Preserve. Report Number 9 to The Nature Conservancy, Des Moines, IA.
- HAMILTON, K. G. A. 1995. Evaluation of leafhoppers and their relatives (Insecta: Homoptera: *Auchenorrhyncha*) as indicators of prairie preserve quality. p. 211–226. *In*: D. C. Harnett (ed.). *Proceedings of the 14th Annual North American Prairie Conference: Prairie biodiversity*. Kansas State University, Manhattan, KS.
- HAZLETT, B. T. 2003. Floristic inventory of Stone State Park, Plymouth and Woodbury Counties, IA. Unpublished report, State Preserve Advisory Board, Iowa Department of Natural Resources, Des Moines, IA.
- HERZBERG, R. and J. A. PEARSON. 2001. The guide to Iowa State preserves. University of Iowa Press, Iowa City.
- HIGGINS, J. J., G. E. LARSON, and K. F. HIGGINS. 2001. Floristic comparisons of tallgrass prairie remnants managed by different land stewardships in eastern South Dakota. p. 21–25. *In*: N. P. Bernstein, and L. J. Ostrander (eds.). *Proceedings of the Seventeenth North American Prairie Conference: seeds for the future ~ roots of the past*. North Iowa Area Community College, Mason City, IA.
- IOWA ADMINISTRATIVE CODE. 2002. Endangered and threatened plant and animal species. Natural Resources Commission, Chapter 77. Iowa Administrative Code, January 9, 2002, Des Moines, IA.
- JUDD, W. S., C. J. CAMPBELL, E. A. KELLOGG, P. F. STEVENS, and M. J. DONOGHUE. 2008. *Plant Systematics*, 3rd edition. Sinauer Associates, Inc., Sunderland, MA.
- KENNEDY, R. K. 1969. An analysis of tallgrass prairie vegetation relative to slope position, Sheeder Prairie, Iowa. M.S. Thesis, Iowa State University, Ames, IA.
- KERR, K. and J. WHITE. 1981. A volunteer-supported effort to find and preserve prairie and savanna remnants in Illinois cemeteries. p. 181–183. *In*: R. L. Stuckey, and K. J. Reese (eds.). *The prairie peninsula—in the “shadow” of Transeau*: *Proceedings of the Sixth North American Prairie Conference*. Ohio State University, Columbus, OH.
- KESSLER, J. E. 2002. The vascular flora and a vegetational analysis of Kish-Ke-Kosh State Preserve: 1981–1983 and 2001. Unpublished floristic inventory.
- KURTZ, C. 1976. Klepinger Sand Prairie. Unpublished floristic inventory.
- LANTZ, C. W. 1969. The Clay Prairie in Butler County, Iowa. *Proceedings of the Iowa Academy of Science* 76:109–112.
- LARSON, K. J. and T. T. WORK. 2003. Differences in ground beetle (Coleoptera: Carabidae) of original and reconstructed tallgrass prairies in northeastern Iowa, USA, and impact of 3-year spring burn cycles. *Journal of Insect Conservation* 7:153–166.
- LARSON, K. J., T. T. WORK, and F. F. PURRINGTON. 2003. Habitat use patterns by ground beetles (Coleoptera: Carabidae) of northeastern Iowa. *Pedobiologia* 47:288–299.
- LEOSCHKE, M. J. 1982. Floristic inventory of Marietta Sand Prairie, Marshall Co., IA. Unpublished report, State Preserves Advisory Board, Iowa Department of Natural Resources, Des Moines, IA.
- LIBERTA, A. E. and R. C. ANDERSON. 1986. Comparison of vesicular-arbuscular mycorrhiza species composition, spore abundance and inoculums potential in an Illinois prairie and adjacent agricultural sites. *Bulletin of the Torrey Botanical Club* 113:178–182.
- MABRY, C. 2002. Floristic inventory of Rolling Thunder Prairie State Preserve. Unpublished report, State Preserves Advisory Board, Iowa Department of Natural Resources, Des Moines, IA.
- MACARTHUR, R. H. and E. O. WILSON. 1967. *The theory of island biogeography*. Princeton University Press, Princeton, NJ.
- MATTHEWS, J. W. 2003. Assessment of the floristic quality index for use in Illinois, USA wetlands. *Natural Areas Journal* 23:53–60.
- MATTHEWS, J. W., P. A. TESSENE, S. M. WIESBROOK, and B. W. ZERCHER. 2005. Effect of area and isolation on species richness and indices of floristic quality in Illinois, USA wetlands. *Wetlands* 25:607–615.
- MENGES-SCHABEN, C. L. 1998. Tallgrass prairie remnants of western Iowa cemeteries. M.S. Thesis, University of Nebraska at Omaha, Omaha, NE.
- MOATS, R. 1988. Floristic inventory of Anderson Prairie State Preserve, Emmet Co., IA. Unpublished report, State Preserves Advisory Board, Iowa Department of Natural Resources, Des Moines, IA.
- MOATS, R. 1989. Management plan for Anderson Prairie, Emmet County, Iowa. Unpublished report, State Preserves Advisory Board, Iowa Department of Natural Resources, Des Moines, IA.
- MOYER, J. F. 1953. Biology of native prairie in Iowa. Ph.D. Dissertation, Iowa State University, Ames, IA.
- NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION (NOAA). 1998. Annual Survey, Iowa. 109:1–13. Ashville, NC.
- NOVACEK-BATES, J. 1989. Management plan and inventory of Turin Loess Hills Preserve, Monona Co., IA. Unpublished report, State Preserves Advisory Board, Iowa Department of Natural Resources, Des Moines, IA.

- NORRIS, W. R. 1995. Ames High School Prairie in *A Natural Areas Inventory of Ames, Iowa*, pp. 327–332. M.S. Thesis, Iowa State University, Ames, IA.
- PALMER, M. W. and J. C. RICHARDSON. 2012. Biodiversity data in the information age: do 21st century floras make the grade? *Castanea* 77:46–59.
- PANZER, R. 2002. Prescribed burning and insect conservation. *Conservation Biology* 16:1296–1307.
- PANZER, R. and M. W. SCHWARTZ. 1998. Effectiveness of a vegetation-based approach to insect conservation. *Conservation Biology* 12:693–702.
- PEARSON, J. A. 1991. Iowa prairie plant list. Unpublished data, Natural Areas Inventory program, Iowa Department of Natural Resources. *In*: W. R. Norris. 1995. A natural area inventory of Ames, Iowa. M.S. Thesis, Iowa State University, Ames, IA.
- PEARSON, J. A. 1996. Management plan for Sylvan Runkel State Preserve, Loess Hills Wildlife Area, Monona Co., IA. Unpublished report, State Preserves Advisory Board, Iowa Department of Natural Resources, Des Moines, IA.
- PEARSON, J. A. and J. W. FLECKINSTEIN. 1987. Management plan for Steele Prairie State Preserve. Unpublished report, State Preserves Advisory Board, Iowa Department of Natural Resources, Des Moines, IA.
- PEARSON, J. A. and J. W. FLECKINSTEIN. 1989. Management plan for Mt. Talbot State Preserve, Plymouth and Woodbury Counties, IA. Unpublished report. State Preserves Advisory Board, Iowa Department of Natural Resources, Des Moines, IA.
- PEARSON, J. A. and M. J. LEOSCHKE. 1987. Floristic inventory of Nestor-Stiles State Preserve, Cherokee Co., IA. Unpublished report, State Preserves Advisory Board, Iowa Department of Natural Resources, Des Moines, IA.
- PHILLIPS, L. J., P. B. MARCUM, D. T. BUSEMEYER, and J. E. EBINGER. 2010. Vascular flora of Short Pioneer Cemetery Prairie Nature Preserve, Grundy County, Illinois: composition and change since 1977. *Erigenia* Number 24:5–11.
- PRIOR, J. C. 1991. Landforms of Iowa. University of Iowa Press, Iowa City.
- RICHARDS, M. 1970. Report on the vegetation of Kalsow Prairie. Unpublished report, State Preserves Advisory Board, Iowa Department of Natural Resources, Des Moines, IA.
- ROOSA, D. M. 1978. Prairie preservation in Iowa – history, present status, and future plans. p. 204–206. *In*: D. Glenn-Lewin, and R. Q. Landers, Jr. (eds.). Fifth Midwest Prairie Conference Proceedings. Ames, IA.
- ROOSA, D. M. 1981. Directory of state preserves. Iowa State Preserves Board (State Preserves Advisory Board), Iowa Conservation Commission (Iowa Department of Natural Resources), Des Moines, IA.
- ROOSA, D. M. 1989. Resource inventory and management plan for Sheeder Prairie State Preserve, Guthrie County, Iowa. A report to the Iowa State Preserves Advisory Board, Parks, Preserves and Recreation Division, Iowa Department of Natural Resources, Des Moines, IA.
- ROSBURG, T. R. 1996. A quantitative plant inventory of Sheeder Prairie and a comparison with its composition in 1968. Unpublished report, State Preserves Advisory Board, Iowa Department of Natural Resources, Des Moines, IA.
- ROSBURG, T. R. 2001. Iowa's prairie heritage: from the past, through the present, and into the future. p. 1–14. *In*: N. P. Bernstein, and L. J. Ostrander (eds.). Proceedings of the Seventeenth North American Prairie Conference: seeds for the future ~ roots of the past. North Iowa Area Community College, Mason City, IA.
- ROSBURG, T. R. 2005a. Floristic and plant community inventories on Dinesen Prairie State Preserve, Shelby County, Iowa. Unpublished report, State Preserve Advisory Board, Iowa Department of Natural Resources, Des Moines, IA.
- ROSBURG, T. R. 2005b. Floristic and plant community inventories on Sylvan Runkel State Preserve, Monona County, IA. Unpublished report, State Preserves Advisory Board, Iowa Department of Natural Resources, Des Moines, IA.
- RUCH, D. G., B. G. TORKE, K. S. BADGER, and P. E. ROTHROCK. 2010. The vascular flora in three prairie cemeteries in Henry County, Indiana. *Proceedings of the Indiana Academy of Science* 119:35–51.
- ST. PIERRE, M. J. and S. D. HENDRIX. 2004. Patch use by a monophagous herbivore in fragmented prairie landscapes. *Prairie Naturalist* 36:231–242.
- SAMSON, F. B. and F. L. KNOPF. 1994. Prairie conservation in North America. *Bioscience* 44:418–21.
- SAUNDERS, D. A., G. W. ARNOLD, A. A. BURBIDGE, and A. J. M. HOPKINS. 1991. The role of remnants of native vegetation in nature conservation: future directions. p. 387–392. *In*: D. A. Saunders, G. W. Arnold, A. A. Burbidge, and A. M. Hopkins (eds.). *Nature conservation: the role of remnants of native vegetation*. Surrey Beatty and Sons Pty Limited, Chipping Norton, New South Wales, Australia.
- SCHENNUM, W. E. 1986. A comprehensive survey for prairie remnants in Iowa: methods and preliminary results. p. 163–167. *In*: Clambey, G. K., and R. H. Pemble (eds.). *The prairie: past, present, and future: proceedings of the Ninth North American Prairie Conference*. Tri-College University, Moorhead, MN.
- SCHENNUM, W. E. and R. DUDLEY. 1984. Floristic inventory of Nestor-Stiles State Preserve, Cherokee Co., IA. Unpublished floristic inventory.
- SELSER, E. J. and P. SCHRAMM. 1992. Comparative species diversity and distribution of butterflies in remnant and restored tallgrass prairie sites. p. 63–65. *In*: D. D. Smith, and C. A. Jacobs (eds.). *Proceedings of the Twelfth North American Prairie Conference: recapturing a prairie heritage*. University of Northern Iowa, Cedar Falls, IA.
- SHAFFER, C. L. 1995. Value and shortcomings of small reserves. *Bioscience* 45:80–88.
- SLAGLE, M. W. and S. D. HENDRIX. 2009. Reproduction of *Amorpha canescens* (Fabaceae) and diversity of its bee community in a fragmented landscape. *Oecologia* 161:813–823.
- SMITH, D. D. 1991. Annual on-site inspection report of Clay Prairie State Preserve. Unpublished report, State Preserves Advisory Board, Iowa Department of Natural Resources, Des Moines, IA.
- SMITH, D. D. 1998. Iowa's prairie: original extent and loss, preservation and recovery efforts. *Journal of the Iowa Academy of Science* 105:94–108.
- SMITH, D. D., C. KERN, and K. DAVIS. 1991. Establishment of long term monitoring of *Lespedeza leptostachya* in Iowa. Unpublished report, State Preserves Advisory Board, Iowa Department of Natural Resources, Des Moines, IA.
- SORENSEN, P. D. 1962. The Williams Prairie: A prairie relict in Johnson County. *Proceedings of the Iowa Academy of Science* 69:45–53.
- SUMMERVILLE, K. S. 2008. Species diversity and persistence in restored and remnant tallgrass prairies of North America: a function of species' life history, habitat type, or sampling bias? *Journal of Animal Ecology* 77:487–494.
- SUMMERVILLE, K. S., C. J. CONOAN, and R. M. STEICHEN. 2006. Species traits as predictors of lepidopteran composition in restored and remnant tallgrass prairies. *Ecological Applications* 16:891–900.
- SWINK, F. and G. S. WILHELM. 1994. *Plants of the Chicago region*. Indiana University Press, Bloomington.
- TAFT, J. B., G. S. WILHELM, D. M. LADD, and L. A. MASTERS. 1997. Floristic quality assessment for vegetation in Illinois: a method for assessing vegetation integrity. *Erigenia* 15:3–95.
- THOMPSON, J. D. 2013. Richard W. Pohl Memorial State Preserve at Ames High School, Story County, Iowa. Vascular plant taxa list. Unpublished floristic inventory.
- TIFFANY, L. H. and G. KNAPHUS. 1995. Fungus pathogens of prairie plants in Iowa. p. 49–54. *In*: D. C. Harnett (ed.). *Proceedings of the 14th Annual North American Prairie Conference: Prairie biodiversity*. Kansas State University, Manhattan, KS.
- THORNWAITE, C. W. 1948. An approach towards a natural classification of climate. *Geographical Review* 38:55–94.
- VAN DER LINDEN, J. and D. R. FARRAR. 1983. An ecological study of the bryophytes of a natural prairie in northwestern Iowa. *The Bryologist* 86:1–13.
- WAGNER, M. D. 1980. Kish-Ke-Kosh survey of vascular plants. Unpublished floristic inventory.

- WEB SOIL SURVEY. United States Department of Agriculture, Natural Resources Conservation Service. Accessed 2015 July 14 from: <http://websoilsurvey.sc.egov.usda.gov/App/WebSoilSurvey.aspx>
- WETZEL, P. R., W. R. NORRIS, and K. M. LYLES. 1999. The vascular flora of Doolittle Prairie State Preserve - a prairie pothole wetland complex. *Journal of the Iowa Academy of Science* 106:26–33.
- WHITE, J. 1978. Illinois natural areas inventory technical reports. Volume 1. Survey methods and results. Illinois Natural Areas Inventory, Urbana, IL.
- WHITE, J. 1988. Protection of Pine Ridge Cemetery Prairie: A story of persistence and cooperation. *Natural Areas Journal* 8:100–106.
- WILSEY, B. J., L. M. MARTIN, and H. W. POLLEY. 2005. Predicting plant extinction based on species-area curves in prairie fragments with high beta richness. *Conservation Biology* 19:1835–1841.
- WILSON, T. 1979. Unpublished floristic inventory of Liska-Stanek Prairie State Preserve.
- WOODLEY, G. R. 1983. The vascular flora of Doolittle Prairie. Herbarium Report Collection 1-37, Ada Hayden Herbarium, Botany Department, Iowa State University, Ames, IA.

APPENDIX

Checklist of Clay Prairie State Preserve flora. Voucher specimens collected by E.L. Freese, M.J. Leoschke and W.R. Norris are deposited in the Ada Hayden Herbarium (ISC) at Iowa State University, and those collected by L.J. Eilers, M.L. Grant and T.G. Lammers are deposited in the Martin L. Grant Herbarium (ISTC) Herbarium at the University of Northern Iowa.

Key

No symbol = Taxa native to Iowa
 * = Taxa not native to Iowa
 Boldface = Native Iowa Prairie Plant (defined in text)
 Nomenclatural synonyms, alternate family names as presented in Eilers and Roosa (1993)
 F = Taxa reported by Freese and Norris
 L = Taxa reported by Lantz
 Abundance codes defined in text
 Collection number of voucher specimens in italics

FERNS AND FERN ALLIES (formerly Pteridophytes)

EQUISETACEAE

Equisetum arvense L. (common horsetail) [L, F] Frequent *Freese 12 May 2009-7*
Equisetum laevigatum A. Br. (smooth scouring-rush) [L, F] Infrequent *Freese 11 Aug 2009-6*

ANGIOSPERMS: MONOCOTYLEDONS

ASPARAGACEAE [formerly LILIACEAE, in part]

**Asparagus officinale* L. (garden asparagus) [F] Sparse *Freese 11 Aug 2009-41*

CONVALLARIACEAE [formerly LILIACEAE, in part]

Polygonatum biflorum (Walter) Ell. (Solomon's seal) [F] Sparse *Freese 11 Aug 2009-7*

CYPERACEAE

Carex aggregata Mack. (smooth clustered sedge) [F] Infrequent *Freese 6 June 2000-1*
Carex bicknellii Britton (copper-shouldered oval sedge) [L, F] Infrequent *Freese 2 June 2001-3*
Carex brevior (Dewey) Mack. ex Lunell (plains oval sedge) [F] Infrequent *Freese 2 June 2001-1*
Carex buxbaumii Wahl. (dark-scaled sedge) [L, F] Infrequent *Freese 21 June 1997-1*
Carex gravida Bailey (long-awned bracted sedge) [F] Infrequent *Freese 2 June 2001-2*

Carex haydenii Dewey (long-scaled tussock sedge) [F] Infrequent *Freese 21 June 1997-3*
Carex meadii Dewey (Mead's stiff sedge) [F] Infrequent *Freese 2 June 2001-4*
Carex normalis Mack. (spreading oval sedge) [F] Infrequent *Freese 2 June 2001-5*
Carex pellita Willd. [*Carex lanuginosa* Michx.] (broad-leaved wooly sedge) [F] Infrequent *Freese 17 July 1998-2*
Carex pennsylvanica Lam. (common oak sedge) [F] Infrequent *Freese 12 May 2009-15*
Carex sartwellii Dewey (running marsh sedge) [F] Infrequent *Freese 17 July 1998-1*
Eleocharis compressa Sulliv. (spike-rush) [F] Infrequent *Freese 21 June 1997-2*
Scirpus atrovirens Willd. (dark green bulrush) [F] Sparse *Freese 17 July 1998-3*
Scleria triglomerata Michx. (tall nut-rush) [F] Sparse *Leoschke 1 July 1983 - ISC 365127*

HYPOXIDACEAE [formerly AMARYLLIDACEAE, in part]

Hypoxis hirsuta (L.) Cov. (yellow stargrass) [L, F] Frequent *Lammers 3345*

IRIDACEAE

Iris shrevei Small (blue flag) [F] Sparse Norris 2013-6-2-8.5
Sisyrinchium campestre Bickn. (prairie blue-eyed grass) [L, F] Frequent *Freese 12 May 2009-3*

JUNCACEAE

Juncus dudleyi Weig. (Dudley's rush) [F] Infrequent *Freese 6 July 2009-4*

LILIACEAE

Lilium philadelphicum L. [*Lilium philadelphicum* L. var. *andinum* (Nutt.) Ker-Gawl] (wood lily) [L, F] Sparse *Freese 6 July 2009-8*

MELANTHIACEAE [formerly, LILIACEAE, in part]

Melanthium virginicum L. (bunch-flower) [L]

POACEAE

**Agrostis gigantea* Roth (redtop) [L, F] Infrequent *Freese 6 July 2009-1*
Andropogon gerardii Vitman (big bluestem) [L, F] Frequent *Eilers 6550*
Bouteloua curtipendula (Michx.) Torrey (side-oats grama) [F] Sparse *Freese 6 July 2009-23*
Bromus ciliatus L. (Canada brome grass) [L, F] Sparse
 **Bromus inermis* Leysser (smooth brome) [F] Sparse *Freese 6 July 2009-24*
Calamagrostis canadensis (Michx.) Beauv. (bluejoint) [F] Infrequent *Freese 20 Sept 2008-43*

- **Dactylis glomerata* L. (orchard grass) [F] Sparse Norris 2011-8-11-36
Dichanthelium leibergii (Vasey) Freckm. (Leiberg's panic grass) [F] Infrequent Freese 8 July 2000-1
Dichanthelium ovale (Elliott) ssp. *praecocious* (Hitch. & Chase) Freckmann & Lelong [*Dichanthelium acuminatum* (Sw.) Gould & Clark var. *villosum* (Gray) Gould & Clark] (long-haired panic grass) [F] Infrequent Freese 6 July 2009-17
Dichanthelium oligosanthes (Schultes) Gould var. *scribnerianum* (Nash) Gould (Scribner's panic grass) [F] Infrequent Freese 8 Aug 2000-1
Elymus canadensis L. (Canada wild rye) [F] Infrequent Freese 20 Sept 2008-30
**Elymus repens* (L.) Gould (*Agropyron repens* (L.) Beauv.) (quack grass) [F] Sparse Freese 11 Aug 2009-32
Elymus virginicus L. (Virginia wild rye) [F] Sparse Freese 8 Aug 2000-2
Hesperostipa spartea (Trin.) Barkworth [*Stipa spartea* Trin.] (porcupine grass) [L, F] Infrequent Freese 6 July 2009-21
Hordeum jubatum L. (squirrel-tail barley) [L, F] Sparse Freese 6 July 2009-25
Koeleria macrantha (Ledeb.) Schultes (June grass) [L]
Muhlenbergia mexicana (L.) Trin. (leafy satin grass) [F] Infrequent Freese 20 Sept 2008-44
Muhlenbergia racemosa (Michx.) BSP. (marsh muhly) [F] Infrequent Freese 20 Sept 2008-45
Panicum dichotomiflorum Michx. (knee grass) [F] Sparse
Phalaris arundinacea L. (reed canary grass) [F] Sparse Freese 6 July 2009-22
**Pbleum pratense* L. (timothy) [L, F] Sparse
**Poa pratensis* L. (Kentucky bluegrass) [L, F] Frequent Freese 8 July 2000-2
Schizachyrium scoparium (Michx.) Nash (little bluestem) [L, F] Infrequent Freese 20 Sept 2008-46
**Setaria faberi* Herrm. (giant foxtail) [F] Sparse Freese Sept 2008-18
**Setaria pumila* (Poiret) Roemer & Schultes ssp. *pumila* [*Setaria glauca* (L.) Beauv.] (yellow foxtail) [F] Sparse Norris 2011-8-15-2
**Setaria viridis* (L.) Beauv. (green foxtail) [F] Sparse Norris 2011-8-15-5
Sorghastrum nutans (L.) Nash (Indian grass) [F] Infrequent Loeschke 1 July 1983 - ISC 365127
Spartina pectinata Link (slough grass) [L, F] Infrequent Freese 8 Aug 2009-3
Sporobolus cryptandrus (Torrey) A. Gray (sand dropseed) [L]
Sporobolus heterolepis (Gray) Gray (prairie dropseed) [F] Infrequent Freese 20 Sept 2008-16

ANGIOSPERMS: EUDICOTYLEDONS (formerly, Dicotyledons)

- ADOXACEAE [includes, CAPRIFOLIACEAE, in part]
Sambucus canadensis L. (Canada elderberry) [F] Sparse Freese 8 Sept 2010-2
AMARANTHACEAE [includes CHENOPODIACEAE]
**Amaranthus retroflexus* L. (pigweed) [F] Sparse Freese 16 Sept 2000-2
Amaranthus tuberculatus (Moquin-Tandon) J.D. Sauer [*Amaranthus rudis* Sauer] (western water hemp) [F] Sparse Freese 16 Sept 2000-4
**Chenopodium album* L. (lamb's quarters) [F] Sparse Freese 16 Sept 2000-3

ANACARDIACEAE

- Rhus glabra* L. (smooth sumac) [L, F] Infrequent Freese 20 Sept 2008-32

APIACEAE

- Cicuta maculata* L. (water hemlock) [L, F] Infrequent Freese 6 July 2009-18
**Daucus carota* L. (Queen Anne's lace) [F] Sparse Freese 20 Sept 2008-31
Eryngium yuccifolium Michx. (rattlesnake master) [L, F] Infrequent Freese 20 Sept 2008-33
**Pastinaca sativa* L. (wild parsnip) [L, F] Sparse Freese 11 Aug 2009-1
Sanicula canadensis L. (black snakeroot) [F] Sparse Freese 11 Aug 2009-2
Sium suave Walter (water parsnip) [L, F] Infrequent Freese 11 Aug 2009-5
Zizia aptera (Gray) Fern. (heart-leafed golden alexander) [L]
Zizia aurea (L.) Koch (golden alexander) [L, F] Common Freese 20 Sept 2008-41

APOCYNACEAE [includes ASCLEPIACEAE]

- Apocynum sibiricum* Jacq. (dogbane) [F] Infrequent Freese 11 Aug 2009-3
Asclepias birtella (Pennell) Woodson (tall green milkweed) [F] Sparse
Asclepias syriaca L. (common milkweed) [L, F] Infrequent Freese 11 Aug 2009-4
Asclepias tuberosa L. ssp. *interior* Woodson (butterfly weed) [F] Sparse Freese 11 Aug 2009-10
Asclepias viridiflora Raf. (green milkweed) [L, F] Sparse

ASTERACEAE

- Achillea millefolium* L. [*Achillea millefolium* L. ssp. *lanulosa* (Nutt.) Piper] (western yarrow) [L, F] Sparse Freese 6 July 2009-14
Ambrosia artemisiifolia L. (common ragweed) [F] Sparse Freese 11 Aug 2009-9
Ambrosia trifida L. (giant ragweed) [L, F] Sparse Freese 16 Sept 2000-1
Antennaria neglecta Greene (pussytoes) [F] Sparse Freese 6 June 1998-3
Antennaria plantaginifolia (L.) Hooker (ladies'-tobacco) [L, F] Sparse Freese 6 June 1998-8
Arnoglossum plantagineum Raf. (*Cacalia plantaginea* (Raf.) Shinnery) (prairie Indian plantain) [L, F] Infrequent Freese 6 July 2009-7
Artemisia ludoviciana Nutt. (white sage) [F] Sparse Freese 11 Aug 2009-11
**Carduus* sp. (musk thistle) [L]
**Cirsium arvense* (L.) Scop. (Canada thistle) [F] Sparse Freese 11 Aug 2009-34
Cirsium flodmanii (Rydb.) Arthur (Flodman's thistle) [L, F] Infrequent Freese 11 Aug 2009-33
**Cirsium vulgare* (Savi) Tenore (bull thistle) [F] Sparse Freese 11 Aug 2009-35
Coreopsis palmata Nutt. (prairie coreopsis) [L, F] Frequent Freese 20 Sept 2008-28
Doellingeria umbellata (Miller) Nees var. *umbellata* (P. Miller) Nees (*Aster umbellatus* Miller) (flat-topped aster) [F] Infrequent Freese 8 Sept 2010-1
Echinacea angustifolia DC. (purple coneflower) [L]
Echinacea pallida Nutt. (pale coneflower) [F] Sparse Lammers 3726

- Erigeron annuus* (L.) Pers. (annual fleabane) [F] Sparse Freese 6 July 2009-20
- Erigeron strigosus* Muhl. ex Willd. (daisy fleabane) [L, F] Infrequent Freese 6 July 2009-11
- Euthamia graminifolia* (L.) Nutt. (common grass-leaved goldenrod) [F] Infrequent Freese 20 Sept 2008-34
- Helianthus grosseserratus* Martens (saw-tooth sunflower) [F] Frequent Freese 11 Aug 2009-12
- Helianthus pauciflorus* Nuttall [*Helianthus rigidus* (Cass.) Desf.] (prairie sunflower) [F] Frequent Freese 11 Aug 2009-13
- Helianthus tuberosus* L. (Jerusalem artichoke) [F] Infrequent Freese 20 Sept 2008-12
- Heliopsis helianthoides* (L.) Sweet (ox-eye) [L, F] Frequent Freese 20 Sept 2008-13
- Krigia biflora* (Walter) Blake (false dandelion) [L, F] Infrequent Freese 6 July 2009-6
- Lactuca canadensis* L. (wild lettuce) [F] Sparse Freese 11 Aug 2009-14
- Liatris pycnostachya* Michx. (prairie blazing star) [L, F] Infrequent Freese 11 Aug 2009-20
- Packera aurea* (L.) A. Love & D. Love [*Senecio aureus* L.] (golden ragwort) [L, F] Sparse Norris 2013-6-2-1
- Parthenium integrifolium* L. (wild quinine) [L, F] Infrequent Freese 20 Sept 2008-20
- Prenanthes racemosa* Michx. (glaucous white lettuce) [F] Infrequent Freese 20 Sept 2008-29
- Ratibida pinnata* (Vent.) Barnh. (gray-headed coneflower) [L, F] Frequent Freese 20 Sept 2008-23
- Rudbeckia hirta* L. (black-eyed Susan) [L, F] Sparse Freese 6 July 2009-12
- Silphium laciniatum* L. (compass plant) [L, F] Frequent Freese 20 Sept 2008-11
- Silphium perfoliatum* L. (cup plant) [F] Sparse Norris 2013-6-2-13
- Solidago canadensis* L. (tall goldenrod) [L, F] Frequent Freese 20 Sept 2008-47
- Solidago gigantea* Aiton (smooth goldenrod) [F] Frequent Norris 2013-6-1-2
- Solidago missouriensis* Nutt. (Missouri goldenrod) [F] Infrequent Freese 11 Aug 2009-15
- Solidago nemoralis* Aiton (field goldenrod) [F] Infrequent Freese 11 Aug 2009-17
- Solidago rigida* L. (stiff goldenrod) [F] Infrequent Freese 20 Sept 2008-7
- Solidago speciosa* Nutt. (showy goldenrod) [F] Infrequent Freese 20 Sept 2008-8
- Symphotrichum ericoides* (L.) G. L. Nesom [*Aster ericoides* L.] (heath aster) [F] Sparse Freese 20 Sept 2008-25
- Symphotrichum leave* (L.) A. Love & D. Love [*Aster laevis* L.] (smooth blue aster) [F] Frequent Lammers 4007
- Symphotrichum lanceolatum* (Willd.) G. L. Nesom [*Aster lanceolatus* Willd.] (panicled aster) [F] Frequent Freese 20 Sept 2008-26
- Symphotrichum novae-angliae* (L.) G. L. Nesom [*Aster novae-angliae* L.] (New England aster) [F] Infrequent Freese 20 Sept 2008-42
- Symphotrichum oolentangiense* (Riddell) G. L. Nesom [*Aster azureus* Lindley] (sky-blue aster) [F] Infrequent Grant 14966
- Symphotrichum pilosum* (Willd.) G. L. Nesom [*Aster pilosus* Willd.] (hairy aster) [F] Sparse Freese 20 Sept 2008-40
- **Taraxacum laevigatum* (Willdenow) de Candolle var. *erythrospermum* (Andrzejowski ex Besser) J. Weiss [*Taraxacum laevigatum* (Willd.) DC.] (red-seeded dandelion) [L]
- **Taraxacum officinale* Weber (common dandelion) [F] Sparse Freese 12 May 2009-12
- **Tragopogon dubius* Scop. (goat's-beard) [L, F] Sparse Freese 11 Aug 2009-16
- BORAGINACEAE**
- Hackelia virginiana* (L.) I.M. Johnston (stickseed) [F] Sparse Freese 11 Aug 2009-22
- Lithospermum canescens* (Michx.) Lehm. (hoary puccoon) [L, F] Infrequent Freese 12 May 2009-1
- BRASSICACEAE**
- **Barbarea vulgaris* R. Br. (yellow rocket) [F] Sparse Freese 6 June 2000-19
- **Brassica rapa* L. [*Brassica campestris* L.] (field mustard) [L, F] Sparse Freese 6 June 2000-20
- **Capsella bursa-pastoris* (L.) Medicus (shepherd's purse) [F] Sparse Freese 8 Sept 2010-3
- Cardamine bulbosa* (Schreber) BSP. (spring cress) [L, F] Infrequent Freese 6 June 2000-4
- CAMPANULACEAE**
- Campanula aparinoides* Pursh (marsh bellflower) [F] Infrequent Freese 8 July 2000-1
- Lobelia spicata* Lam. (spiked lobelia) [L, F] Infrequent Freese 6 July 2009-26
- Triodanis perfoliata* (L.) Nieuw. (Venus's looking-glass) [F] Sparse Freese 6 June 1998-10
- CARYOPHYLLACEAE**
- **Cerastium fontanum* Baumgarten [*Cerastium vulgatum* L.] (mouse-eared chickweed) [F] Sparse Freese 6 June 1998-7
- **Silene latifolia* Poiret [*Silene pratensis* (Rafn.) Gren. & Godron] (white campion) [F] Sparse Freese 6 July 2009-27
- CISTACEAE**
- Helianthemum bicknellii* Fern. (hoary frostweed) [L]
- CONVOLVULACEAE**
- Calystegia sepium* (L.) R.Br. (bindweed) [F] Sparse Freese 17 June 1998-6
- **Convolvulus arvensis* L. (European bindweed) [L, F] Sparse Freese 11 Aug 2009-21
- CORNACEAE**
- Cornus foemina* P. Miller ssp. *racemosa* (Lam.) J.S. Wilson (gray dogwood) [F] Common Freese 11 August 2009-18
- EUPHORBIACEAE**
- Euphorbia corollata* L. (flowering spurge) [L, F] Infrequent Freese 11 August 2009-17
- **Euphorbia cyparissias* L. (cypress spurge) [L]
- Euphorbia maculata* L. (carpet spurge) [F] Sparse Freese 16 Sept 2000-3
- Euphorbia nutans* Lag. (nodding spurge) [F] Sparse Freese 16 Sept 2000-4
- FABACEAE**
- Amorpha canescens* Pursh (leadplant) [L, F] Frequent Freese 20 Sept 2008-19
- Baptisia bracteata* Muhl. ex Ell. var. *glabrescens* (Larisey) Isley (cream wild indigo) [L, F] Infrequent Freese 20 Sept 2008-15

Baptisia lactea (Raf.) Thieret (white wild indigo) [L, F] Infrequent Freese 20 Sept 2008-14

Dalea candida Willd. (white prairie clover) [L, F] Sparse Lammers 3733

Dalea purpurea Vent. (purple prairie clover) [L, F] Infrequent Freese 6 July 2009-28

Desmodium canadense (L.) DC. (showy tick-trefoil) [L, F] Infrequent Freese 11 Aug 2009-23

Lathyrus palustris L. (marsh vetchling) [L, F] Infrequent Freese 6 July 2009-31

Lespedeza capitata Michx. (round-headed bush clover) [L, F] Infrequent Freese 20 Sept 2008-48

**Medicago sativa* L. (alfalfa) [L]

**Melilotus alba* Medicus (white sweet clover) [F] Sparse Freese 6 July 2009-32

**Trifolium arvense* L. (rabbit foot clover) [L]

**Trifolium pratense* L. (red clover) [F] Sparse Freese 11 Aug 2009-39

**Trifolium repens* L. (white clover) [L]

FAGACEAE

Quercus rubra L. [*Quercus borealis* Michx. f. var. *maxima* (Marsh.) Ashe] (northern red oak) [L, F] Sparse

GENTIANACEAE

Gentiana andrewsii Griseb. (bottle gentian) [F] Infrequent Freese 20 Sept 2008-10

Gentiana x billingtonii Farw. (soapwort gentian) [F] Sparse Freese 20 Sept 2008-50

Gentiana puberulenta J. Pringle (downy gentian) [L, F] Sparse Freese 20 Sept 2008-51

HYDROPHYLLACEAE

Ellisia nyctelea L. (wild tomato) [F] Sparse Freese 6 July 2009-29

LAMIACEAE

**Glechoma hederacea* L. (creeping Charlie) [F] Sparse Freese 6 June 1998-1

Monarda fistulosa L. (horsemint) [F] Frequent Freese 6 Aug 1997-2

**Nepeta cataria* L. (catnip) [F] Sparse Freese 16 Sept 2000-5

Pycnanthemum virginianum (L.) Dur. & Jackson (common mountain mint) [L, F] Infrequent Freese 20 Sept 2008-35

Stachys palustris L. (woundwort) [F] Infrequent Freese 11 Aug 2009-30

MALVACEAE

**Malva* sp. (mallow) [F] Sparse Freese 16 Sept 2000-6

MORACEAE

**Morus alba* L. (white mulberry) [F] Sparse Freese 11 Aug 2009-27

NYCTAGINACEAE

Mirabilis nyctaginea (Michx.) MacM. (wild four-o'clock) [F] Sparse

ONAGRACEAE

Oenothera biennis L. ssp. *centralis* Munz (evening primrose) [F] Sparse Freese 11 Aug 2009-24

OROBANCHACEAE [formerly SCROPHULARIACEAE, in part]

Castilleja coccinea (L.) Sprengel (Indian paintbrush) [L, F] Infrequent Lammers 2560

Pedicularis canadensis L. (lousewort) [L, F] Infrequent Freese 12 May 2009-4

OXALIDACEAE

Oxalis stricta L. (yellow wood sorrel) [F] Sparse Freese 17 July 2013-2

Oxalis violacea L. (violet wood sorrel) [L, F] Frequent Freese 12 May 2009-2

PLANTAGINACEAE [formerly SCROPHULARIACEAE, in part]

Veronicastrum virginicum (L.) Farwell (Culver's root) [L, F] Frequent Freese 6 July 2009-3

POLEMONIACEAE

Phlox pilosa L. (prairie phlox) [L, F] Infrequent Freese 6 July 2009-33

POLYGONACEAE

Persicaria pensylvanica (L.) M. Gomez [*Polygonum pensylvanicum* L. var. *laevigatum* Fern.] (pinkweed) [F] Sparse Norris 2011-8-11-21

**Rumex crispus* L. (curly dock) [L, F] Sparse Freese 11 Aug 2009-37

Rumex acetosella L. (red sorrel) [F] Sparse Norris 2013-6-25-4

**Rumex altissimus* Wood (pale dock) [F] Sparse Norris 2013-6-1-9

PRIMULACEAE

Dodecatheon meadia L. (shooting star) [L, F] Common Lammers 3338

Lysimachia ciliata L. (fringed loosestrife) [L, F] Infrequent Freese 20 Sept 2008-21

Lysimachia quadriflora Sims (whorled loosestrife) [F] Infrequent Freese 6 July 2009-5

RANUNCULACEAE

Anemone canadensis L. (Canada anemone) [L, F] Frequent Freese 20 Sept 2008-35

Anemone cylindrica Gray (thimbleweed) [F] Sparse Freese 20 Sept 2008-36

Anemone patens L. var. *multifida* Pritzel [*Pulsatilla patens* (L.) P. Mill.] (pasque flower) [L]

Thalictrum dasycarpum Fischer & Ave-Lall. (purple meadow-rue) [L, F] Infrequent Freese 20 Sept 2008-37

RHAMNACEAE

Ceanothus americanus L. var. *pitcheri* T. & G. (New Jersey tea) [L, F] Frequent Freese 20 Sept 2008-27

ROSACEAE

Fragaria virginiana Duchesne (wild strawberry) [L, F] Frequent Freese 12 May 2009-11

Geum triflorum Pursh (prairie smoke) [L, F] Frequent Lammers 3561

Potentilla arguta Pursh (prairie cinquefoil) [F] Infrequent Lammers 3730

**Potentilla recta* L. (sulphur cinquefoil) [F] Sparse Freese 6 June 1998-4

Potentilla simplex Michx. (common cinquefoil) [F] Sparse Norris 2-13-6-2-10

Prunus americana Marsh. (wild plum) [F] Sparse Norris 2013-6-1-6

Prunus serotina Ehrh. (wild black cherry) [F] Sparse Norris 2011-8-11-31

Prunus virginiana L. (choke cherry) [L, F] Sparse Freese 11 Aug 2009-25

Rosa arkansana Porter var. *suffulta* (Greene) Cockerell (sunshine rose) [L, F] Infrequent Freese 20 Sept 2008-31

Rosa blanda Aiton (meadow rose) [F] Infrequent Freese 20 Sept 2008-39

**Rubus idaeus* L. (cultivated red raspberry) [L]

Rubus occidentalis L. (black raspberry) [F] Infrequent *Freese* 17 July 1998-4
Spiraea alba Du Roi (meadowsweet) [L, F] Infrequent *Freese* 11 Aug 2009-36

RUBIACEAE

Galium aparine L. (cleavers) [F] Sparse *Norris* 2013-6-2-7
Galium boreale L. (northern bedstraw) [L, F] Frequent *Lammers* 3565
Galium obtusum Bigelow (wild madder) [F] Sparse *Freese* 17 June 1998-2

SALICACEAE

Populus deltoides Bartram ex Marsh. (cottonwood) [F] Frequent *Freese* 20 Sept 2008-5
Populus tremuloides Michx. (quaking aspen) [L, F] Frequent *Freese* 20 Sept 2008-4
Salix discolor Muhl. (pussy willow) [L, F] Infrequent *Lammers* 3724
Salix humilis Marsh. (prairie willow) [L, F] Infrequent *Lammers* 3336
Salix petiolaris Smith (meadow willow) [F] Infrequent *Freese* 20 Sept 2008-3

SAPINDACEAE [formerly ACERACEAE]

Acer saccharinum L. (silver maple) [F] Sparse

SANTALACEAE

Comandra umbellata (L.) Nutt. (bastard toadflax) [L, F] Infrequent *Freese* 12 May 2009-6

SAXIFRAGACEAE

Heuchera richardsonii R. Br. (alumroot) [L, F] Infrequent *Freese* 6 July 2009-34
Micranthes pensylvanica (L.) Haworth [*Saxifraga pensylvanica* L.] (swamp saxifrage) [L, F] Infrequent *Lammers* 3333

SCROPHULARIACEAE

Scrophularia lanceolata Pursh (early figwort) [F] Sparse *Freese* 17 June 1998-5

SOLANACEAE

Physalis heterophylla Nees (ground cherry) [F] Sparse *Freese* 11 Aug 2009-25
Physalis virginiana P. Miller (ground cherry) Sparse [F] *Norris* 2011-8-11-15
Solanum americanum P. Miller (black nightshade) [F] Sparse *Freese* 16 Sept 2000-7

ULMACEAE

**Ulmus pumila* L. (Siberian elm) [F] Sparse *Freese* 11 Aug 2009-30
Ulmus rubra Muhl. (red elm) [F] Sparse *Norris* 2011-8-11-34

URTICACEAE

Parietaria pensylvanica Muhl. ex Willd. (pellitory) [F] Sparse *Norris* 2011-8-11-23
Urtica dioica L. (stinging nettle) [F] Sparse *Norris* 2013-6-1-4

VERBENACEAE

Verbena urticifolia L. (white vervain) [F] Sparse *Norris* 2011-8-11-16

VIOLACEAE

Viola nephrophylla Greene (northern bog violet) [F] Frequent *Freese* 11 Aug 2009-31
Viola pedata L. (bird's-foot violet) [L, F] Frequent *Lammers* 3341
Viola pedatifida G. Don. (prairie violet) [L, F] Frequent *Lammers* 3342
Viola sororia Willd. (hairy blue violet) [F] Sparse *Freese* 6 June 1998-9

VITACEAE

Parthenocissus sp. [F] Sparse *Norris* 2013-6-1-1
Vitis riparia Michx. (riverbank grape) [F] Sparse *Norris* 2011-8-11-27