

1955

## B-Vitamins Stimulatory to Cellulose Digestion by Washed Suspensions of Rumen Microorganisms

Glen Hall  
*Iowa State College*

Edmund W. Cheng  
*Iowa State College*

Wise Burroughs  
*Iowa State College*

*Let us know how access to this document benefits you*

Copyright ©1955 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

---

### Recommended Citation

Hall, Glen; Cheng, Edmund W.; and Burroughs, Wise (1955) "B-Vitamins Stimulatory to Cellulose Digestion by Washed Suspensions of Rumen Microorganisms," *Proceedings of the Iowa Academy of Science*, 62(1), 273-278.

Available at: <https://scholarworks.uni.edu/pias/vol62/iss1/32>

This Research is brought to you for free and open access by the IAS Journals & Newsletters at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact [scholarworks@uni.edu](mailto:scholarworks@uni.edu).

**Offensive Materials Statement:** Materials located in UNI ScholarWorks come from a broad range of sources and time periods. Some of these materials may contain offensive stereotypes, ideas, visuals, or language.

## **B-Vitamins Stimulatory to Cellulose Digestion by Washed Suspensions of Rumen Microorganisms\***

By GLEN HALL, EDMUND W. CHENG AND WISE BURROUGHS

Numerous investigators have reported the presence of unidentified factors in certain feedstuffs and other materials which stimulate rumen microbial activity in vitro (1, 2, 3, 6, 8, 9, 10). Recently Ruf et al. (11) showed that extracts of yeast, cow manure and several feedstuffs stimulated the rate of cellulose digestion when they were added to a purified medium used for the in vitro culture of rumen microorganisms. Since many of the sources of the unidentified factors are known to contain several vitamins of the B group, it appears desirable to study the influence of certain of these vitamins on the rate of in vitro cellulose digestion. Preliminary reports of Hall et al. (7) and of Bentley et al. (1) seem to indicate that biotin, vitamin B<sub>12</sub> and possibly other vitamins exert favorable influences on the rate of cellulose digestion by rumen microorganisms. The present investigation is an attempt to find out the stimulatory effects, if any, of each of the B-vitamins or the combination of them on cellulose digestion by washed suspensions of rumen microorganisms.

### **EXPERIMENTAL**

To study the influence of vitamins on the cellulytic activity of rumen microorganisms it is essential that the medium as well as the inoculum used is free from such vitamins. For this purpose the washed suspension technique of Cheng et al. (4) was used. Briefly, this technique consists of separating the rumen microorganisms from the rumen fluid and partially digested feed debris by a process of differential centrifugation. Rumen contents obtained from a fistulated steer were strained through four layers of cheese cloth which constituted the first step in obtaining the rumen inoculum in liquid form. In a typical experiment a sample of 1200 ml. of rumen fluid was taken. After several processes of centrifugation and washing, the bacterial cells thus obtained were suspended in 600 ml. of a nutrient solution prepared according to the formula presented in Table 1. Three grams of Solka-Floc, a commercial

\*Journal paper No. J-2786 of the Iowa Agricultural Experiment Station, Ames, Iowa, Project No. 1208.

**Table 1**  
Composition of Basal Medium

Constituent	gm/2 liters	Constituent	gm/2 liters
Cellulose <sup>a</sup>	12.00	MgSO <sub>4</sub>	0.15
Urea	2.00	FeSO <sub>4</sub> .7H <sub>2</sub> O	0.075
Na <sub>2</sub> HPO <sub>4</sub> .H <sub>2</sub> O	1.20	CuSO <sub>4</sub> .5H <sub>2</sub> O	0.002
KH <sub>2</sub> PO <sub>4</sub>	0.60	CoCl <sub>2</sub>	0.002
NaHCO <sub>3</sub>	3.50	MnSO <sub>4</sub>	0.0004
KCl	0.75	ZnSO <sub>4</sub>	0.00008
NaCl	0.75		

<sup>a</sup>Solka-Floc—Brown Company, Berlin, New Hampshire

preparation of cellulose, were added to the suspension making the concentration of this insoluble cellulose 0.5 percent. The pH was adjusted to 7.0 with a saturated solution of sodium carbonate. Aliquots of 20 ml. each were pipetted into 75 ml. centrifuge tubes. The suspension was maintained anaerobic by passing through a constant stream of carbon dioxide gas. These tubes were placed in a constant temperature water bath at 39° C. At the end of 24 hours, the fermentation was terminated and the cellulose was determined on the entire contents of each tube using the procedure of Crampton and Maynard (5) with slight modifications.

Each of the B-vitamins was added to the rumen fermentation tubes at levels ranging from 0.1 to 100 micrograms. Various combinations of the B-vitamins were also studied. All the B-vitamins used were crystalline products. All treatments were triplicated.

### RESULTS AND DISCUSSION

In a basal medium containing no known B-vitamins but 0.5 percent of pure cellulose, approximately one-half of the cellulose was usually digested during an incubation period of 24-hr. For the convenience of comparison, the amount of cellulose digested for the basal group in an experiment was assigned a value of 100 and the amounts of cellulose digested for the test groups were computed proportionally to this basal value.

The results of the effects of vitamin B<sub>12</sub>, pseudo-vitamin B<sub>12</sub>, biotin, folic acid, para-aminobenzoic acid, pyridoxine and riboflavin on cellulose digestion by rumen microorganisms are shown in Table 2. All these vitamins appeared to increase the amounts of cellulose digested when added at levels ranging from 0.1 to 100

**Table 2**  
The Stimulatory effect of Certain B-Vitamins upon Cellulose Digestion by Rumen Microorganisms

B-Vitamin	Amount added $\mu\text{g}/20$ ml. medium	Cellulose digested (relative value)
B <sub>12</sub>	None	100
	1	109
	5	124
	10	116
Pseudo B <sub>12</sub>	None	100
	1	107
	5	119
	10	113
	25	126
	50	132
	100	127
Biotin	None	100
	0.1	123
	1	122
	10	124
Folic Acid	None	100
	1	121
	10	136
	100	142
para-Aminobenzoic acid	None	100
	1	109
	10	115
	100	122
Pyridoxine	None	100
	1	119
	10	119
	100	145
Riboflavin	None	100
	10	107
	100	126

micrograms per 20 ml. of washed suspension. Apparently these vitamins were required either by the cellulolytic microorganisms or other microorganisms in symbiosis with them.

The additions of either choline chloride, inositol, niacin, pantothenic acid or thiamine at levels ranging from 1 to 100 micro-

grams did not result in a significant increase in cellulose digestion. It appears that either these vitamins were not required by rumen microorganisms or that they were synthesized in sufficient quantities under the conditions of these experiments.

Further studies involved the addition of more than one individual vitamin into the medium. Various combinations of B-vitamins were tried but only the combination of vitamin B<sub>12</sub> and biotin was found to stimulate microbial cellulose digestion more than did either of these vitamins alone. This synergistic effect of vitamin B<sub>12</sub> and biotin is shown in Table 3. It is further shown in Table 3 that yeast

**Table 3**  
Effect of Biotin, Vitamin B<sub>12</sub>, and Yeast Extract Upon  
Cellulose Digestion by Rumen Microorganisms

Addition to basal medium	cellulose digested (relative value)
None	100
5 µg vitamin B <sub>12</sub>	115
1 µg biotin	112
5 µg vitamin B <sub>12</sub> + 1 µg biotin	124
50 mg. yeast extract	148
50 mg. yeast extract + 5 µg. vitamin B <sub>12</sub> + 1µg. biotin	148

extract was more stimulatory to cellulose digestion than the combination of biotin and vitamin B<sub>12</sub>. However, the addition of these two vitamins did not enhance the stimulatory effect of the yeast extract. Apparently the yeast extract contains additional cellulolytic factors other than biotin and vitamin B<sub>12</sub>.

The influences of the additions of other B-vitamins to the combination of vitamin B<sub>12</sub> and biotin into the basal medium on cellulose digestion are shown in Table 4. It is seen that folic acid, p-

**Table 4**  
Influence of Various B-Vitamins in Addition to B<sub>12</sub> and Biotin  
on Cellulose Digestion by Rumen Microorganisms

Addition to basal medium	Cellulose digested (relative value)
5 µg B <sub>12</sub> + 1 µg biotin	100
5 µg B <sub>12</sub> + 1 µg biotin + 10 µg folic acid	96
5 µg B <sub>12</sub> + 1 µg biotin + 1 µg inositol	103
5 µg B <sub>12</sub> + 1 µg biotin + 10 µg p-aminobenzoic acid	95
5 µg B <sub>12</sub> + 1 µg biotin + 100 µg pyridoxine	92
5 µg B <sub>12</sub> + 1 µg biotin + 10 µg riboflavin	108
5 µg B <sub>12</sub> + 1 µg biotin + 10 µg thiamine	97

aminobenzoic acid, pyridoxine and thiamine did not stimulate any increase in cellulose digestion. The slightly stimulatory effect

of inositol and riboflavin was probably not significant. It is possible that in the presence of B<sub>12</sub> and biotin, the microorganisms were able to synthesize other B-vitamins in sufficient quantities to supply their needs.

These data confirm our earlier report (7) and that of Bentley et al. (1) which demonstrated that biotin, vitamin B<sub>12</sub>, and para-aminobenzoic acid stimulated cellulose digestion by rumen microorganisms in vitro. The stimulatory action of the B-vitamins was not equal to that obtained from such sources as yeast extract and autoclaved rumen liquid. Thus, while these B-vitamins are not the complete answer to the unidentified cellulolytic factors, it would appear that they are at least partially responsible for the cellulolytic properties of such substances as yeast extract, rumen liquid, and possibly other sources of these factors.

It is not yet known whether the addition of these B-vitamins which stimulated cellulose digestion in vitro to ruminant rations would increase rumen microbial activity in vivo. Normally, rumen microorganisms are thought to synthesize sufficient quantities of the B-vitamins to meet their needs as well as that of their host. It is conceivable that these B-vitamins required by cellulolytic microorganisms may be supplied by the microbial synthesis of other types of rumen bacteria in as much as symbiotic relationships undoubtedly exist in the normal rumen. It should also be mentioned that throughout these studies carbohydrates which furnish readily available energy for microbial activity were not added into the medium. The effects of such materials upon in vitro cellulose digestion will be reported later.

#### SUMMARY

Certain B-vitamins were found to stimulate cellulose digestion by washed suspensions of rumen microorganisms. These were: Vitamin B<sub>12</sub>, pseudo-vitamin B<sub>12</sub>, biotin, para-aminobenzoic acid, pyridoxine, riboflavin, and folic acid. Of the combinations of vitamins studied, vitamin B<sub>12</sub> and biotin were the only vitamins in which the effect of one was synergistic to that of the other. Despite the stimulatory effect of these B-vitamins, no combination of them was found which stimulated cellulose digestion as much as that obtained from yeast extract. This substance apparently contains unidentified factors stimulatory to cellulose digestion in addition to the B-vitamins.

#### References

- (1) Bentley, O. G., Johnson, R. R., Vanecko, S., and Hunt, C. H. Studies on Factors Needed by Rumen Microorganisms for Cellulose Digestion In Vitro. *J. Animal Sci.*, 13:581. 1954.
- (2) Burroughs, W., Headley, H. G., Bethke, R. M., and Gerlaugh, P. Cellulose Digestion in Good and Poor Quality Roughages Using an Artificial Rumen. *J. Animal Sci.*, 9:514. 1950.

- (3) Burroughs, W., Latona, A., DePaul, P., Gerlaugh, P., and Bethke, R. M. Mineral Influences upon Urea Utilization and Cellulose Digestion by Rumen Microorganisms Using the Artificial Rumen Technique. *J. Animal Sci.*, 10:693. 1951.
- (4) Cheng, E. W., Hall, G., and Burroughs, W. A Method for the Study of Cellulose Digestion by Washed Suspensions of Rumen Microorganisms. In Press. *J. Dairy Sci.* 1955.
- (5) Crampton, E. W., and Maynard, L. A. The Relation of Cellulose and Lignin Content to the Nutritive Value of Animal Feeds. *J. Nutrition.* 15:383. 1938.
- (6) Doetsch, R. N., Robinson, R. Q., and Shaw, J. C. Techniques Employed in Cellural Investigations of the Bacteriology of Bovine Rumen Contents. *J. Animal Sci.*, 11:536. 1952.
- (7) Hall, G., Cheng, E. W., and Burroughs, W. B-vitamins and Other Factors Stimulatory to Cellulose Digestion by Washed Suspensions of Rumen Microorganisms. *J. Animal Sci.*, 12:918. 1953.
- (8) Huhtanen, C. N., Rogers, M. R., and Gall, L. S. Improved Techniques for Isolating and Purifying Rumen Organisms. *J. Bacteriol.*, 64:17. 1952.
- (9) Hungate, R. E. The Anaerobic Mesophilic Cellulolytic Bacteria. *Bacteriol Revs.*, 14:1. 1950.
- (10) McNeill, J. J., Doetsch, R. N., and Shaw, J. C. Some Nutritional Requirements of Bovine Rumen Bacteria. *J. Dairy Sci.*, 37:81. 1954.
- (11) Ruf, E. W., Hale, W. H., and Burroughs, W. Observations upon an Unidentified Factor in Feedstuffs Stimulatory to Cellulose Digestion in the Rumen and Improved Liveweight Gains in Lambs. *J. Animal Sci.*, 12: 731. 1953.

DEPARTMENT OF ANIMAL HUSBANDRY

IOWA STATE COLLEGE

AMES, IOWA