Proceedings of the Iowa Academy of Science

Volume 60 | Annual Issue

Article 60

1953

Generalizations of Continuity

Buchanan Cargal
Iowa State College

Let us know how access to this document benefits you

Copyright ©1953 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Cargal, Buchanan (1953) "Generalizations of Continuity," *Proceedings of the Iowa Academy of Science*, 60(1), 477-481.

Available at: <https://scholarworks.uni.edu/pias/vol60/iss1/60>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

Generalizations of Continuity

By BUCHANAN CARGAL

Let X be a Hausdorff space in which each neighborhood is an inexhaustible set (a set of Baire's second category), and let $\{N\}$ denote the system of neighborhoods in X . Thus $N(x)$ is a neighborhood of the element x . Let Y be a regular separable Hausdorff space and let $\{M\}$ denote the system of neighborhoods in Y . If f is a function on X into Y and ξ is a point of X , then f is continuous at ξ if for every $M(f(\xi))$ and for every $N(\xi)$, the set $N(\xi)E[x:f(x) \in M(f(\xi))]$ contains ξ as an interior point. A direct method for generalizing continuity is to weaken the requirements on the set $N(\xi)E[x:f(x) \in M(f(\xi))]$. For example, if $N(\xi)E[x:f(x) \in M(f(\xi))]$ contains an N (not necessarily an $N(\xi)$), then f is neighborly at ξ [1]¹. Other generalizations are obtained by requiring that $N(\xi)E[x:f(x) \in M(f(\xi))]$ contains an inexhaustible set, a set residual on a neighborhood (that is, a set whose complement on the neighborhood is exhaustible, or of Baire's first category), a non-denumerable set, or a non-null dense-in-itself subset. The purpose of this paper is to unify the study of generalized continuity by developing a concept which includes each of the generalizations mentioned above as special cases. It should be observed that properties of $N(\xi)E[x:f(x) \in M(f(\xi))]$ do not characterize functions which are semi-continuous or cliquish [2] at ξ .

DEFINITION I a. Let λ be a set property defined for subsets of X and let S be a subset of X . The point ξ is said to be a *point of λ approach by S* if for every $N(\xi)$, $N(\xi)S$ has property λ .²

b. The point ξ is said to be a *point of $k\lambda$ approach (concentrated λ approach) by S* if there exists an $N(\xi)$ such that for every N contained in $N(\xi)$, NS has property λ .

LEMMA I. For any property λ and for any set S in X , the set E of points of λ approach by S which are not points of $k\lambda$ approach by S is nowhere dense.

PROOF. If a neighborhood N contains a point of E , then there exists a neighborhood N_1 contained in N such that N_1S does not have property λ . Thus no element of N_1 belongs to E .

DEFINITION II a. Let f be a function on X into Y . The point

¹Numbers in brackets refer to the list of references.

²Definitions I and II are essentially those introduced by Blumberg [3] for the case when the property λ is inexhaustibility

ξ is said to be a *point of λ f -approach by S* if for every $M(f(\xi))$ and for every $N(\xi)$, $N(\xi)S \cap E[x:f(x) \in M(f(\xi))]$ has property λ .

b. The point ξ is said to be a *point of $k\lambda$ (concentrated λ) f -approach by S* if for every $M(f(\xi))$, there exists an $N(\xi)$ such that for every N contained in $N(\xi)$, $NSE[x:f(x) \in M(f(\xi))]$ has property λ .

DEFINITION III. The property λ is said to be an ascending set property if for every set S which has property λ and for every subset A of X , it is true that $(S+A)$ has property λ .

LEMMA II. Let λ be an ascending set property, let $f(x)$ be defined on X and let S be a subset of X . The set E of points of λ f -approach by S which are not points of $k\lambda$ f -approach by S is an exhaustible set.

PROOF. Let $\{M^r\}$ be a countable set of neighborhoods which are equivalent to $\{M\}$ and choose M_j^r from $\{M^r\}$. By Lemma I, the set E_j of points of λ approach by $SE[x:f(x) \in M_j^r]$ which are not points of $k\lambda$ approach by $SE[x:f(x) \in M_j^r]$ is nowhere dense. Thus $E_r = \sum_j E_j$ is exhaustible. We need only to show that $E \subseteq E_r$. Choose ξ in E . Since ξ is not a point of $k\lambda$ f -approach by S , there exists an $M_1(f(\xi))$ such that for every $N(\xi)$, there exists an N contained in $N(\xi)$ such that $NSE[x:f(x) \in M_1(f(\xi))]$ does not have property λ . Choose $M_k^r(f(\xi))$ from $\{M^r\}$ such that $M_k^r(f(\xi)) \subseteq M_1(f(\xi))$. Since λ is ascending, $NSE[x:f(x) \in M_k^r(f(\xi))]$ does not have property λ . Hence $\xi \in E_r$. Thus E is exhaustible.

COROLLARY 1. The set of points of inexhaustible f -approach by X which are not points of concentrated inexhaustible f -approach by X is an exhaustible set [3].

Let ρ metrize Y and let $M(f(\xi); \epsilon)$ denote the set $E[y:\rho(y, f(\xi)) < \epsilon]$.

THEOREM I. Let λ be an ascending set property and S be a set in X . If the sequence of functions $\{f_n(x)\}$ is defined on X and converges uniformly to $F(x)$, and if there exists a point ξ in S such that for every n , ξ is a point of λ f_n -approach by S , then ξ is a point of λ F -approach by S .

PROOF. Let $M(F(\xi))$ and $N(\xi)$ be given. We must show that $N(\xi)SE[x:F(x) \in M(F(\xi))]$ has property λ . Choose a positive number ϵ such that $M(F(\xi); \epsilon) \subseteq M(F(\xi))$ and choose n such that for every x in S , $\rho(f_n(x); F(x)) < \epsilon/3$. Now $N(\xi)SE[x:f_n(x) \in M(\xi; \epsilon/3)]$ has property λ . Suppose $\eta \in N(\xi)SE[x:f_n(x) \in M(\xi; \epsilon/3)]$. Then $\rho(f_n(\eta); f_n(\xi)) < \epsilon/3$. Hence $\rho(F(\xi); F(\eta)) \leq \rho(F(\xi); f_n(\xi)) + \rho(f_n(\xi); f_n(\eta)) + \rho(f_n(\eta); F(\eta)) < \epsilon$. Thus $\eta \in N(\xi)SE$

$[x:F(x) \in M(F(\xi))]$. Since λ is an ascending set property, $N(\xi)SE[x:F(x) \in M(F(\xi))]$ has property λ .

COROLLARY II. *Let λ be the property that a set contains ξ as an interior point. Then Theorem I yields the well-known result that the uniform limit of a sequence of functions which are continuous at ξ is a function which is continuous at ξ .*

For other particular results, λ may be taken to be the property that a set contain a neighborhood, that a set be infinite, or any other ascending set property.

Let the complement with respect to X of a set S be denoted by $S^{\sim}$ and the closure of S be designated cS .

DEFINITION IV. The property λ is said to be a *locally characterizing set property* if for every set S which has property λ and for every N such that NS has property λ , there exists an N_1 contained in N such that N_1S has property λ and $N_1(S^{\sim})$ does not have property λ .

For example, the property that a set be residual on a neighborhood is a locally characterizing set property while the property that a set be inexhaustible is not.

LEMMA III. *Let λ be an ascending locally characterizing set property. Let N be a neighborhood and R and S be sets in X such that NRS has property λ . Then there exists an N_1 contained in N such that $N_1R(S^{\sim})$ does not have property λ .*

PROOF. NS has property λ because λ is ascending. But since λ is locally characterizing, there exists an N_1 contained in N such that $N_1(S^{\sim})$ does not have property λ . From the ascending property of λ , it now follows that $N_1R(S^{\sim})$ does not have property λ .

LEMMA IV. *Let λ be an ascending locally characterizing set property. Let f be a function on X onto a subset of Y and let each element x of a set S in X be a point of λ f -approach by S . If ξ is a point of λ f -approach by S , then for every $M(f(\xi))$ and for every $N(\xi)$, there exists an N contained in $N(\xi)$ such that for every x in NS , $f(x) \in M(f(\xi))$.*

PROOF. Let $M(f(\xi))$ and $N(\xi)$ be given. Choose $M_1(f(\xi))$ such that $cM_1(f(\xi)) \in M(f(\xi))$. Now $N(\xi)SE[x:f(x) \in M_1(f(\xi))]$ has property λ . By Lemma III, there exists an N contained in $N(\xi)$ such that $NSE[x:f(x) \text{ not } \in M_1(f(\xi))]$ does not have property λ . Suppose η , a point of NS , is such that $f(\eta) \text{ not } \in M(f(\xi))$. Choose $M(f(\eta))$ disjoint from $M_1(f(\xi))$. Now $NSE[x:f(x) \in M(f(\eta))]$ has property λ . Since λ is ascending $NSE[x:f(x) \text{ not } \in M_1(f(\xi))]$ has property λ , a contradiction. Hence $f(\eta) \in M(f(\xi))$.

COROLLARY III. *Since ρ metrizes the space Y , under the conditions of Lemma IV, for every positive number ϵ , and for every $N(\xi)$, there exists an N contained in $N(\xi)$ such that for every x in NS , $\rho(f(x), f(\xi)) < \epsilon$.*

THEOREM II. *Let λ be an ascending locally characterizing set property. Let $f(x)$ be defined on X and let each element x of a set S in X be a point of λ f -approach by S . If ξ is a point of $k\lambda$ f -approach by S , then $f(x)$ is continuous at ξ with respect to S .*

PROOF. Let $M(f(\xi))$ and $N(\xi)$ be given. Choose $M_1(f(\xi))$ such that $\epsilon M_1(f(\xi)) \in M(f(\xi))$. Since ξ is a point of $k\lambda$ f -approach by S , there exists an $N_1(\xi)$ in $N(\xi)$ such that for every N in $N_1(\xi)$, $NS\epsilon[x: f(x) \in M_1(f(\xi))]$ has property λ . Suppose η , a point of $N_1(\xi)S$, is such that $f(\eta) \notin M(f(\xi))$. Choose $M(f(\eta))$ disjoint from $M_1(f(\xi))$. Now $N_1(\xi)SE[x: f(x) \in M(f(\eta))]$ has property λ . By Lemma III, there exists an N_1 in $N_1(\xi)$, such that $N_1SE[x: f(x) \notin M(f(\eta))]$ does not have property λ . Since λ is ascending, $N_1SE[x: f(x) \in M_1(f(\xi))]$ does not have property λ , a contradiction. Thus $f(\eta) \in M(f(\xi))$. Hence $f(x)$ is continuous at ξ with respect to S .

Using Lemma II and Theorem II, we may state the following theorem.

THEOREM II'. *If λ is an ascending locally characterizing set property, if $f(x)$ is defined on X , and if each element x of S is a point of λ f -approach by S , then $f(x)$ is continuous over S with respect to S except on an exhaustible set.*

THEOREM III. *Let the sequence of functions $\{f_n(x)\}$ be defined and converge to $F(x)$ on X . Let λ be an ascending locally characterizing set property. Let R be a set residual in X and let it be required that for every n , each x in R is a point of λ f_n -approach by R . Then $F(x)$ is continuous over R with respect to R except on an exhaustible set.*

REMARK. If we say that a set has property λ if it contains a neighborhood and let $R = X$, we obtain Bledsoe's results for neighborhoodly functions [2]. The proof of Theorem III is similar to the proof given by Bledsoe, but we include it here because our theorem is more general.

PROOF. By Theorem II', $f_n(x)$ is continuous on a residual set R_n with respect to R . Thus for every n , $f_n(x)$ is continuous on the residual set $R' = \Pi_n R_n$ with respect to R . Let $w(x) = \limsup \eta \rightarrow x \rho(F(\eta), F(x))$ for η 's in R if this limit exists, and let $w(x) = 1$ at all other points of R . Let n and p be positive integers and

let $E_p^n = E[x \in R' : w(x) \geq 1/p \text{ and for every } m \geq n, \rho(f_m(x), F(x)) < 1/7p]$. Suppose there are integers n and p and a neighborhood N such that E_p^n is dense on N . Choose ξ in $E_p^n N$. Then $\rho(f_n(\xi), F(\xi)) < 1/7p$. Since $f_n(x)$ is continuous at ξ with respect to R , there exists an $N(\xi)$ contained in N such that for every x in $N(\xi)R$, $\rho(f_n(x), f_n(\xi)) < 1/7p$. Since $w(\xi) \geq 1/p$, there exists an x_1 in $N(\xi)R$ such that $\rho(F(\xi), F(x_1)) \geq 6/7p$. Choose $m \geq n$ such that $\rho(f_m(x_1), F(x_1)) < 1/7p$. Since x_1 is a point of λ f_m -approach by R , using Corollary III, we can conclude that there exists an N_1 contained in $N(\xi)$ such that for every x in N_1R , $\rho(f_m(x), f_m(x_1)) < 1/7p$. In particular let x_2 be contained in $N_1E_p^n$. Then $\rho(f_m(x_2), f_m(x_1)) < 1/7p$. But $\rho(f_m(x_2), f_n(x_2)) < 2/7p$. We have $\rho(F(\xi), F(x_1)) \leq \rho(F(\xi), f_n(\xi)) + \rho(f_n(\xi), F(x_1)) + \rho(f_n(\xi), f_n(x_2)) + \rho(f_n(x_2), f_m(x_2)) + \rho(f_m(x_2), f_m(x_1)) < 6/7p$, a contradiction. Thus $E = \sum_{n,p} E_p^n$ is exhaustible and $F(x)$ is continuous on $(R' - E)$ with respect to R .

References

1. W. W. Bledsoe, *Neighborly Functions*, Proc. Amer. Math. Soc., Vol. 3, No. 1 (Feb. 1952) pp. 114-115.
2. H. P. Thielman, *Types of Functions*, Amer. Math. Monthly, Vol. 60 (1953) pp. 156-161. Also see *Pathological Functions*, Proc. Iowa. Academy of Science, Vol. 59, (1952) pp. 338-343.
3. H. Blumberg, *Arbitrary Point Transformations*, Duke Math. Jour., Vol. 11 (1944) pp. 671-685.

DEPARTMENT OF MATHEMATICS

IOWA STATE COLLEGE

AMES, IOWA