

1950

## Bernoulli Numbers

R. B. McClenon  
*Grinnell College*

*Let us know how access to this document benefits you*

Copyright ©1950 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

---

### Recommended Citation

McClenon, R. B. (1950) "Bernoulli Numbers," *Proceedings of the Iowa Academy of Science*, 57(1), 315-319.

Available at: <https://scholarworks.uni.edu/pias/vol57/iss1/39>

This Research is brought to you for free and open access by the IAS Journals & Newsletters at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact [scholarworks@uni.edu](mailto:scholarworks@uni.edu).

**Offensive Materials Statement:** Materials located in UNI ScholarWorks come from a broad range of sources and time periods. Some of these materials may contain offensive stereotypes, ideas, visuals, or language.

## Bernoulli Numbers

By R. B. McCLENON

The first appearance of the set of rational numbers of which I am speaking was in James Bernoulli's *Ars Conjectandi*. This work was published in 1713, eight years after the death of its author. In the 2nd chapter, Bernoulli deals with permutations and combinations. He starts with the formula

$$C(n, k) = C(1, k-1) + C(2, k-1) + \dots + C(n-1, k-1)$$

where naturally

$$c(r, s) = 0 \text{ for } r < s;$$

or

$$\frac{n(n-1)\dots(n-k+1)}{k!} = \sum_{h=1}^n \frac{(h-1)(h-2)\dots(h-k+1)}{(k-1)!}$$

For  $k = 3$ , for instance,

$$\frac{n(n-1)(n-2)}{3!} = \sum_{h=1}^n \frac{(h-1)(h-2)}{2!} =$$

$$\sum \left( \frac{1}{2}h^2 - \frac{3}{2}h + 1 \right) = \frac{1}{2} \sum h^2 - \frac{3}{2} \sum h + n$$

$\sum n$  was of course known, long before Bernoulli's time, to equal  $\frac{n(n+1)}{2}$ .

$$\begin{aligned} \therefore \sum h^2 &= \frac{n(n-1)(n-2)}{3} + \frac{3n(n+1)}{2} - 2n \\ &= \frac{1}{3}n^3 + \frac{1}{2}n^2 + \frac{1}{6}n \end{aligned}$$

Likewise, for  $k = 4$ ,

$$\frac{n(n-1)(n-2)(n-3)}{4!} = \sum_{h=1}^n \frac{(h-1)(h-2)(h-3)}{3!}$$

$$= \frac{1}{6} \sum n^3 - \sum n^2 + \frac{11}{6} \sum n - n$$

from which follows

$$\sum n^3 = \frac{1}{4} n^4 + \frac{1}{2} n^3 + \frac{1}{4} n^2$$

Bernoulli gives thus a table of the values of  $\sum n^c$  (the notation he used) as far as  $c=10$ . Thereupon appears the remarkable generalization:

$$\sum n^c = \frac{n^{c+1}}{c+1} + \frac{n^c}{2} + \frac{c}{2} A n^{c-1}$$

$$+ \frac{c(c-1)(c-2)}{2 \cdot 3 \cdot 4} B n^{c-3}$$

$$+ \frac{c(c-1)(c-2)(c-3)(c-4)}{2 \cdot 3 \cdot 4 \cdot 5 \cdot 6} C n^{c-5} + \dots$$

in which  $A = 1/6$ ,  $B = -1/30$ ,  $C = 1/42$ ,  $D = -1/30$ ,  $E = 5/66 \dots$ , the first five of the numbers  $B_2, B_4, B_6 \dots$ . Bernoulli does not indicate how he obtained this generalization. It seems, however, from the coefficients which appear in the formulas as far as  $\sum n^{10}$ , that the discovery of the first five numbers should not be too difficult. But to arrive at any general formula, or recurrence relation, seems hopeless by merely following Bernoulli's tracks.

Euler took a different line, and with his tremendous power of computation managed to determine the value of the first 15 non-vanishing values.

Passing on at once to a relatively simple method for computing the numbers, the following has been published in various places and is perhaps the one most frequently used:

Starting with the function  $\frac{x}{e^x - 1}$ , which is certainly analytic in

the vicinity of  $x=0$  if the value of  $f(0)$  is defined to be unity, we may write

$$\frac{x}{e^x - 1} = \sum_{n=0}^{\infty} \frac{B_n x^n}{n!} = B_0 + \frac{B_1 x}{1!} + \frac{B_2 x^2}{2!} + \dots \tag{1}$$

Thus,

$$1 = \left(1 + \frac{x}{2!} + \frac{x^2}{3!} + \dots\right) \left(B_0 + \frac{B_1}{1!} x + \frac{B_2}{2!} x^2 + \dots\right)$$

so that  $B_0 = 1$ ,  $B_1 = -\frac{1}{2} \dots$  and in general ( $n > 1$ )

$$\begin{aligned} \frac{1}{n!} B_0 + \frac{1}{(n-1)!} \frac{B_1}{1!} + \frac{1}{(n-2)!} \frac{B_2}{2!} + \dots \\ + \frac{1}{1!} \frac{B_{n-1}}{(n-1)!} = 0 \end{aligned}$$

If we multiply by  $n!$ , we get

$$\binom{n}{0} B_0 + \binom{n}{1} B_1 + \binom{n}{2} B_2 + \dots + \binom{n}{n-1} B_{n-1} = 0$$

or symbolically,  $(B + 1)^n - B^n = 0$ , replacing  $B_k$  by  $B^k$ . Returning to

$$\begin{aligned} f(x) &= \frac{x}{e^x - 1}, \quad f(x) + \frac{x}{2} = \frac{x}{2} \left(1 + \frac{2}{e^x - 1}\right) \\ &= \frac{x}{2} \frac{e^x + 1}{e^x - 1} = \frac{x}{2} \frac{e^{\frac{x}{2}} + e^{-\frac{x}{2}}}{e^{\frac{x}{2}} - e^{-\frac{x}{2}}} \end{aligned} \tag{2}$$

which is an even function. Thus  $B_3 = B_5 = \dots = B_{2k-1} = 0$  and for a few of the others:

for  $n = 2, (B + 1)^2 - B^2 = 2B_1 + 1 = 0$   
 $n = 3, 3B_2 + 3B_1 + 1 = 0 \quad B_2 = 1/6$   
 $n = 4, 4B_3 + 6B_2 + 4B_1 + 1 = 0 \quad B_3 = 0$  (known)  
 $n = 5, 5B_4 + 10B_3 + 10B_2 + 5B_1 + 1 = 0 \quad B_4 = -1/30$   
 $n = 7, 7B_6 + 21B_5 + 36B_4 + 35B_3 + 21B_2 + 7B_1 + 1 = 0$   
 $B_6 = 1/42$

The case  $n = 125$  would seem to be a good problem for one of the super-machines now being constructed. For this would check what is, so far as I know, the last Bernoulli number to have been computed.

The uses of the Bernoulli numbers are almost innumerable. Among the most interesting are the series expansions for  $\tan x$ ,  $\cot x$ , and  $\csc x$ . These are found without much difficulty, starting with (2). The results are

$$\begin{aligned} x \cot x &= 1 - \frac{2^2 B_2}{2!} x^2 + \frac{2^4 B_4}{4!} x^4 - + \dots \\ &\quad + (-1)^k \frac{2^{2k} B_{2k}}{(2k)!} x^{2k} + \dots \\ &= 1 - \frac{1}{3} x^2 - \frac{1}{45} x^4 - \frac{2}{945} x^6 - \dots \end{aligned} \tag{3}$$

$$\begin{aligned} \tan x &= \sum_{k=1}^{\infty} (-1)^{k-1} \frac{2^{2k} (2^{2k} - 1) B_{2k}}{(2k)!} x^{2k-1} \\ &= x + \frac{1}{3} x^3 + \frac{2}{15} x^5 + \frac{17}{315} x^7 + \dots \end{aligned} \tag{4}$$

$$\begin{aligned} x \csc x &= \sum_{k=0}^{\infty} (-1)^{k-1} \frac{(2^{2k} - 2) B_{2k}}{(2k)!} x^{2k} \\ &= 1 + \frac{1}{6} x^2 + \frac{7}{360} x^4 + \frac{31}{15120} x^6 + \dots \end{aligned} \tag{5}$$

We have so far assumed that the  $B_{2n}$  in the series for  $\frac{x}{e^x - 1}$  were the same  $B_{2n}$  which Bernoulli gave as A, B, C . . . in the expansion of  $\sum n^x$  in powers of  $n$ . The proof of this is not very difficult, since we have Euler's expansion of  $\cot x$  in a series of partial fractions. From this we are led by a direct road to the identity

$$\frac{1}{e^x - 1} = \frac{1}{x} - \frac{1}{2} + \sum_{n=1}^{\infty} \frac{2x}{x^2 + 4n^2\pi^2}$$

Now for  $|x| < 2\pi$  each fraction can be expanded in powers of  $x$ , giving

$$\frac{2x}{x^2 + 4n^2\pi^2} = \frac{x}{2n^2\pi^2} \left[ 1 - \frac{x^2}{4n^2\pi^2} + \frac{x^4}{4^2n^4\pi^4} - + \dots \right] \tag{6}$$

The double series (6) being absolutely convergent within its circle of convergence, can be summed "vertically" in powers of  $x$ , so that

$$\frac{1}{e^x - 1} = \frac{1}{x} - \frac{1}{2} + \frac{x}{2\pi^2} \left[ \frac{1}{n^2} - \frac{x^3}{2^4\pi^4} \left[ \frac{1}{n^4} \dots \right] \right]$$

Thus

$$B_{2n} = (-1)^{n-1} \frac{(2n)!}{2^{2n-1} \pi^{2n}} \sum_1^{\infty} \frac{1}{n^{2n}}$$

and we have the connection between the Bernoulli numbers as obtained by their discoverer, and the coefficients in the expansion of

$\frac{x}{e^x - 1}$ . Incidentally we note that the successive Bernoulli numbers increase greatly as  $r$  increases.

DEPARTMENT OF MATHEMATICS  
GRINNELL COLLEGE