

1949

A Bignell (?) Loess Section in Western Iowa

Robert V. Ruhe
Iowa State College

Let us know how access to this document benefits you

Copyright ©1949 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Ruhe, Robert V. (1949) "A Bignell (?) Loess Section in Western Iowa," *Proceedings of the Iowa Academy of Science*, 56(1), 229-231.

Available at: <https://scholarworks.uni.edu/pias/vol56/iss1/30>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

A Bignell(?) Loess Section in Western Iowa

By ROBERT V. RUHE

INTRODUCTION

The Bignell loess was named and first described by Schultz and Stout (1945). At the type section of the Bignell formation, near Bignell, Lincoln County, Nebraska, the upper loess (Bignell) is separated from the Peorian loess by a prominent soil and overlain by a complex topsoil.

Condra, Reed, and Gordon (1947) have also described the Bignell loess. Various profile sections in this paper (figures 3, 7, 12 and 14) show that the Bignell loess is restricted to flood plains, terraces, and upland shoulders of the valleys of the eastern half of Nebraska.

Schultz and Stout (p. 243) have proposed the following correlation of the Bignell loess:

W-3 Mankato	Soil Y Bignell loess
Two Creeks forest bed	Soil X
W-2 Tazewell-Cary	
"Peorian"	"Peorian" cycle complex
W-1 Iowan	

Condra, Reed, and Gordon concur in this interpretation.

The Bignell loess has not been previously recognized in Iowa. The lack of a soil break in the buff loess of western and southwestern Iowa has been somewhat perplexing when one realizes that on the western side of the Missouri River valley in the vicinity of Blair, Nebraska, the soil separating the Bignell and Peorian loesses is well developed.

SECTION IN WESTERN IOWA

In the SC SW $\frac{1}{4}$ sec. 8, T. 81N., R. 44W., Harrison County, Iowa a Bignell(?) - Peorian loess section is located. The loesses are exposed in a vertical face near the top of a bluff approximately 100 feet above the grade of the Pisgah road. A road cut at the base of this bluff shows Pleistocene sands and gravels; this road cut is located approximately two-tenths of a mile northeast of the intersection of the Pisgah road (county trunk D) and county trunk V.

The description of the section is as follows:

- | | |
|---|---|
| <ol style="list-style-type: none"> 3. Bignell(?) loess : approximately
 Modern soil profile, 30 inches
 Buff, calcareous, well sorted loess, 10 feet
 Buff, calcareous, fossiliferous, well sorted loess, 12 feet 2. Peorian loess : exposed
 Dark gray to black, slightly calcareous, well sorted
 silt (A horizon of buried soil profile), 12 inches
 Light gray to buff, heavier textured, leached silt
 (B horizon), 15 inches
 Buff, slightly calcareous, well sorted loess
 (C horizon), 9 inches 1. Slump | <p>Thickness
24½ feet</p> <p>3 feet</p> |
|---|---|

An analysis of the clay and organic contents of the buried soil profile (Fig. 1) shows the horizon differentiation. Accumulation of organic material has occurred in the A horizon ; slight alluviation of clay has occurred in the B horizon. There is secondary carbonate in the A horizon (from overlying calcareous loess) ; the B horizon is leached ; the C horizon contains primary carbonate material.




Fig. 1. Buried Soil "X" profile analysis.

A somewhat marked physical dissimilarity exists between the lower buff loess of this described section and the Loveland loess. The Loveland loess is exposed at several locations in the general area of the Pisgah road section ; it is generally differentiated from the buff loess by its pink to red color and heavier mass texture.

The contact between the lower buff loess and the Loveland loess was not determined. A more detailed study of this area is necessary in order to show the exact stratigraphic relationship. The loess below the buried soil in the Pisgah road section, however, does not have the color or the texture characteristic of the Loveland loess in that area.

CONCLUSION

The similarity of the above described section to the Bignell type section, as described by Schultz and Stout, the observation by the writer of similar loess sections in eastern Nebraska which are designated as Bignell-Peorian, and the existence of the soil interbedded in buff loess, which has been previously recognized as Peorian loess, indicate that the loess overlying the buried soil in the Pisgah road section is probably Bignell loess.

Literature Cited

- Condra, G. E., Reed, E. C., and Gordon, E. D. 1947. Correlation of the Pleistocene deposits of Nebraska. Nebraska Geol. Survey Bull. 15, pp. 13, 33.
- Schultz, C. B. and Stout, T. M. 1945. Pleistocene loess deposits of Nebraska. Am. Jour. Sci., Vol. 243, pp. 241, 243.

DEPARTMENT OF GEOLOGY
IOWA STATE COLLEGE
AMES, IOWA