

1945

A Preliminary Report on the Geometridae of Iowa

Elizabeth Jerrel
Iowa Wesleyan College

H. E. Jaques
Iowa Wesleyan College

Copyright © Copyright 1945 by the Iowa Academy of Science, Inc.
Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Jerrel, Elizabeth and Jaques, H. E. (1945) "A Preliminary Report on the Geometridae of Iowa," *Proceedings of the Iowa Academy of Science*: Vol. 51: No. 1 , Article 52.
Available at: <https://scholarworks.uni.edu/pias/vol51/iss1/52>

This Research is brought to you for free and open access by UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

A PRELIMINARY REPORT ON THE GEOMETRIDAE OF IOWA

ELIZABETH JERREL AND H. E. JAQUES

For the convenience of anyone desiring information on the family Geometridae, order Lepidoptera, the following list of species reported from Iowa has been prepared. The Geometridae are universally distributed in all parts of the world where vegetation is found. They are rather frail in structure, with considerable expanse of wing in relation to the size of the body. They are either semi-diurnal or crepuscular. Over 800 species of Geometrids are known to occur within the limits of the United States and Canada, of which 61 have been found in Iowa. This number could doubtless be greatly enlarged by more intensive collecting.

The larvae are commonly known as measuring worms or loopers, because, the anterior pairs of prolegs being totally aborted, they progress by bringing the posterior somites close to the anterior ones and looping the median somites. This, combined with the elongated form, causes these larvae when at rest to assume positions in which the body extends from the posterior prolegs in a twig-like manner. The front parts are not really free, however, as they are connected by a slender thread of silk to some point nearby. The larvae also suspend themselves from branches on a long thread of silk to escape perching birds. Dichromatism is often revealed among the larvae, part of a brood being green, the remainder brown or yellowish.

In some species of this family the adult females are without wings, such as the spring and fall cankerworms: *Paleacrita vernata* Peck. and *Alsophila pomataria* Har. This characteristic considerably facilitates their control, the females, after emerging from pupae in the ground, being caught in tanglefoot or some other similar substance when they attempt to climb the trees to deposit their eggs.

Haematopsis grataria Fab., the Chickweed Moth, (Fig. 1.) is very common, and is often seen by the roadsides where it has the habit of clinging to grasses and flying up when disturbed. The larvae feed upon *Stellaria media* (L.), the common chickweed. It ranges from the Atlantic to the Mississippi and beyond. *Euchlaena obtusaria* Hub., (Fig. 2) is, like the preceding species, a native of the eastern half of the continent.

For some twenty years the department has been making general collections of insects throughout the state in an effort to determine their geographical and seasonal distribution. The accumulated information of this specific group must be recognized as only fragmentary. It should be added that the Order Lepidoptera has not been as intensively studied as has several other orders.




Fig. 1. *Haematopsis grataria* Fab.




Fig. 2. *Euchlaena obtusaria* Hub.

- 3901—PTYCHOPODA INDUCTATA
GN.
- 4020—DREPANA ARCUATA WLK.
- 4026—ALSOPHILA POMETARIA
HARRIS
- 4046—NEMORIA BISTRIARIA HBN.
- 4087—CHLORISSA PISTASCIARIA
GN.
- 4095—CHLOROCHLAMYS CHLORO-
LEUCARIA GN.
- 4149—SCOPULA ENUCLEATA GN.
- 4158—S. INDUCTATA GN.
- 4204—HAEMOTOPIS GRATARIA
FABR.
- 4205—TIMANDRA AMATURARIA
WLK.
- 4234—DYSPTERIS ABORTIVARIA
H. S.
- 4401—LYGRIS DIVERSILINEATA
HBN.
- 4535—PERCNOPTILOTA OBSTIPATA
FABR.
- 4559—EUPHYIA CENTROSTRIGARIA
WOLL.
- 4573—EULYPE HASTATA L.
- 4599—EUDELE MENDICA WLK.
- 4603—E. UNICOLOR ROB.
- 4605—BAPTA SEMICLARATA WLK.
- 4606—B. VESTALIATA GN.
- 4640—PTEROSPODA OPUSCULARIA
HLST.
- 4654—FERNALDELLA FIMETARIA
G&H.
- 4657—MELLILLA ANTHOMETATA
WLK.
- 4663—PHYSOSTEGANIA PUSTU-
LARIA GN.
- 4665—PHILOBIA AEMULATARIA
WLK.
- 4688—SEMIOTHISA DENTICULATE
GRT.
- 4723—S. OCELLINATA G&.
- 4726—S. MELLISTRIGATA GRT.
- 4735—S. SNOVIATA PACK.
- 4739—S. S-SIGNATA PACK.
- 4743—S. TENEBROSATA HLST.
- 4746—ITAME RIBEARIA FITCH
- 4751—I. SULPHUREA PACK.
- 4756—I. COORTARIA HLST.
- 4774—I. LATIFERRUGATA WLK.
- 4807—PARAPHIA UNIPUNCTATA
HAW.
- 4812—TORNOS SCOLOPACINARIUS
GN.
- 4908—ANAVITRINELLA PAM-
PINARIA GN.
- 4918—ANACAMPTODES VELLIVO-
LATA HLST.
- 4952—NACOPHORA YPSILON
FORBES
- 4958—PHIGALIA TITEA CRAM.
- 4960—PALEACRITA VERNATA PECK.
- 4966—LYCIA URSARIA WLK.
- 4968—AMPHIDASIS COGNATARIA
GN.
- 4991—EUGONOBAPTA NIVOSARIA
GN.
- 4993—LYTROSIS UNITARIA H. S.
- 4994—EUCHLAENA SERRATA DRU.
- 4995—E. OBTUSARIA HBN.
- 4997—E. JOHNSONARIA FIRCH.
- 5001—E. ASTYLUSARIA WLK.
- 5005—E. TIGRINARIA GN.
- 5007—XANTHATYPE SOSPETA
DRU.
- 5010—X. URTICARIA SWETT.
- 5044—NEMATOCAMPA LIMBATA
HAW.
- 5054—METANEMA INATOMARIA
GN.
- 5059—ENNOMOS SUBSIGNARIUS
HBN.
- 5072—PERO HONESTARIUS WLK.
- 5082—P. MARMORATUS GROSSE.
- 5161—SICYA MACULARIA HARR.
- 5170—DEUTERONOMOS MAGNARIUS
GN.
- 5184—APICIA CONFUSARIA HBN.
- 5211—PROCHOERODES TRANS-
VERSATA DRURY.

IOWA WESLEYAN COLLEGE
MT. PLEASANT, IOWA