

1945

A List of the Chloropidae (Diptera) Known to Occur in Iowa

Leonard Redlinger
Iowa Wesleyan College

H. E. Jaques
Iowa Wesleyan College

Let us know how access to this document benefits you

Copyright ©1945 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Redlinger, Leonard and Jaques, H. E. (1945) "A List of the Chloropidae (Diptera) Known to Occur in Iowa," *Proceedings of the Iowa Academy of Science*, 51(1), 471-472.

Available at: <https://scholarworks.uni.edu/pias/vol51/iss1/54>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

A LIST OF THE CHLOROPIDAE (DIPTERA) KNOWN TO OCCUR IN IOWA

LEONARD REDLINGER AND H. E. JAQUES

The Dipterous family Chloropidae is composed of small to very small, bare or nearly bare flies. These frit-flies as they are called are very numerous and representatives of the family may be collected almost anywhere. Adults of a considerable number of species are common in meadows and other places where there is rank growing grass and several of them are of considerable importance economically. They may be readily collected with a sweeping net in any of their ideal habitats.

The larvae often live in the stems of our cultivated and wild grasses as well as some other plants. Occasionally species feed within the burrows in plants made by other insects. A few feed upon the eggshells and cast-off skins of insects. These larvae are thick and cylindrical, with stout mouth hooks, and antennae of two segments. They move about by fleshy abdominal protuberances.

The adults of this family are minute but plump, and are variously colored, sometimes blackish, sometimes yellowish or both. The wings are of moderate length or very short. The head is usually hemispherical, but sometimes more or less triangular. The antennae are usually short, with rounded third segment. The unusually large frontal or vertical triangular offers one of the best characters for recognizing the family. The legs are short, and the femora rarely greatly thickened, but usually somewhat curved.


The figure of *Meromyza americana*, the Wheat Stem Maggot, illustrates the general characters and appearance of the members of the family.

The chief purpose of this paper is to list the known Iowa species. Specimens of all of these are found in the Iowa Insect Survey collection at Iowa Wesleyan College. The numbers accompanying the names are the catalog numbers assigned these species in a special catalog of Iowa Diptera.

210-1	MEROMYZA AMERICANA FITCH	211-5	C. MELANOCERA Loew
		211-6	C. OBSCURICORNIS Loew
210-21	EPICHLOROPS EXILIS (COQ.)	211-11	C. CROCOTA Loew
		211-12	C. PALPALIS Adams
210-31	CHLOROPISCA GLABRA MEIG.	211-13	C. RUGINOSA Becker
		211-14	C. CINERAPENNIS Adams
210-31a	C. GLABRA var. CLYPEATA Mall.	211-15	C. RUFESCENS Coq.
		211-31	ECTECEPHALA ALBISTYLUM Macq.
210-51	DIPLOTOXA ANCHORA Lw.		E. SULCATA Sabr.
210-53	D. VERSICOLOR Lw.	211-32	
210-92	ANTHRACOPHAGA INTERRUPTA Becker	211-41	PARECTECEPHALA EUCERA Lw.
		211-81	GAURAX DORSALIS Lw.
211-2	CHLOROPS CERTIMA Adams	212-2	CRASSISETA COSTATA Lw.

212-2a	C. COSTATA var. (red pleura)	213-2a	O. COXENDIX var. PULICORNIS Sab.
212-3	C. NIGRICEPS Lw.	213-3	O. COXENDIX OBSCURA Coq.
212-4	C. LONGULA Lw.	213-5	O. FRIT L.
212-5	C. EUNOTA Lw.	213-5a	O. FRIT var. NITIDISSIMA Mg.
212-43	HIPPELATES NOBILIS Loew	213-5b	O. FRIT var. "B"
212-44	H. PLEBEJUS Lw.	213-5c	O. FRIT var. "A"
212-45	H. SUBVITTATUS Mall.	213-6	O. INCERTA Becker
212-46	H. PALLIPES Lw.	213-10	O. UMBROSA Lw.
212-47	H. STRAMINEA Lw.	213-11	O. LONGIPES Lw.
212-48	H. DISSIDENS Tucker	213-12	O. ABDOMINALIS Becker
212-49	H. BISHOPII Sabr.	213-13	O. MINOR Adams
212-61	MADIZA CINEREA Lw.	213-14	O. AINSLIEI Cur.
212-61 var.	M. CINEREA var.	213-15	O. MELANCHOLICA Becker
212-63	M. SETULOSA Mall.	213-16	O. LUTICEPS Sabrosk.
212-64	M. PROVOCANS Beck.	213-17	O. SCABRA Ald.
212-65	M. LATIFRONS Lw.	213-18	O. FLAVESCENS Tucker
212-66	M. NIGRIPALPIS Mall.	213-19	O. NUDIUSCULA Loew
212-67	M. QUINQUELINEATA Adams	213-21	DICRAEUS INCONGRUUS Ald.
212-68	M. PARVA Ad.	213-31	TRICIMBA LINEELLA Fall.
213-2	OSCINELLA COXENDIX Fitch.	213-32	T. BRUNNICOLLIS Beck.

IOWA WESLEYAN COLLEGE
Mt. PLEASANT, IOWA



Meromyza americana Fitch