

1941

The Use of Tetrachlorethane in the Eradication of the European Bindweed (Abstract)

A. L. Bakke

Iowa State College

Copyright © Copyright 1941 by the Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Bakke, A. L. (1941) "The Use of Tetrachlorethane in the Eradication of the European Bindweed (Abstract)," *Proceedings of the Iowa Academy of Science*: Vol. 48: No. 1 , Article 22.

Available at: <https://scholarworks.uni.edu/pias/vol48/iss1/22>

This Research is brought to you for free and open access by UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

tests indicate that the method may be useful for germinating the pollen of many species, providing an easily performed, stimulating experiment for elementary students.

BOTANY DEPARTMENT,
IOWA STATE COLLEGE,
AMES, IOWA.

THE USE OF TETRACHLORETHANE IN THE ERADICATION OF THE EUROPEAN BINDWEED (ABSTRACT)

A. L. BAKKE

Tetrachlorethane, made by the addition of chlorine to acetylene, has proved to be effective in killing the European bindweed. By making holes 18 inches deep and having the holes 18 inches apart and placing two ounces of tetrachlorethane in each hole, a complete eradication of the European bindweed may be made with one application.

BOTANY DEPARTMENT,
IOWA STATE COLLEGE,
AMES, IOWA.

THE EARLIEST KNOWN EPIPHYTOTIC OF RUST IN IOWA (ABSTRACT)

MARIE A. CORKLE AND I. E. MELHUS

A rust epiphytotic in 1858 apparently has been overlooked in the study of disease prevalence in Iowa. Early Iowa agricultural records and newspapers contain numerous reports of the calamitous wheat and oat failures caused by rust in 1858; yet this rust year has not been reported in any scientific publication from Iowa. Wheat yielded an average of 4.2 bushels per acre, oats 5.4 bushels, while the yields for 1855 were 14.09 and 32.09 bushels per acre, respectively. County agricultural societies reported either total wheat crop failures or very poor yields, with the exception of Van Buren and Woodbury Counties, which reported "medium" and "fair" yields, respectively. In general, oats were reported as a total failure or seriously injured. Throughout the state grain fields were left uncut, not being worth the expense of harvesting. Losses were attributed to rust and wet