

Proceedings of the Iowa Academy of Science

Volume 48 | Annual Issue

Article 125

1941

The Inheritance of Pitted Ear (Abstract)

Karl A. Stiles
Coe College

Copyright ©1941 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Stiles, Karl A. (1941) "The Inheritance of Pitted Ear (Abstract)," *Proceedings of the Iowa Academy of Science*, 48(1), 483-483.

Available at: <https://scholarworks.uni.edu/pias/vol48/iss1/125>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

It was found that the radiation on the first and second days had prevented subsequent development of all cells. Irradiation after the second day inhibited the development only of the cells of the embryo proper.

It is therefore indicated that the yolk and serosa cells undergo a marked increase in their resistance to X-rays after the second day. The yolk and serosa cells are apparently able to complete the process of differentiation following exposure to the radiation. The most evident morphological expression of this differentiation is a marked increase in size, especially of the nucleus. In some of the eggs irradiated on the second day this process of differentiation was so affected that large masses of nuclear material were produced.

DEPARTMENTS OF RADIOLOGY AND ZOOLOGY,
STATE UNIVERSITY OF IOWA,
IOWA CITY, IOWA.

THE INHERITANCE OF PITTED EAR
(ABSTRACT)

KARL A. STILES

Pitted ear is a relatively rare human anomaly defined as a pit in the proximal end of the upper part of the helix. This hereditary malformation has been studied in three different families, not known to be related. Pedigree charts prepared for these families show that one hundred and twenty-seven individuals are involved, of which twenty-four show pitted ear. This physical trait varies greatly throughout the families: the size of the pits ranges from approximately that of a pin head to about that of a match stick; and in depth, from about one-sixteenth to one-half inch. This trait is usually unilateral in its expression, but appears about as often in one ear as in the other; and there are some cases in which it is found in both ears. This character is not inherited as a simple Mendelian dominant, but rather an irregular dominance is suggested.

COE COLLEGE,
CEDAR RAPIDS, IOWA.