

1940

Ecological and Floristic Aspects of Clay and Palo Alto Counties

Ada Hayden
Iowa State College

Let us know how access to this document benefits you

Copyright ©1940 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Hayden, Ada (1940) "Ecological and Floristic Aspects of Clay and Palo Alto Counties," *Proceedings of the Iowa Academy of Science*, 47(1), 155-155.

Available at: <https://scholarworks.uni.edu/pias/vol47/iss1/20>

This Research is brought to you for free and open access by the IAS Journals & Newsletters at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

Offensive Materials Statement: Materials located in UNI ScholarWorks come from a broad range of sources and time periods. Some of these materials may contain offensive stereotypes, ideas, visuals, or language.

ECOLOGICAL AND FLORISTIC ASPECTS OF CLAY
AND PALO ALTO COUNTIES

ADA HAYDEN

A survey of the native flora of Clay and Palo Alto Counties with reference to plants useful to waterfowl includes the plants useful for food, for nesting materials and for cover. The plants are listed systematically as well as referred to associations. The range of distribution for flora of the state has been extended and some plants apparently new to the state have been recorded.

IOWA STATE COLLEGE,
DEPARTMENT OF BOTANY,
AMES, IOWA.

A CHIMERIC TULIP

KARL A. STILES

An unusual yellow Cottage tulip which was collected by Norman Stookey in the spring of 1934 was given to the author. All the petals of this tulip were yellow except one which was one-half yellow and one-half red.

Inasmuch as the bulb from which this tulip developed has not produced another such chimeric flower it appears probable that it is due to a somatic mutation or a somatic non-disjunction of the chromosomes.

BIOLOGY DEPARTMENT,
COE COLLEGE,
CEDAR RAPIDS, IOWA.

A "TWIN-EMBRYO" ABNORMALITY IN MAIZE

JOHN E. SASS AND GEORGE F. SPRAGUE

A mutant strain of corn obtained by ultra violet treatment has twin growing points in the embryo. The flattened emerging coleoptile is a continuous sheath, containing two distinct stem apices,