

1940

## New Synthetic Tools in the Pyridine, Quinoline and Other Nitrogen-Heterocyclic Series

Sydney M. Spatz  
*Iowa State College*

H. Gilman  
*Iowa State College*

Copyright ©1940 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

---

### Recommended Citation

Spatz, Sydney M. and Gilman, H. (1940) "New Synthetic Tools in the Pyridine, Quinoline and Other Nitrogen-Heterocyclic Series," *Proceedings of the Iowa Academy of Science*, 47(1), 262-263.  
Available at: <https://scholarworks.uni.edu/pias/vol47/iss1/51>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact [scholarworks@uni.edu](mailto:scholarworks@uni.edu).

## DYES CONTAINING LONG-CHAINED ALIPHATIC RADICALS

FRED W. HOYT AND H. GILMAN

The investigation of the synthesis and properties of compounds containing long-chained aliphatic radicals has been extended to the preparation of several azo dyes, coupling products of  $\beta$ -naphthol with the appropriately substituted aromatic diazonium chlorides. The dyes prepared include those from a series of three *p*-alkoxyanilines,  $\text{ROC}_6\text{H}_4\text{NH}_2$ , where R is *n*- $\text{C}_{12}\text{H}_{25}$ —, *n*- $\text{C}_{14}\text{H}_{29}$ —, and *n*- $\text{C}_{16}\text{H}_{33}$ —, all deep red in color; and two brighter red compounds from *p*-(N-alkylsulfonamido)-anilines,  $\text{H}_2\text{NC}_6\text{H}_4\text{SO}_2\text{NHR}$ , where R is *n*- $\text{C}_{12}\text{H}_{25}$ — and *n*- $\text{C}_{18}\text{H}_{37}$ —.

These compounds are crystalline solids, soluble in the usual organic solvents and in fats and oils. The melting points of the compounds are such that they might serve as derivatives for the corresponding anilines.

The compounds prepared and their melting points are: 1-(*p*-dodecoxyphenylazo)-2-naphthol, m. 80-82°; 1-(*p*-tetradecoxyphenylazo)-2-naphthol, m. 81.5-83°; 1-(*p*-hexadecoxyphenylazo)-2-naphthol, m. 82-84°; 1-[*p*-(N-dodecylsulfonamido)-phenylazo]-2-naphthol, m. 163-164°; and 1-[*p*-(N-octadecylsulfonamido)-phenylazo]-2-naphthol, m. 158-159.5°.

DEPARTMENT OF CHEMISTRY,  
IOWA STATE COLLEGE,  
AMES, IOWA.

## NEW SYNTHETIC TOOLS IN THE PYRIDINE, QUINOLINE AND OTHER NITROGEN- HETEROCYCLIC SERIES

SYDNEY M. SPATZ AND H. GILMAN

The halogen-metal interconversion reaction now makes possible the preparation of hitherto difficultly or completely inaccessible organolithium compounds from the corresponding bromo- or iodo-nuclear substituted nitrogen heterocycles by means of other organolithium compounds like *n*-propyllithium and *n*-butyllithium:



These organolithium compounds of the nitrogen heterocycles are of a relatively high order of reactivity with other functional groups.

Some of the types reported on in the pyridine series are:

1. 3—pyridyllithium in 70 per cent yield from 3—bromopyridine and *n*—butyllithium,
2. 3—quinollythium in 52 per cent yield from 3—bromopyridine,
3. 5—bromo—3—pyridyllithium in 41 per cent yield from 3, 5—dibromopyridine and slightly more than two equivalents of *n*—butyllithium.

In the carbazole series, some of the organolithium compounds reported are:

1. N—ethyl—2, 8—dilithiocarbazole in 84-91 per cent yield from N—ethyl—2, 8—dibromocarbazole, and in 79-92 per cent yield from the corresponding di-iodo-compound.
2. N—ethyl—2—lithiocarbazole<sup>1</sup> from the corresponding iodo-carbazole in 68 per cent yield.

<sup>1</sup> Studies by Irving Banner.

DEPARTMENT OF CHEMISTRY,  
IOWA STATE COLLEGE,  
AMES, IOWA.

---

## DIRECT SYNTHESSES OF AMIDES FROM LONG-CHAINED ALIPHATIC AMINES

BYRON A. HUNTER, WILLIAM I. HARBER AND H. GILMAN

An examination has been made of direct procedures for the synthesis of amides. Dodecylamine, tetradecylamine, hexadecylamine and octadecylamine have been treated with a variety of aliphatic and aromatic acids. Direct heating of the amine and acid, in open containers and at elevated temperatures, gave quite satisfactory yields of amides.

This direct procedure has been resolved into two stages. First, a mixture of amine and acid, dissolved or suspended in warm petroleum ether, gives promptly and in good yield the corresponding salt. These salts can be used for the characterization of amine and acid. Second, when the salts are heated they are converted to the amides.