

1940

## Pyrolidines from Primary Amines

George H. Coleman  
*State University of Iowa*

Chris C. Schulze  
*State University of Iowa*

Harold A. Hoppens  
*State University of Iowa*

Copyright © Copyright 1940 by the Iowa Academy of Science, Inc.  
Follow this and additional works at: <https://scholarworks.uni.edu/pias>

---

### Recommended Citation

Coleman, George H.; Schulze, Chris C.; and Hoppens, Harold A. (1940) "Pyrolidines from Primary Amines," *Proceedings of the Iowa Academy of Science*: Vol. 47: No. 1 , Article 53.  
Available at: <https://scholarworks.uni.edu/pias/vol47/iss1/53>

This Research is brought to you for free and open access by UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact [scholarworks@uni.edu](mailto:scholarworks@uni.edu).

As an example, *n*-octadecyl amine and acetic acid gives a 24 per cent yield of salt melting at 84.5-85°. Upon heating this salt at 225° for fifteen minutes there is a quantitative conversion to *n*-octadecylacetamide (m.p., 78-78.5°).

DEPARTMENT OF CHEMISTRY,  
IOWA STATE COLLEGE,  
AMES, IOWA.

---

## PYRROLIDINES FROM PRIMARY AMINES

GEORGE H. COLEMAN, CHRIS C. SCHULZE AND  
HAROLD A. HOPPENS

Pyrrolidine has been prepared from *n*-butylamine by heating *N*-chloro-*N*-acyl derivatives of the amine with concentrated sulfuric acid.

*N*-chloro-*N*-butylacetamide underwent ring closure with the formation of pyrrolidine in 50 per cent yields when heated with 95 per cent sulfuric acid for one hour at 130-140°. With 99.5 per cent acid the percentage yield was much lower.

*N*-chloro-*N*-*n*-butyl-*p*-toluenesulfonamide when heated with 95 per cent sulfuric acid for thirty minutes at 140° formed pyrrolidine in 50 per cent yields.

In both cases considerable amounts of *n*-butylamine were also formed.

DEPARTMENT OF CHEMISTRY,  
STATE UNIVERSITY OF IOWA,  
IOWA CITY, IOWA.

---

## BEHAVIOR OF SOME *p*-HYDROXYBENZALDEHYDE DERIVATIVES TOWARD OXIDIZING AGENTS

R. P. PERRY AND L. CHARLES RAIFORD

Benzaldehyde<sup>1</sup> and its alkyl, alkoxy and halogen substitution products are readily oxidized to the corresponding acids. Contact with air often brings about the change.<sup>2</sup> Bücking,<sup>3</sup> and Fittig and

<sup>1</sup> Wöhler and Liebig, *Ann.*, 3,250 (1832).

<sup>2</sup> Bornemann, *Ber.*, 17, 1466 (1884).

<sup>3</sup> Bücking, *Ber.*, 9,529 (1876).