

1940

Melt-Water Volume of Iowa's Last Ice-Sheet

Charles Keyes

Let us know how access to this document benefits you

Copyright ©1940 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Keyes, Charles (1940) "Melt-Water Volume of Iowa's Last Ice-Sheet," *Proceedings of the Iowa Academy of Science*, 47(1), 270-270.

Available at: <https://scholarworks.uni.edu/pias/vol47/iss1/62>

This Research is brought to you for free and open access by the IAS Journals & Newsletters at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

Offensive Materials Statement: Materials located in UNI ScholarWorks come from a broad range of sources and time periods. Some of these materials may contain offensive stereotypes, ideas, visuals, or language.

MELT-WATER VOLUME OF IOWA'S LAST ICE-SHEET

CHARLES KEYES

The old-fashioned notion of a gargantuan ice-cap, so huge as to occupy half of the earth's hemisphere and miles in thickness, was as we now well know, not a creation of the geologist but of the zoologist, a fancy drawn upon to support the theological idea of special creation of life on the globe. It was zoologist Agassiz's last stand against the rapidly growing Darwinian revolution through evolution.

Lately, this all but forgotten concept has been revived to explain submarine terraces, 50 to 300 feet below sea-level, the continental shelf at 600 feet and even deep-sea plains trenched by canyons a mile below sea-level. But an ice-cap is, as we now know, not the gargantuan affair so often pictured, which latter is really a composition of half a dozen different glaciations.

On the other hand, the dimensions of Iowa's last great ice-sheet, about the largest known, are fairly well determined. It probably did not exceed a thousand feet in thickness. At this figure the ice-cap when melted, would return to the ocean water sufficient to raise the sea level only about 2 feet. If, for fancy, the ice were half a mile thick, the sea level would scarcely rise 5 feet. These are maximum figures. The flooded benches under sea, on the continental shelf, will have to be accounted for by some other means than as directrices of melt-waters from a great ice sheet.

DES MOINES, IOWA.

PETROGRAPHY OF THE TERTIARY IGNEOUS ROCKS,
NIGGER HILL DISTRICT, WYOMING-SOUTH DAKOTA

J. ROBERT BERG

The Nigger Hill District, located on the Wyoming-South Dakota border in the northwestern Black Hills, is an area of Paleozoic sedimentaries, dipping outward from a central core of various alkaline and sub-alkaline Tertiary intrusives, and pre-Cambrian pegmatitic granite and schists.

The Tertiary igneous rocks are of interest for three reasons: (1) they may furnish data that will contribute to the much debated problem of the origin of alkaline rocks; (2) when their structural relations are better understood, it will be possible to interpret more