

Proceedings of the Iowa Academy of Science

Volume 47 | Annual Issue

Article 78

1940

A Heat Vaporization Experiment for Elementary Laboratory

L. T. Earls

Iowa State College

Let us know how access to this document benefits you

Copyright ©1940 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Earls, L. T. (1940) "A Heat Vaporization Experiment for Elementary Laboratory," *Proceedings of the Iowa Academy of Science*, 47(1), 287-287.

Available at: <https://scholarworks.uni.edu/pias/vol47/iss1/78>

This Research is brought to you for free and open access by the IAS Journals & Newsletters at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

Offensive Materials Statement: Materials located in UNI ScholarWorks come from a broad range of sources and time periods. Some of these materials may contain offensive stereotypes, ideas, visuals, or language.

graph screen leads us to suppose that the high yield (line) at 150 ekv is due to emission of a homogeneous long-range group.

PHYSICS DEPARTMENT,
STATE UNIVERSITY OF IOWA,
IOWA CITY, IOWA.

DISTRIBUTION IN ANGLE OF ALPHA PARTICLES FROM $F^{19} + H^1$

A. ELLETT, W. B. McLEAN, V. J. YOUNG, G. J. PLAIN

The distribution in angle of long range alpha particles from fluorine bombarded by protons has been studied in the range 270 - 440 ekv. The distribution shows a very strong concentration in the forward direction. Intensity as a function of angle in the center of mass system may be represented by the equation

$$I(\Theta) = 1 + .77 \cos \Theta + .17 \cos^2 \Theta$$

for a bombarding energy of 375 ekv. The distribution shows little, if any, energy dependence and in particular is not observably different at 330 ekv bombarding energy.

Targets were prepared by electrolyzing hydrogen fluoride on tantalum and were fairly thin, the apparent half width of the 330 ekv gamma ray line being 40 ekv or less.

DEPARTMENT OF PHYSICS,
STATE UNIVERSITY OF IOWA,
IOWA CITY, IOWA.

A HEAT VAPORIZATION EXPERIMENT FOR ELEMENTARY LABORATORY

L. T. EARLS

An electric heating coil is used in an ordinary pint vacuum bottle to evaporate water under steady state conditions. The system is used on a platform balance; the heat of vaporization is determined from total energy input and total change of weight. Sources of error are discussed and accuracy obtainable in actual use is indicated.

PHYSICS DEPARTMENT,
IOWA STATE COLLEGE,
AMES, IOWA.