

Proceedings of the Iowa Academy of Science

Volume 46 | Annual Issue

Article 130

1939

Blood Studies in Chronic CO Asphyxia

Marian Davis
Iowa State College

Erma Smith
Iowa State College

Let us know how access to this document benefits you

Copyright ©1939 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Davis, Marian and Smith, Erma (1939) "Blood Studies in Chronic CO Asphyxia," *Proceedings of the Iowa Academy of Science*, 46(1), 443-443.

Available at: <https://scholarworks.uni.edu/pias/vol46/iss1/130>

This Research is brought to you for free and open access by the IAS Journals & Newsletters at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

Offensive Materials Statement: Materials located in UNI ScholarWorks come from a broad range of sources and time periods. Some of these materials may contain offensive stereotypes, ideas, visuals, or language.

BLOOD STUDIES IN CHRONIC CO ASPHYXIA

MARIAN DAVIS AND ERMA SMITH

A closed circuit respiration apparatus was devised by means of which albino rats may be exposed to varying percentages of CO-air mixture and maintained in comfort for hourly intervals. The apparatus enclosed 31 liters of air. By trial and error it was found that addition of 0.31 per cent CO to this volume of air made a mixture just sub-lethal for adapted rats placed therein for one hour. Adaptation was brought about by gradually lengthening the daily exposure interval from an initial ten minutes on the first day to one hour in the course of about three weeks.

Blood studies. (a) Co saturation. The saturation of the hemoglobin with carbon monoxide was measured by the method of Sayers and Yant (1). The percentage saturation for different intervals from two minutes to one hour after one hour exposure will be presented. The saturation at death was about 85 per cent, which is the usual lethal saturation for pure CO.

(b) Coagulation time. The coagulation time was measured in gassed and control rats using the capillary method of Mills (2). Coagulation time appears to be unchanged by exposure to CO.

REFERENCES

1. SAYERS, R. R. AND W. P. YANT. U. S. Bureau of Mines Reports of Investigations, Serial No. 2476, 1923.
2. LEVINSON, S. A. AND R. P. MACFATE. Clinical Lab. Diag., 384, 1937.

DEPARTMENT OF ZOOLOGY AND ENTOMOLOGY,
IOWA STATE COLLEGE
AMES, IOWA.

SOME PHASES OF WOUND HEALING IN AN INSECT (MELANOPLUS DIFFERENTIALIS)

GARTH JOHNSON

A study was made of the inflammatory processes and stages of repair following the production of wounds by incision and excision in the fifth abdominal sternite of both nymphs and adults. The following changes may be observed to take place: A clot is formed