

1939

Effects of X-Rays on the Early Development of the Grasshopper

Titus C. Evans
State University of Iowa

Let us know how access to this document benefits you

Copyright ©1939 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Evans, Titus C. (1939) "Effects of X-Rays on the Early Development of the Grasshopper," *Proceedings of the Iowa Academy of Science*, 46(1), 456-456.

Available at: <https://scholarworks.uni.edu/pias/vol46/iss1/141>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

author hopes is simpler than that in "Fresh Water Biology" by Ward and Whipple.

IOWA LAKESIDE LABORATORY, AND
DEPARTMENT OF ZOOLOGY,
STATE UNIVERSITY OF IOWA,
IOWA CITY, IOWA.

EFFECTS OF X-RAYS ON THE EARLY DEVELOPMENT
OF THE GRASSHOPPER

TITUS C. EVANS

Eggs of *Melanoplus differentialis* were irradiated with different dosages of X-rays on the first day of development. Effects were noted on the subsequent increase in number and kinds of cells, weight and rate of oxygen consumption.

DEPARTMENT OF ZOOLOGY,
STATE UNIVERSITY OF IOWA,
IOWA CITY, IOWA.

THE TIME OF EMBRYONIC DETERMINATION OF
SENSORIA AND ANTENNAL COLOR, AND THEIR
RELATION TO THE DETERMINATION OF
WINGS, OCELLI, AND WING MUSCLE
IN APHIDS

KARL A. STILES

Intermediate-winged aphids of the species *Macrosiphum solani-folii* were studied in an effort to determine the time of embryonic segregation of the sensoria and antennal color, and their relation to the determination of wings, ocelli, and wing muscle.

Dark antennal color and increased sensoria of winged aphids are considered to be characters closely correlated in development; for, in general, when there was a darkening of antennae, there was a corresponding increase in the number of sensoria. In practically all cases increased sensoria were correlated with a relatively large amount of nondegenerate wing muscle. It is concluded that embryonic determination of dark antennal color and increased sensoria takes place in a comparatively short period of time as compared with that of wings, ocelli, and wing muscle. The data make