

1938

Some Cytological Details of Ceresan Poisoning in Seedlings

J. E. Sass
Iowa State College

Let us know how access to this document benefits you

Copyright ©1938 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Sass, J. E. (1938) "Some Cytological Details of Ceresan Poisoning in Seedlings," *Proceedings of the Iowa Academy of Science*, 45(1), 99-99.

Available at: <https://scholarworks.uni.edu/pias/vol45/iss1/15>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

THE MICROFOSSILS OF TWO EOCENE COAL
DEPOSITS IN WYOMING

L. R. WILSON

Samples of coal were secured at six inch intervals through two brown coal deposits near Wamsutter, Wyoming. These were macerated and studied microscopically. The coal is of Green River age and contains an abundance of pollens, spores and invertebrate fragments. The plant fossils were studied statistically and stratigraphically and a high degree of correlation between the spectra of the two deposits is evident. Species of the Cycadaceae, Coniferae, Gnetaceae, Myricaceae, Betulaceae, and Tiliaceae have been recognized in the material.

COE COLLEGE,

CEDAR RAPIDS, IOWA.

SOME CYTOLOGICAL DETAILS OF CERESAN POIS-
ONING IN SEEDLINGS

J. E. SASS

Heavy overdoses of Ceresan induces the formation of giant cells in seedlings of corn and small grains. Cell division is inhibited. Nuclear division occurs, with more or less normal pro-phases. Anaphase separation of halves of split chromosomes fails to take place, accompanied by apparent failure of the sprindle mechanism. Cell wall formation may be initiated, but the evidence suggests subsequent dissolution of partial cell walls.

DEPARTMENT OF BOTANY,

IOWA STATE COLLEGE,

AMES, IOWA.

FURTHER STUDIES OF NON-NUCLEAR STRUCTURES
IN THE BASIDIUM

J. E. SASS

Pursuing a further analysis of the so-called Golgi apparatus in the basidium, tests were made with some accepted chondriosome techniques. Typical chondriosomes were found to be present, quite unlike the bodies associated with the Golgi apparatus. It is suggested that the Nebenkern and Golgi material are discrete cell