

1937

Current Views with Reference to the Nature of Liquid Structure

G. W. Stewart
State University of Iowa

Let us know how access to this document benefits you

Copyright ©1937 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Stewart, G. W. (1937) "Current Views with Reference to the Nature of Liquid Structure," *Proceedings of the Iowa Academy of Science*, 44(1), 149-149.

Available at: <https://scholarworks.uni.edu/pias/vol44/iss1/49>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

mental work for this paper was performed at Physics Department of the State University of Iowa.

DEPARTMENT OF PHYSICS,
PARSONS COLLEGE,
FAIRFIELD, IOWA.

CURRENT VIEWS WITH REFERENCE TO THE NATURE OF LIQUID STRUCTURE

G. W. STEWART

This is a report of progress in the study of the structure of liquids, analogous to the crystal structure of solids. Prins has given a mathematical treatment wherein the crystal structure is approximated to an expressed degree of probability. Bernal has made this treatment more general. These efforts advance the subject by giving models capable of mathematical description. Experiments in Raman effect continue to show that the intermolecular forces in the liquid are much like those in the crystal solid. The study of glasses shows their crystallite structure and emphasizes the difference between glass and liquid. These results, as others, emphasize the essential correctness of the cybotactic view, which emphasizes the fluctuating character of the structure, the liquid possessing at any instant minute spots approximating crystallites.

DEPARTMENT OF PHYSICS,
STATE UNIVERSITY OF IOWA,
IOWA CITY, IOWA.

PHYSICAL DEVELOPMENT OF PHOTOGRAPHIC PLATES

GEORGE HIGGINS

Physical development of photographic plates is not as efficient as chemical development. While it produces very fine grain negatives, longer exposure than normal is required and plates must be developed an hour to produce satisfactory printing densities. pH of the developer is important.

DEPARTMENT OF PHYSICS,
IOWA STATE COLLEGE,
AMES, IOWA.