

1936

Effects of Catalysts on the Preparation and Properties of Organometallic Compounds

M. Lichtenwalter
Iowa State College

J. C. Bailie
Iowa State College

A. J. Carter
Iowa State College

Let us know how access to this document benefits you

Copyright ©1936 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Lichtenwalter, M.; Bailie, J. C.; and Carter, A. J. (1936) "Effects of Catalysts on the Preparation and Properties of Organometallic Compounds," *Proceedings of the Iowa Academy of Science*, 43(1), 205-205. Available at: <https://scholarworks.uni.edu/pias/vol43/iss1/49>

This Research is brought to you for free and open access by the IAS Journals & Newsletters at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

Offensive Materials Statement: Materials located in UNI ScholarWorks come from a broad range of sources and time periods. Some of these materials may contain offensive stereotypes, ideas, visuals, or language.

Bromination of methyl dibenzofuran-4-carboxylate gives methyl 2-bromodibenzofuran-6-carboxylate (m.p., 166°-167°); and nitration gives heteronuclear substitution isomers, the nitro group going to the 2-position and to the 3-position to yield methyl 2-nitrodibenzofuran-6-carboxylate (m.p., 205.5°) and methyl 3-nitrodibenzofuran-6-carboxylate (m.p., 158°). Nitration of 4-methoxydibenzofuran gives 1-nitro-4-methoxydibenzofuran.

DEPARTMENT OF CHEMISTRY,
IOWA STATE COLLEGE,
AMES, IOWA.

EFFECTS OF CATALYSTS ON THE PREPARATION AND PROPERTIES OF ORGANOMETALLIC COMPOUNDS

M. LICHTENWALTER, J. C. BAILIE AND A. J. CARTER

It has been shown that the formation of RM compounds can be accelerated photochemically and by the use of varying quantities of selected reagents. Also, the rates of reaction between RM compounds and various reactants can be accelerated by small quantities of catalysts like cuprous chloride. For example, the time required for complete interaction of benzonitrile and phenylmagnesium bromide is reduced significantly by the addition of small amounts of compounds like cuprous chloride. It appears that these particular catalysts function by first forming very unstable organometallic compounds like phenylcopper, in the case mentioned. Then the phenylcopper decomposes to give copper and diphenyl, probably *via* free phenyl radicals which then set up chain reactions.

DEPARTMENT OF CHEMISTRY,
IOWA STATE COLLEGE,
AMES, IOWA.

THE FRIEDEL-CRAFTS REACTION WITH FURANS

M. McCORKLE AND J. A. V. TURCK, JR.

2-Furoic acid, aluminum chloride and substituted benzenes give 6-substituted-1-naphthoic acids. For example, with chlorobenzene the product is 6-chloro-1-naphthoic acid; with anisole, 6-methoxy-1-naphthoic acid; with toluene, 6-methyl-1-naphthoic acid.