

1936

## Hydroquinone Monoacetate

H. S. Olcott  
*State University of Iowa*

Copyright ©1936 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

---

### Recommended Citation

Olcott, H. S. (1936) "Hydroquinone Monoacetate," *Proceedings of the Iowa Academy of Science*, 43(1), 206-206.

Available at: <https://scholarworks.uni.edu/pias/vol43/iss1/51>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact [scholarworks@uni.edu](mailto:scholarworks@uni.edu).

Benzene, aluminum chloride and methyl 2-furoate give methyl  $\alpha$ -naphthoate in 56 per cent yield.

Benzene, aluminum chloride and methyl 2-methyl-3-furoate give methyl 4-phenyl-4, 5-dihydro-2-methyl-3-furoate; and substituted benzenes and other aryl types give related products.

A very wide variety of branched and straight chained alkyl halides react with ethyl 5-bromo-2-furoate in the presence of aluminum chloride to give ethyl 4-*tert.*-butyl-5-bromo-2-furoate. However, *n*-amyl bromide (unlike *n*-C<sub>5</sub>H<sub>11</sub>Cl, *n*-C<sub>5</sub>H<sub>11</sub>I, and the other RX compounds) gives ethyl 5-*tert.*-butyl-2-furoate exclusively, unless a purer grade (resublimed) aluminum chloride is used in which case the ethyl 4-*tert.*-butyl-5-bromo-2-furoate is obtained.

DEPARTMENT OF CHEMISTRY,  
IOWA STATE COLLEGE,  
AMES, IOWA.

---

## HYDROQUINONE MONOACETATE

H. S. OLCOTT

Alkyl monoesters of hydroquinone and pyrocatechol have not hitherto been prepared. Hydroquinone monoacetate has been obtained by the following series of reactions. Hydroquinone plus one molar equivalent of carbobenzoxy chloride gives hydroquinone monobenzyl carbonate. This compound is acetylated, and then hydrogenated in alcohol with Pd or Ni catalyst. Hydroquinone monoacetate, plates or needles from petroleum ether, M. P. 57-59°C., is obtained from the residue.

BIOCHEMICAL LABORATORY,  
STATE UNIVERSITY OF IOWA,  
IOWA CITY, IOWA.

---

## MIXED DIACYL DERIVATIVES OF *O*-AMINOPHENOL CONTAINING AN ACYL DERIVED FROM A SULFONIC ACID

L. CHAS. RAIFORD AND J. REID SHELTON

In previous work in this Laboratory it was found that, in general, only one mixed diacyl derivative can be obtained from *o*-aminophenol, regardless of the order of introduction of these acyls, provided both are derived from carboxylic acids. In one