

1934

## Comparison of X-Ray Diffraction in H<sub>2</sub>O and H<sub>2</sub>/2O

G. W. Stewart  
*State University of Iowa*

*Let us know how access to this document benefits you*

Copyright ©1934 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

---

### Recommended Citation

Stewart, G. W. (1934) "Comparison of X-Ray Diffraction in H<sub>2</sub>O and H<sub>2</sub>/2O," *Proceedings of the Iowa Academy of Science*, 41(1), 251-251.

Available at: <https://scholarworks.uni.edu/pias/vol41/iss1/80>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact [scholarworks@uni.edu](mailto:scholarworks@uni.edu).

phenomenon has already been observed by Herkert K. Ward with cyclohexane in benzene. Just to what extent these phenomena may be said to be representative is not known.

DEPARTMENT OF PHYSICS,  
STATE UNIVERSITY OF IOWA,  
IOWA CITY, IOWA.

---

### COMPARISON OF X-RAY DIFFRACTION IN H<sub>2</sub>O AND H<sub>2</sub>O

G. W. STEWART

It is to be assumed that the forces in a molecule are approximately invariant for an isotopic change in mass. Consequently the x-ray diffraction curves for H<sub>2</sub>O and H<sub>2</sub>O, determined by the structure of the corresponding liquid, may be expected to be nearly alike. Experiment by x-ray diffractions shows this to be the case, for the differences are well within experimental error and are not as much as 1% over the greater part of the diffraction-intensity-angle curves.

DEPARTMENT OF PHYSICS,  
STATE UNIVERSITY OF IOWA,  
IOWA CITY, IOWA.

---

### HYPERFINE STRUCTURE AND THE DEPOLARIZATION OF RESONANCE RADIATION BY A MAGNETIC FIELD

A. ELLETT

The depolarization of Sodium D line resonance radiation resulting from the Larmor precession of the excited atoms has been carefully determined by a photographic method. Results are compared with a former visual determination (Jour. Opt. Soc. 10, 427, 1925). The effect of hyperfine structure and especially of the degeneracy due to the finite breadth of adjacent hyperfine levels is discussed following Breit, Rev. Mod. Phys. 5:91, 1933, especially Section 4, p. 117 ff.

DEPARTMENT OF PHYSICS,  
STATE UNIVERSITY OF IOWA,  
IOWA CITY, IOWA.