

1934

Effect of Additions of Cadmium on Growth of Zinc Crystals

E. P. T. Tyndall
State University of Iowa

Let us know how access to this document benefits you

Copyright ©1934 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Tyndall, E. P. T. (1934) "Effect of Additions of Cadmium on Growth of Zinc Crystals," *Proceedings of the Iowa Academy of Science*, 41(1), 252-252.

Available at: <https://scholarworks.uni.edu/pias/vol41/iss1/83>

This Research is brought to you for free and open access by the IAS Journals & Newsletters at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

Offensive Materials Statement: Materials located in UNI ScholarWorks come from a broad range of sources and time periods. Some of these materials may contain offensive stereotypes, ideas, visuals, or language.

A MECHANICAL METHOD FOR THE ANALYSIS
OF COMPLEX SPECTRA

J. V. ATANASOFF AND A. E. BRANDT

This paper discusses the application of the punched card equipment, now commonly used in statistical and actuarial work, to the analysis of complex spectra into terms. The spectral frequencies are punched on one set of cards and as complements on another. These cards are then passed in pairs through the tabulator which is provided with a summary punch. The cards from the summary punch are sorted and tabulated. The common differences as well as a code number indicating the lines giving rise to these common differences may be easily read from the tabulation sheet. With the aid of certain refinements, the method becomes very rapid and requires only routine attention on the part of the operator.

DEPARTMENT OF MATHEMATICS,
IOWA STATE COLLEGE,
AMES, IOWA.

EFFECT OF ADDITIONS OF CADMIUM ON GROWTH
OF ZINC CRYSTALS

E. P. T. TYNDALL

Using the Czochralski-Gomperz method and various specimens of zinc containing certainly less than 1 per cent Cd., Schilling¹ encountered mosaic crystals and failed to find the region of successful growth of Hoyem and Tyndall.² It has been found, however, that by using zinc containing several tenths of one per cent of Cadmium the mosaics are largely prevented and a region of growth similar to the previous one reappears.

DEPARTMENT OF PHYSICS,
STATE UNIVERSITY OF IOWA,
IOWA CITY, IOWA.

ELECTRICAL RESISTIVITIES OF ZINC SINGLE AND
MOSAIC CRYSTALS

W. J. POPPY

Continuation of the work on electrical resistivity of single crystal zinc completely confirms the value of the ratio of principal

¹ Physics 5: 1, 1934.

² Phys. Rev. 33: 81, 1929.