

1934

Electron collisions in Mercury Vapor - The 9.8 Volt Loss

C. L. Cross
State University of Iowa

John A. Eldridge
State University of Iowa

Let us know how access to this document benefits you

Copyright ©1934 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Cross, C. L. and Eldridge, John A. (1934) "Electron collisions in Mercury Vapor - The 9.8 Volt Loss," *Proceedings of the Iowa Academy of Science*, 41(1), 257-257.

Available at: <https://scholarworks.uni.edu/pias/vol41/iss1/93>

This Research is brought to you for free and open access by the IAS Journals & Newsletters at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

Offensive Materials Statement: Materials located in UNI ScholarWorks come from a broad range of sources and time periods. Some of these materials may contain offensive stereotypes, ideas, visuals, or language.

the same time to accomplish as a certain laboratory experiment. The transfer values of projects versus laboratory experiments have been tested in seven student groups of known intelligence. It has been difficult to find concepts and skills that did not transfer better in the more intelligent groups from project than from experiment in the laboratory. In the groups of lower intelligence the reverse is true. These data have been treated statistically.

DEPARTMENT OF PHYSICS,
STATE UNIVERSITY OF IOWA,
IOWA CITY, IOWA.

ELECTRON COLLISIONS IN MERCURY VAPOR — THE 9.8 VOLT LOSS

C. L. CROSS AND JOHN A. ELDRIDGE

Magnetic analysis of electrons which have passed through mercury vapor has shown the energy losses corresponding to the principle levels of the normal atom. A very prominent energy loss of 11.1 volts (and so greater than the ionization potential) is not accounted for. It is now shown that the energy loss of 9.8 volts is not due to a double 4.9 excitation.

DEPARTMENT OF PHYSICS,
STATE UNIVERSITY OF IOWA,
IOWA CITY, IOWA.

MEAN FREE PATHS OF MOLECULES AND THE WAVE MECHANICS

JOHN A. ELDRIDGE

Direct experimental determination of the mean free paths of hydrogen, nitrogen and oxygen have led to a value of the effective cross section of the molecule between 4 and 5 times greater than that obtained from viscosity measurements. Massey and Mohr have shown that assuming the rigid sphere molecule the wave mechanics leads to a difference in the values; the theoretical ratio, however, is not greater than 2.

DEPARTMENT OF PHYSICS,
STATE UNIVERSITY OF IOWA,
IOWA CITY, IOWA.