

1930

The Effect of Physical Factors on the Development of Woodland and Prairie Associations in Iowa

J. M. Aikman
Iowa State College

Let us know how access to this document benefits you

Copyright ©1930 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Aikman, J. M. (1930) "The Effect of Physical Factors on the Development of Woodland and Prairie Associations in Iowa," *Proceedings of the Iowa Academy of Science*, 37(1), 209-209.

Available at: <https://scholarworks.uni.edu/pias/vol37/iss1/31>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

THE EFFECT OF PHYSICAL FACTORS ON THE
DEVELOPMENT OF WOODLAND AND
PRAIRIE ASSOCIATIONS IN IOWA

J. M. AIKMAN

The physical factors within the transition zone between the typical prairie association of the state and the deciduous forest communities occurring along the Des Moines River have been studied. The conditions for plant growth at the source of the river in northwestern Iowa have also been compared with those near the mouth of the river in southeastern Iowa. The purpose of this paper is to report the effect of the modification of factors on the distribution and grouping of woodland species and on the general reactions of the communities to the varying habitat factors.

IOWA STATE COLLEGE,
AMES, IOWA.

THE ANATOMY OF THE CROWNS AND ROOTS OF
THE ANNUAL AND BIENNIAL WHITE SWEET
CLOVER COMPARED

J. N. MARTIN

In case of the annual, all structures die at the end of first season's growth. In the biennial only the aerial shoot dies at end of first season's growth. The root and the crown of the biennial prepare, during the first season, for the second season's growth which consists chiefly in the development and fruiting of the flowering shoots that are present on the crown but held in check during first season's growth.

The flowering shoots develop early in the seedling stage in the axils of the cotyledons in both annual and biennial, but in the annual they are not permitted to develop beyond a rudimentary stage. The annual has transferred the flowering function to the first year aerial shoot and suppressed the second year aerial shoot.

In preparation for the second season's growth the root and crown of the biennial develop much cortex and other storage tissues characterized by thin cellulose walls. During the second