

1927

The Action of the Alkyl Chlorides in the Wurtz Reaction

Harry F. Lewis

Let us know how access to this document benefits you

Copyright ©1927 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Lewis, Harry F. (1927) "The Action of the Alkyl Chlorides in the Wurtz Reaction," *Proceedings of the Iowa Academy of Science*, 34(1), 222-222.

Available at: <https://scholarworks.uni.edu/pias/vol34/iss1/52>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

o-nitrobenzaldehyde finds no support on the basis of gas evolved when treated with alkylmagnesium halides.

Benzylmagnesium halides, and related RMgX compounds like α -naphthylmethylmagnesium chloride (where the -MgX group is attached to a carbon that is attached in turn to an unsaturated carbon) have been found to give rearrangement products with new compounds like ethyl chlorocarbonate and ethylene oxide. In some cases (with benzylmagnesium chloride) rearrangement takes place to the para position. A study is in progress of the mechanism of such rearrangements.

In connection with the proof of the *non*-addition of RMgX compounds to an ethylenic linkage, it has been shown that cinnyl chloride when treated with magnesium gives largely a Grignard reagent which when treated with carbon dioxide gives methyl atropic acid. The three other cases in the literature that have been offered as apparent proofs for the addition of RMgX compounds to an ethylenic linkage are also being investigated.

IOWA STATE COLLEGE,
AMES, IOWA.

THE ACTION OF THE ALKYL CHLORIDES IN THE WURTZ REACTION

HARRY F. LEWIS

At the last meeting of the Iowa Academy, a report was made on the mechanism of the Wurtz reaction using butyl and isobutyl bromides. The preparation of octane in this manner is costly and an attempt has been made to replace the bromide by the chloride. This report covers such experiments. Either in the presence or absence of ether as a solvent, the reaction is difficult to control. Yields of octane up to 20% based on the butyl chloride have been obtained. An extremely inflammable by-product is produced and it is difficult to complete a preparation without at least one fire. Using octane itself as a solvent, several bad explosions resulted.

STERIC HINDRANCE IN THE BEHAVIOR OF PHENYL ALKYL ETHER AND DERIVATIVES

L. CHAS. RAIFORD AND D. M. BIROSEL

It is known that phenyl alkyl ethers substitute in the phenyl radical less easily than phenol; nevertheless, phenetol will give a