

1929

The Production of Propionic Acid from Pentoses by *Propionibacterium pentosaceum*

C. H. Werkman
Iowa State College

R. M. Hixon
Iowa State College

E. I. Fulmer
Iowa State College

C. H. Rayburn
Iowa State College

Let us know how access to this document benefits you

Copyright ©1929 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Werkman, C. H.; Hixon, R. M.; Fulmer, E. I.; and Rayburn, C. H. (1929) "The Production of Propionic Acid from Pentoses by *Propionibacterium pentosaceum*," *Proceedings of the Iowa Academy of Science*, 36(1), 111-112.

Available at: <https://scholarworks.uni.edu/pias/vol36/iss1/18>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

THE EFFECT OF STEFFEN WASTE ON THE FERMENTATION OF PENTOSANS FROM THE CORN-STALK

E. I. FULMER, C. H. WERKMAN, R. M. HIXON, AND
A. L. WILLIAMS

It was found that Steffen waste furnishes a suitable source of nitrogen, salts, and buffers for the growth of *Aerobacter pectinovorum* on pentosan material prepared from corn-stalks.

PHYSIOLOGICAL BEHAVIOR OF THE PROPIONIC ACID GROUP OF BACTERIA

SARA E. KENDALL AND C. H. WERKMAN

The group constitutes a number of species of bacteria producing large quantities of propionic acid from carbohydrates. The generic diagnosis of the group is: *Propionibacterium*, Orla-Jensen, 1909. Gram positive, non-sporulating non-motile short rods showing marked morphological variation in acid media or when grown under aerobic conditions; normal growth anaerobic. Cultures are catalase positive. Carbohydrates, glucosides and alcohols attacked with the production of propionic acid, acetic acid and CO₂. The species are differentiated on the basis of sugar fermentation, nitrate reduction, pigment production and morphology. A key to the species is given with a description of each.

IOWA STATE COLLEGE,
AMES, IOWA.

THE PRODUCTION OF PROPIONIC ACID FROM PENTOSE BY *PROPIONIBACTERIUM PENTOSACEUM*

C. H. WERKMAN, R. M. HIXON, E. I. FULMER, C. H. RAYBURN

Propionibacterium pentosaceum attacks pentoses (xylose, arabinose) with the production of propionic and acetic acids. The medium employed was as follows:

| | |
|---------------------------------|---------|
| Dried yeast (Harris Lab.) | 10 gms. |
| K ₂ HPO ₄ | 1 gm. |
| CaCO ₃ | 5 gms. |
| Pentose (xylose or arabinose) | 15 gms. |
| Water | 750 cc. |

The medium was brought to a boil, cooled and adjusted to pH 7.1 electrometrically. It was then sterilized for fifteen minutes at 20 lbs. pressure. The inoculum consisted of 25 cc. of a 10 day culture of *P. pentosaceum* in Sherman's sodium lactate medium. Incubation was at 30°C. for 10 days. The xylose employed was prepared from corn cobs by a method of acid hydrolysis. In addition to the pentoses a number of other fermentable substances were employed to furnish comparative data. Table 1 gives the results

Table 1—Production of Propionic acid by *P. pentosaceum*

| FLASK No. | FERMENTABLE MATERIAL 15 Gms. | PROPI-ONIC ACID ¹ | PER-CENT OF FER-MENTA-ABLE MATERIAL | ACE-TIC ACID ¹ | PER-CENT OF FER-MENTA-ABLE MATERIAL | PROPI-ONIC ACID ² | PER-CENT OF FER-MENTA-ABLE MATERIAL | ACE-TIC ACID ² | PER-CENT OF FER-MENTA-ABLE MATERIAL |
|-----------|------------------------------|------------------------------|-------------------------------------|---------------------------|-------------------------------------|------------------------------|-------------------------------------|---------------------------|-------------------------------------|
| | | Gms. | | Gms. | | Gms. | | Gms. | |
| 1 | Xylose | 3.8 | 25.3 | 1.4 | 9.3 | 4.15 | 27.7 | 1.11 | 7.4 |
| 2 | Arabinose | 4.8 | 32.0 | 1.4 | 9.3 | 5.15 | 34.3 | 1.10 | 7.3 |
| 3 | Sod. lactate | 6.1 | 40.6 | 2.0 | 13.3 | 5.7 | 38.0 | 2.30 | 15.3 |
| 4 | Dextrose | 6.6 | 44.0 | 1.67 | 11.1 | 7.4 | 49.3 | 0.99 | 6.6 |
| 5 | Starch | 1.6 | 10.6 | 3.8 | 25.3 | 1.4 | 9.3 | 2.70 | 18.0 |
| 6 | Glycerol | 3.8 | 25.3 | 0.06 | 0.4 | 3.6 | 24.0 | 0.24 | 1.6 |

of one experiment. Small quantities of apparently succinic acid were found present in all cases, probably derived from the proteins. Both propionic and acetic acids are produced from the pentoses, xylose and arabinose by *P. pentosaceum*. Only propionic acid (no acetic) is produced from glycerol by this organism in significant quantity.

IOWA STATE COLLEGE,
AMES, IOWA.