

1927

Gastric Motility in Vitamin Deficiency

Erma Anita Smith
Iowa State College

Copyright ©1927 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Smith, Erma Anita (1927) "Gastric Motility in Vitamin Deficiency," *Proceedings of the Iowa Academy of Science*, 34(1), 337-337.

Available at: <https://scholarworks.uni.edu/pias/vol34/iss1/119>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

mal is confined in an air-tight vessel also gave results which indicate that the fish uses the air in respiration.

(Preliminary report.)

STATE UNIVERSITY OF IOWA,
 IOWA CITY, IOWA.

GASTRIC MOTILITY IN VITAMIN DEFICIENCY

ERMA ANITA SMITH

Guinea pigs in vitamin C deficiency and dogs in vitamin B deficiency were studied.

I. Guinea pigs:

The emptying time of the stomach after a standard barium meal was determined by means of the X-ray. The experiments were conducted upon four groups of twelve animals each as follows:

	CONTROLS	CHRONIC SCURVY	ACUTE SCURVY
NUMBER OF ANIMALS	16	12	16
Ration	Basal ration plus 4 to 5 cc. orange juice daily	Basal ration plus 0.5 cc. orange juice daily	Basal ration only
Total number of x-ray observations	84	58	50
Average emptying time in minutes	110	113	116

The observations show that the emptying time of the stomach in experimental scurvy is not significantly different from the emptying time in normal control animals.

II. Dogs:

Dogs were fed beri beri producing diets and after loss of appetite for the diet gastric peristalsis was studied by Carlson's balloon method. Graphic records from the stomach of animals before being placed upon the diet — compared to records from the same animals in a state of anorexia, after 100 days on the vitamin B deficient ration, demonstrate the same type of gastric peristalsis. Spastic paralysis develops after the period of pseudo-anorexia. When in this state a hunger balloon cannot be kept in the stomach long enough to obtain records.

IOWA STATE COLLEGE,
 AMES, IOWA.