

1926

A Simple Derivation of the Equation for Mean Free Path

George E. Davis
Iowa State College

Let us know how access to this document benefits you

Copyright ©1926 Iowa Academy of Science, Inc.

Follow this and additional works at: <https://scholarworks.uni.edu/pias>

Recommended Citation

Davis, George E. (1926) "A Simple Derivation of the Equation for Mean Free Path," *Proceedings of the Iowa Academy of Science*, 33(1), 250-250.

Available at: <https://scholarworks.uni.edu/pias/vol33/iss1/70>

This Research is brought to you for free and open access by the Iowa Academy of Science at UNI ScholarWorks. It has been accepted for inclusion in Proceedings of the Iowa Academy of Science by an authorized editor of UNI ScholarWorks. For more information, please contact scholarworks@uni.edu.

A SIMPLE DERIVATION OF THE EQUATION FOR
MEAN FREE PATH

GEORGE E. DAVIS

(ABSTRACT)

According to simple kinetic theory, in which all the gas molecules but one are considered stationary, the mean free path L , is given by $L = \frac{1}{N \pi \sigma^2}$, where N is the number of molecules per cc. and σ is the molecular diameter. Clausius, considering all molecules to be moving with the mean relative velocity, finds $L = \frac{3}{4N \pi \sigma^2}$. Again, Maxwell, applying his law of velocity distribution, obtains the result $L = \frac{1}{\sqrt{2}N \pi \sigma^2}$. This value found by Maxwell is the one most generally accepted. If we consider it to be correct, then the value from simple kinetic theory is 41% too large, while that found by Clausius is 6% too large.

An approximate value of L , much more nearly correct than the first one given, may be derived very simply by considering the gas molecules to be vibrating in sheets. The result obtained is $L = \frac{1}{4 N \sigma^2}$. This value is 11% higher than that found by Maxwell, as compared to 41% for the first value given above. The mathematical difficulties encountered by Clausius and Maxwell in deriving their formulas are entirely avoided in this development.

IOWA STATE COLLEGE,
AMES, IOWA.

A NEW ANALYSIS OF THE ACTION OF AMSLER'S
POLAR PLANIMETER

GEORGE E. DAVIS

(ABSTRACT)

The theory of Amsler's polar planimeter, as commonly given, leads to the expression $A = L h$, where A is the area circumscribed, L is the length of the tracer arm, and h is the net distance of translation of the tracer arm in a direction perpendicular to its length. However, it can be shown that the area may also be given by $A = L^2 \theta$, where θ is the net angle through which the tracer arm has rotated about either of its ends. But as the tracer point passes around the area A , the tracer arm does not, in general,